

Research Article

# Improving Stabilizer Approximation with Quantum Strategy

Fen Zuo<sup>1</sup>

1. Hefei MiQro Era Digital Technology Co. Ltd., China

We introduce a quantum strategy from nonlocal games to improve the stabilizer approximation we proposed previously. The resulting approach turns out to be a qubit-by-qubit gauging procedure for standard stabilizers, which could involve discrete or continuous gauge parameters. We take examples from many-body physics and quantum chemistry to show such a procedure leads to an improvement of the performance.

Corresponding author: Fen Zuo, [zuofen@miqroera.com](mailto:zuofen@miqroera.com)

## I. A brief introduction

The famous toric code<sup>[1]</sup> presents a beautiful connection between a stabilizer code<sup>[2]</sup> and a quantum many-body problem: we could use the stabilizers to engineer a Hamiltonian, such that its ground states constitute exactly the logical subspace, or the stabilized subspace. Reversing the logic, we would like to get the ground states of some Hamiltonian with stabilizers. For realistic Hamiltonians, we wouldn't be so luck to obtain the exact ground states anymore. So we settle for approximate ground states. For Hamiltonians expressed as sums of Pauli terms, a practical approach would be to select an appropriate commuting subset of Pauli terms as the stabilizers, such that the subspace they stabilize would be an approximate ground state. This is the so-called stabilizer approximation that we proposed in<sup>[3]</sup> and developed in<sup>[4][5]</sup>, based on previous studies<sup>[6][7]</sup>. See<sup>[8][9][10]</sup> for related investigations.

While stabilizer approximation performs rather well in most cases, it could go bad when two non-commuting terms possess coefficients of close magnitude. In such a situation choosing either operator as the stabilizer would not be good enough, as discussed in<sup>[8]</sup>. This in fact should have been

anticipated. Stabilizer formalism alone is not adequate to handle the Hamiltonian problem, in the same spirit that Clifford circuits could not provide universal quantum computation<sup>[11][12]</sup>. To do better, we need to introduce more “quantumness” into it.

The development of nonlocal games, especially the Clause–Horne–Shimony–Holt (CHSH) game<sup>[13]</sup>, gives us a clue as how to achieve a quantum advantage. So we start with the CHSH game, and try to borrow the proper quantum strategy from it to improve our approximation.

## II. The CHSH game

We follow the presentation of the CHSH game in Thomas Vidick’s lecture notes<sup>[14]</sup>. Two players in the game are given binary queries  $i$  and  $j$  respectively, and give also binary answers  $a$  and  $b$ . They win if the answers satisfy

$$a_i \oplus b_j = i \wedge j, \quad (1)$$

where  $\oplus$  is logical XOR,  $\wedge$  is logical AND. Assuming the queries are uniformly chosen, we could easily calculate the classical bias and winning probability. To do this, we linearly transform the binary values of  $a$  and  $b$  into spin values  $\{+1, -1\}$ , but still use the same labels to denote them. Then, we have for the bias

$$\beta_c = \max \frac{1}{4}(a_0 b_0 + a_0 b_1 + a_1 b_0 - a_1 b_1) = \frac{1}{2}, \quad (2)$$

and the winning probability

$$\omega_c = \frac{1}{2} + \frac{1}{2}\beta_c = \frac{3}{4}. \quad (3)$$

So classically we have at most a chance of  $\frac{3}{4}$  to win the game.

Quantumly, we could do strictly better. We promote  $\{a_0, a_1\}$  and  $\{b_0, b_1\}$  into

$$A_0 = Z, \quad A_1 = X; \quad B_0 = H = \frac{1}{\sqrt{2}}(X + Z), \quad B_1 = \tilde{H} \equiv \frac{1}{\sqrt{2}}(Z - X). \quad (4)$$

All of them have eigenvalues  $\{+1, -1\}$ . Now we let the two players share an EPR pair:

$$|\Phi\rangle = \frac{|00\rangle + |11\rangle}{\sqrt{2}}, \quad (5)$$

and take the values of the respective measurements  $\{A_i, B_j\}$  as their answers. Therefore, the quantum bias reads

$$\beta_q = \frac{1}{4} \langle \Phi | A_0 \otimes B_0 + A_0 \otimes B_1 + A_1 \otimes B_0 - A_1 \otimes B_1 | \Phi \rangle = \frac{\sqrt{2}}{2}. \quad (6)$$

The winning probability of the quantum strategy is then

$$\omega_q = \frac{1}{2} + \frac{1}{2} \beta_q = \frac{1}{2} + \frac{\sqrt{2}}{4} \approx 0.853, \quad (7)$$

which is indeed greater the classical value. Moreover, it is proved to be the best one.

### III. CHSH vs. stabilizer

The CHSH game has a nice interpretation in terms of the stabilizer formalism, as shown in<sup>[15]</sup>. We slightly modify the notations and presentation in<sup>[15]</sup> to show the significance for the ground state problem.

Say we have a Hamiltonian:

$$H_0 = -X \otimes X - X \otimes Z + Z \otimes X - Z \otimes Z. \quad (8)$$

To approximate its ground state with stabilizer states, we choose a commuting subset of terms, say  $X \otimes X, Z \otimes Z$ , as stabilizers. The corresponding stabilizer state is  $|\Phi\rangle$ , and the corresponding energy, or expectation value, of the Hamiltonian is  $-2$ . Now we make the following transformation:

$$X' = R_y^\dagger\left(\frac{\pi}{4}\right) X R_y\left(\frac{\pi}{4}\right), \quad (9)$$

$$Z' = R_y^\dagger\left(\frac{\pi}{4}\right) Z R_y\left(\frac{\pi}{4}\right), \quad (10)$$

with

$$R_y(\theta) = \cos\left(\frac{\theta}{2}\right) I - i \sin\left(\frac{\theta}{2}\right) Y. \quad (11)$$

Essentially, we are taking the rotation around the  $\hat{y}$  axis by  $\pi/4$ . Explicitly, this gives

$$X' = \frac{X + Z}{\sqrt{2}} = H, \quad Z' = \frac{Z - X}{\sqrt{2}} = \tilde{H}. \quad (12)$$

Then we could rewrite the Hamiltonian as

$$H_0 = -\sqrt{2}(X \otimes X' + Z \otimes Z'). \quad (13)$$

Now if we choose  $\{X \otimes X', Z \otimes Z'\}$  as our new stabilizers, we get a lower energy  $-2\sqrt{2}$ . In fact, this is the lowest eigenvalue of  $H_0$ . The corresponding new stabilizer state is simply<sup>[15]</sup>

$$|\Phi'\rangle = I \otimes R_y^\dagger\left(\frac{\pi}{4}\right) |\Phi\rangle. \quad (14)$$

It is not difficult to check that  $X \otimes X'$  and  $Z \otimes Z'$  indeed stabilize  $|\Phi'\rangle$ . Therefore, the quantum strategy used in the CHSH game helps us improve the stabilizer approximation for the ground state energy. The essential point behind the strategy is that, for different qubits we could use completely different sets of Pauli operators to define the stabilizers. In other words, we could gauge the Pauli operators for each qubit individually, with the single-qubit rotations. I am not sure whether “gauge” is the proper word here to describe the procedure exactly, but I prefer to use it anyhow.

## IV. Gauging stabilizers

### A. Individual gauging

Now we apply the above procedure to some realistic problems, to see whether it is of help more generally. The first example is the Ising model with both longitudinal and transverse magnetic fields<sup>[8]</sup>:

$$H_{Ising} = \sum_{ij} J Z_i \otimes Z_j + \sum_i (g_x X_i + g_z Z_i). \quad (15)$$

When  $g_x \sim g_z \gg J$ , we would have difficulty to select our stabilizers properly. Taking an extremal situation,  $g_x = g_z = 1, J = 0$ , we have

$$H'_{Ising} = \sum_i (X_i + Z_i). \quad (16)$$

At each site, we may choose either  $-Z_i$  or  $-X_i$  as our stabilizer. This gives an energy  $-N$ , with  $N$  the number of sites. Now we gauge the Pauli operators uniformly as before, and get

$$H'_{Ising} = \sqrt{2} \sum_i X'_i. \quad (17)$$

Choosing  $-X'_i$  as the new stabilizer at each site, we get a lower energy  $-\sqrt{2}N$ , which is actually the lowest eigenvalue of  $H'_{Ising}$ . Similar discussion for the generalized toric code model with external magnetic fields is done in<sup>[9]</sup>.

### B. Sequential gauging

The above example looks a little bit trivial. Now we give a slightly involved example. Consider the electronic Hamiltonian of the hydrogen molecule. We transform it into the qubit form with a proper transformation, say the parity transformation<sup>[16]</sup>. In the bounding region, the Hamiltonian looks like this

$$H_{H_2} = 2I_1 \otimes Z_2 - 2Z_1 \otimes I_2. \quad (18)$$

We could easily select the stabilizers as  $-I_1 \otimes Z_2, Z_1 \otimes I_2$ , and get the corresponding state

$$|\Psi_{HF}\rangle = |01\rangle. \quad (19)$$

This is a product/Hartree-Fock (HF) state, and the corresponding energy  $-4$  is exact. In the asymptotic region, the correlation/resonating term dominates, and the Hamiltonian looks like:

$$H'_{H_2} = 2X_1 \otimes X_2 \quad (20)$$

We would get a single stabilizer  $-X_1 \otimes X_2$ , which gives a degenerate state space with the exact energy  $-2$ . In the region in between, the Hamiltonian would be certain combination of the above two, say

$$H''_{H_2} = I_1 \otimes Z_2 - Z_1 \otimes I_2 + 2X_1 \otimes X_2. \quad (21)$$

Choosing either the HF stabilizers  $-I_1 \otimes Z_2, Z_1 \otimes I_2$  or the correlation stabilizer  $-X_1 \otimes X_2$  gives an energy  $-2$ . But this is not so satisfiable, because either choice ignores completely the contributions from the alternative side.

To properly include contributions from both sets of stabilizers, we run the following gauging procedure. First, we gauge  $X_1$  and  $Z_1$  as before, and discard  $Z'_1$ , say, by choosing  $X'_1$  as the first stabilizer. We are left with the reduced Hamiltonian:

$$H''_{H_2} \sim -\frac{1}{\sqrt{2}}I_2 + Z_2 + \sqrt{2}X_2. \quad (22)$$

Now we gauge  $X_2$  and  $Z_2$ , and get

$$H''_{H_2} \sim -\frac{1}{\sqrt{2}}I_2 + \frac{1+\sqrt{2}}{\sqrt{2}}X'_2 + \frac{1-\sqrt{2}}{\sqrt{2}}Z'_2. \quad (23)$$

Choosing  $-X'_2$  as the second stabilizer and discarding  $Z'_2$ , we obtain an energy  $-1 - \sqrt{2} \approx -2.414$ , which is significantly lower than the naive stabilizer value  $-2$ . Still, this is higher than the lowest eigenvalue of  $H''_{H_2}$ , which is  $-2\sqrt{2} \approx -2.828$ .

### C. Continuous gauging

All the previous steps could be generalized by introducing a gauging parameter, which could vary continuously. This could be easily done with a general rotation  $R_y(\theta)$  around the  $\hat{y}$  axis. The gauge transformation would then be given by:

$$X' = R_y^\dagger(\theta)XR_y(\theta), \quad (24)$$

$$Z' = R_y^\dagger(\theta) Z R_y(\theta). \quad (25)$$

When  $\theta = \pi/4$ , we reproduce the previous special transformation as expected.

Now we could repeat the whole calculation for the hydrogen molecule with the general transformation. By properly choosing the gauge parameters, we could further lower the energy from  $-1 - \sqrt{2} \approx -2.414$  to  $-2.5$ , even closer to the exact value  $-2\sqrt{2} \approx -2.828$ .

## V. Summary

In this short note we employ the quantum strategy in the CHSH game to improve the stabilizer approximation for groundstates. The resulting approach turns out to be a gauging procedure for the Pauli operators  $\{Z, X\}$ . As expected, it significantly improves the performance when the original approximation deteriorates. The new stabilizer states in the gauged basis could then provide better initialization for further quantum algorithms<sup>[7][8]</sup>.

Such an approach also inspires a natural question: could it be enhanced to a universal computation model in the Heisenberg picture<sup>[12]</sup>? Perhaps the answer is already hidden in the framework of the so-called ZX-calculus<sup>[17]</sup>, I guess.

## Acknowledgments

I got the idea of gauging stabilizers when I was reading the series of blogposts “A brief introduction to quantum PCP conjecture”, written in Chinese by the blogger “Climber.PI” on his homepage “[Complexity Meets Quantum](#)”. I don’t know the author personally, and I am not even aware of his name. So I would like to take this opportunity to convey my appreciation to him.

## References

1. <sup>△</sup>Kitaev AY (1997). "Fault-tolerant quantum computation by anyons". *Annals Phys.* 303: 2–30. [[arXiv: quant-ph/9707021](#)].
2. <sup>△</sup>Gottesman D (1996). "Class of quantum error-correcting codes saturating the quantum Hamming bound". *Phys. Rev. A* 54 (1996): 1862. *arXiv: [quant-ph/9604038](#)*.
3. <sup>△</sup>Li X, Wang J, Wu C, Zuo F (2022). "Stabilizer Approximation". *arXiv: [2209.09564](#)*.
4. <sup>△</sup>Wang J, Wu C, Zuo F (2023). "Stabilizer Approximation II: From H<sub>2</sub>O To C<sub>6</sub>H<sub>6</sub>". *arXiv. Available from: [arXiv: 2302.11734](#)*.

5. <sup>△</sup>Wu C, Wang J, Zuo F (2023). "Stabilizer Approximation III: Maximum Cut". *arXiv*. Available from: [arXiv: 2303.17215](https://arxiv.org/abs/2303.17215).
6. <sup>△</sup>Shang ZX, Chen MC, Yuan X, Lu CY, Pan JW (2021). "Schrödinger–Heisenberg Variational Quantum Algorithms". *Phys. Rev. Lett.* **131**: 060406. *arXiv*: [2112.07881](https://arxiv.org/abs/2112.07881).
7. <sup>a, b</sup>Gokul Subramanian Ravi, Pranav Gokhale, Yi Ding, William M. Kirby, Kaitlin N. Smith, Jonathan M. Baker, Peter J. Love, Henry Hoffmann, Kenneth R. Brown, Frederic T. Chong (2022). "CAFQA: A classical simulation bootstrap for variational quantum algorithms". *Proceedings of 28th Conference on Architectural Support for Programming Languages and Operating Systems (ASPLOS)*. ACM, New York, NY, USA, 15 pages. doi:[10.1145/3567955.3567958](https://doi.org/10.1145/3567955.3567958). *arXiv*: [2202.12924](https://arxiv.org/abs/2202.12924).
8. <sup>a, b, c, d</sup>Cheng MH, Khosla KE, Self CN, Lin M, Li BX, Medina AC, Kim MS (2022). "Clifford Circuit Initialization for Variational Quantum Algorithms". *arXiv*: [2207.01539](https://arxiv.org/abs/2207.01539).
9. <sup>a, b</sup>Sun J, Cheng L, Zhang S (2024). "Stabilizer ground states for simulating quantum many-body physics: theory, algorithms, and applications". *arXiv*. Available from: [arXiv: 2403.08441](https://arxiv.org/abs/2403.08441).
10. <sup>△</sup>Anand A, Brown KR (2024). "Stabilizer configuration interaction: Finding molecular subspaces with error detection properties". *arXiv*: [2410.21125](https://arxiv.org/abs/2410.21125).
11. <sup>△</sup>Gottesman D (1997). "Stabilizer Codes and Quantum Error Correction". Ph.D. thesis, California Institute of Technology, Pasadena, CA, 1997. *arXiv*: [quant-ph/9705052](https://arxiv.org/abs/quant-ph/9705052).
12. <sup>a, b</sup>Gottesman D (1998). "The Heisenberg Representation of Quantum Computers". <https://arxiv.org/abs/quant-ph/9807006>.
13. <sup>△</sup>Clauser JF, Horne MA, Shimony A, Holt RA (1969). "Proposed Experiment to Test Local Hidden-Variable Theories". *Phys. Rev. Lett.* **23**: 880.
14. <sup>△</sup>Vidick T (2018). "Quantum multiplayer games, testing and rigidity". *Lecture notes of Spring school on Quantum Computation, March 19–22, 2018, University of California, San Diego*.
15. <sup>a, b, c</sup>Zhengfeng Ji (2016). "Classical verification of quantum proofs". *STOC '16: Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*. 885–898.
16. <sup>△</sup>Seeley JT, Richard MJ, Love PJ (2012). "The Bravyi–Kitaev transformation for quantum computation of electronic structure". *J. Chem. Phys.* **137**: 224109 (2012). [[arXiv: 1208.5986](https://arxiv.org/abs/1208.5986)].
17. <sup>△</sup>Coecke B, Kissinger A (2017). *Picturing quantum processes: A first course in quantum theory and diagrammatic reasoning*. Cambridge University Press.

## **Declarations**

**Funding:** No specific funding was received for this work.

**Potential competing interests:** No potential competing interests to declare.