

v1: 14, March 2024

Commentary

On Heated Tobacco Products and the Importance of Science-Based Assessments and Product Classification

Peer-reviewed: 14, March 2024

© The Author(s) 2024. This is an Open Access article under the CC BY 4.0 license.

Qeios, Vol. 6 (2024)
ISSN: 2632-3834

Carrie Wade¹, Markus Nordlund¹, Gizelle Baker¹¹. Philip Morris International, Neuchâtel, Switzerland

Proper assessment of heated tobacco products (HTPs), which are alternatives to cigarettes that heat rather than burn tobacco, is necessary to gain a full understanding of both the potential risks and benefits of these newer products. This includes understanding differences in HTP aerosol compared to cigarette smoke. Recent publications have attempted to characterize HTP emissions in order to provide guidance on relative risk and classification. However, improper representation of studies and misinterpretation of how products are designed and used has led to confusion about the relative risk of HTPs compared to cigarettes. This commentary provides clarification on these important issues with a focus on classification of emissions from the device.

Commentary

It is well accepted that the primary health risks from smoking are not related to nicotine, but instead a consequence of chronic inhalation of the high level of toxicants generated from combustion when tobacco is burned [Benowitz, 2010; RCP 2016]. While the best way for people who smoke to reduce their risk of developing smoking-related disease is to quit tobacco and nicotine altogether, there remain people who do not. Following the principle of tobacco harm reduction [RCP, 2016], over the past two decades, alternative nicotine and tobacco products such as electronic cigarettes and heated tobacco products (HTPs) have been developed that aid and encourage those adult smokers who do not quit to completely switch to using a potentially lower-risk, non-combustible nicotine product. To evaluate the relative harms of these products, it is both essential to categorize them correctly and reasonable to compare their effects to those of the cigarettes that they aim to replace.

Recently, there has been much interest in the composition and classification of the aerosol of HTPs [Uguna and Snape 2022; Bekki *et al.* 2017; Auer *et al.* 2017; Davis, Williams, and Talbot 2019]. Extensive evidence indicates that emissions from HTPs are fundamentally different from those from a combustible cigarette and should be considered an aerosol rather than smoke [FDA 2019; VG Braunschweig 2021; Kärkelä *et al.* 2021; Kärkelä, Tapper, and Kajolinna 2022; Amorós-Pérez *et al.* 2021, 2022; Pratte, Cosandey, and Goujon 2017; Schaller *et al.* 2016; Haziza *et al.* 2020, 2017; Haziza, de La Bourdonnaye, Skiada, *et al.* 2016; Lüdicke *et al.* 2018; Haziza, de La Bourdonnaye, Merlet, *et al.* 2016]. Nevertheless, some authors are arguing that the HTP emissions can be considered smoke [Uguna and Snape 2022; Auer *et al.* 2017] and, further, that emissions from HTPs and cigarettes should be compared on a tobacco-weight rather than a unit-consumption basis [Uguna and Snape 2022]. These recent publications raise concerns about incorrect product classification and risk comparisons based on inaccurate scientific arguments.

The intention of this communication is not to describe in detail the generation and properties of emissions from combustible cigarettes and HTPs, which has been expertly covered by [Sussman *et al.* 2023], but rather to comment on recent issues raised about aerosol categorization and product comparisons with a focus on PMI's HTP, which is marketed as IQOS in some countries.

Differentiating between cigarette smoke and HTP aerosol

To determine whether the emissions of a product should be classified as smoke, it is fundamental (1) to examine the physicochemical processes that generate the emissions; and (2) to characterize the aerosol composition in terms of its chemical and physical characteristics. The recent argument that HTP aerosol can be considered "smoke" is founded on the detection of "smoke" components in the emissions from the HeatStick [Uguna and Snape 2022; Auer *et al.* 2017] without considering their levels or whether these components are produced by chemical reactions, which would indicate creation of a unique substance, or transferred via physical phase-change processes during product operation, which not involve a change at the molecular level.

An analytical study (which has since faced concerns regarding the validity of the methods) [Maeder and Peitsch 2017; Caruso and Polosa 2017; Fujita, Cozzani, and Peitsch 2017; FDA, 2019] of polycyclic aromatic hydrocarbons (PAHs) and volatile organic compounds (VOCs) reported that HTP emissions contain elements from pyrolysis and thermogenic degradation that are the same harmful constituents present in conventional tobacco smoke [Auer *et al.* 2017]. In the case of PAHs, small quantities are present in cured tobacco, and the very low levels observed in HTP emissions have been shown to be proportional to the PAH content of the tobacco materials in the HeatSticks, indicating direct transfer to the aerosol from the tobacco biomass [Goujon *et al.* 2020]. This contrasts with the large amounts of PAHs formed when tobacco burns in a cigarette, and which are the precursors of soot formation [Poget *et al.* 2021; Goujon *et al.* 2020; Pratte, Cosandey, and Goujon 2017; Bentley *et al.* 2020]. Without the production of PAHs at the high concentrations needed to initiate soot particle formation, smoke particles do not form in HTP emissions, and the aerosol produced is liquid-based.

A recent review also highlighted, in addition to PAHs, the presence of black carbon (suggesting soot) and appreciable amounts of "tar" (also called nicotine-free dry particulate matter; NFDPM) as indicators that HTP emissions can be considered smoke [Uguna and Snape 2022]. Although an NFDPM value can be formally calculated for HTPs in the same way as for combustible cigarettes, the compositions of the two types of substance are fundamentally different [Mallock *et al.* 2018]. NFDPM is derived from total particulate matter (TPM), defined as the portion of the smoke/aerosol that is trapped on the filter during standard smoke or aerosol analysis [ISO 4387:2000]. The majority of the TPM of cigarette smoke constitutes smoke-related toxicants, whereas that of HTP aerosol constitutes the humectants needed to generate the aerosol, with far lower levels of tobacco-related toxicants [Bentley, 2020]. Regarding the presence of black carbon in environmental HTP emissions, the cited study [Ruprecht *et al.* 2017] did not detect black carbon that is indicative of soot, but did detect 0.73%–0.79% of the level of 370nm black carbon that is present in cigarette smoke, concluding that the only carbon content in HTP emissions is derived from low levels of organic compounds.

For a full understanding of the vast difference in the physicochemical properties of emissions from combustible cigarettes, HTPs and e-cigarettes, we refer the reader to [Sussman *et al.* 2023]. In brief, tobacco smoke is a highly complex aerosol comprised of solid particles and liquid droplets generated through various processes that are initiated and sustained through ignition and combustion of the cigarette rod; HTP emissions are a complex vapour mixture derived by heating biomass at a temperature below the ignition point [Sussman *et al.* 2023].

Comparative analysis of HTP aerosol with cigarette smoke

In terms of evaluating the relative health implications of tobacco product emissions – irrespective of their classification as smoke or not – the product-use behaviour of the consumers must be factored in. [Uguna and Snape 2022] have stated that the yields of harmful and potentially harmful constituents (HPHCs) in HTP emissions have been underestimated by a factor of 3.2–3.6–

fold because they have been compared with those in cigarette smoke on a per-unit rather than a tobacco-weight basis. However, a comparison based on the weight of tobacco in the products fails to account for the average daily consumption of HTP users versus smokers.

Heated tobacco products were developed as a potentially less harmful alternative to cigarettes. Data from numerous studies show that, when adult smokers switch from cigarettes to HTP, the number of HeatSticks that they use is comparable to the number of cigarettes that they smoked prior to switching [FDA 2019; Haziza *et al.* 2020, 2017; Haziza, de La Bourdonnaye, Skiada, *et al.* 2016]. A clinical study conducted over 90 days has further demonstrated that exposure to nicotine after using one HeatStick is comparable to that after smoking one cigarette [Haziza *et al.* 2020].

Research based on this ratio of use (i.e., 1 cigarette to 1 HeatStick) clearly and reproducibly shows that emissions of HPHCs in HTP aerosol that users are exposed to are significantly lower than those in combustible tobacco smoke [FDA 2019; Kärkelä, Tapper, and Kajolinna 2022; Mallock *et al.* 2018; Schaller *et al.* 2016]. Further, reductions in biomarkers of exposure to HPHCs measured in clinical studies are aligned with the reduction in formation of the respective compounds in the aerosol compared to the levels in cigarette smoke on a per-unit basis [FDA 2019; Znyk, Jurewicz, and Kaleta 2021; Haziza *et al.* 2020; Lüdicke *et al.* 2018; Haziza *et al.* 2017; Haziza, de La Bourdonnaye, Skiada, *et al.* 2016; Haziza, de La Bourdonnaye, Merlet *et al.* 2016].

Conclusions

Based on our scientific assessment of HTPs, together with the results from several independent peer-reviewed studies and independent combustion and aerosol science experts, HTP aerosol is fundamentally different to cigarette smoke and does not fit the definition of smoke, being a liquid-droplet-based aerosol that lacks solid particles. Furthermore, HTP aerosol contains only 532 compounds at levels above 100 ng/HeatStick [Bentley *et al.* 2020], as compared approximately 4800 compounds that have been detected in cigarette smoke at the same threshold [Bentley *et al.* 2020, Rodgman and Green 2003, Rodgman and Perfetti 2013].

When assessing smoke-free products that may reduce the public health burden relative to continued cigarette smoking, it is important that the totality of evidence is taken into consideration, that individual studies are not misrepresented, and that products are evaluated in accordance with how they are used by consumers. Regarding the totality of evidence on THR aerosol, of which only a small portion has been cited herein, the conclusions are clear: it is not smoke, and while not risk-free, it contains only a fraction of the HPHCs found in cigarette smoke.

Statements and Declarations

The authors are employees of Philip Morris International.

References

- Amorós-Pérez A, Cano-Casanova L, del Carmen Román-Martínez M, Lillo-Ródenas MA (2021). "Comparison of Particulate Matter Emission and Soluble Matter Collected from Combustion Cigarettes and Heated Tobacco Products Using a Setup Designed to Simulate Puffing Regimes." *Chemical Engineering Journal Advances* 8: 100144. doi: 10.1016/j.cej.2021.100144.
- Amorós-Pérez A, Cano-Casanova L, del Carmen Román-Martínez M, Lillo-Ródenas MA (2022). "Solid Matter and Soluble Compounds Collected from Cigarette Smoke and Heated Tobacco Product Aerosol Using a Laboratory Designed Puffing Setup." *Environmental Research* 206: 112619. doi: 10.1016/j.envres.2021.112619
- Auer R, Concha-Lozano N, Jacot-Sadowski I, Cornuz J, Berthet A (2017). Heat-Not-Burn Tobacco Cigarettes: Smoke by Any Other Name. *JAMA Internal Medicine* 177: 1050–1052. doi: 10.1001/jamainternmed.2017.1419
- Bekki K, Inaba Y, Uchiyama S, Kunugita N (2017). Comparison of Chemicals in Mainstream Smoke in Heat-Not-Burn Tobacco and Combustion Cigarettes. *Journal of UOEH* 39: 201–207. doi: 10.7888/juoeh.39.201
- Benowitz NL (2010). Nicotine addiction. *New England Journal of Medicine* 362: 2295–2303. doi: 10.1056/NEJMra0809890.
- Bentley MC, Almstetter M, Arndt D, Knorr A, Martin E, Pospisil P, Serge Maeder S (2020). Comprehensive Chemical Characterization of the Aerosol Generated by a Heated Tobacco Product by Untargeted Screening." *Analytical and Bioanalytical Chemistry* 412: 2675–2685. doi: 10.1007/s00216-020-02502-1
- Caruso M, Polosa R (2017). Perplexing Conclusions Concerning Heat-Not-Burn Tobacco Cigarettes." *JAMA Internal Medicine* 177: 1699.

- doi:10.1001/jamainternmed.2017.5843
- Davis B, Williams M, Prue Talbot P (2019). IQOS: Evidence of Pyrolysis and Release of a Toxicant from Plastic. *Tobacco Control* 28: 34–41. doi: 10.1136/tobaccocontrol-2017-054104.
- FDA (2019). PMTA Technical Product Lead Review. <https://www.fda.gov/media/124247/download>
- Fujita, Cozzani, Peitsch MC (2017). Pubpeer Comments on the Manuscript 'Heat-Not-Burn Tobacco Cigarettes: Smoke by Any Other Name.' Pubpeer. 2017. <https://pubpeer.com/publications/EE000F6D5B8732A85FE0366654F95C#2>
- Goujon C, Kleinhans S, Maeder S, Poget L, Schaller J-P (2020). Robustness of HPHC Reduction for THS 2.2 Aerosol Compared with 3R4F Reference Cigarette Smoke Under High Intensity Puffing Conditions. *Contributions to Tobacco & Nicotine Research* 29: 66–83. doi: 10.2478/cttr-2020-0008
- Haziza C, de La Bourdonnaye G, Donelli A, Poux V, Skiada D, Weitkunat R, Baker G, Picavet P, Lüdicke F (2020). Reduction in Exposure to Selected Harmful and Potentially Harmful Constituents Approaching Those Observed Upon Smoking Abstinence in Smokers Switching to the Menthol Tobacco Heating System 2.2 for 3 Months (Part 1)." *Nicotine & Tobacco Research* 22: 539–548. doi: 10.1093/ntr/ntz013.
- Haziza C, de La Bourdonnaye G, Merlet S, Benzimra M, Ancerewicz J, Donelli A, Baker G, Picavet P, Lüdicke F (2016). "Assessment of the Reduction in Levels of Exposure to Harmful and Potentially Harmful Constituents in Japanese Subjects Using a Novel Tobacco Heating System Compared with Conventional Cigarettes and Smoking Abstinence: A Randomized Controlled Study in Confinement." *Regulatory Toxicology and Pharmacology* 81: 489–499. doi: 10.1016/j.yrtph.2016.09.014.
- Haziza C, de La Bourdonnaye G, Skiada D, Ancerewicz J, Baker G, Picavet P, Lüdicke F (2016). "Evaluation of the Tobacco Heating System 2.2. Part 8: 5-Day Randomized Reduced Exposure Clinical Study in Poland." *Regulatory Toxicology and Pharmacology* 81 Suppl 2: S139–S150. doi: 10.1016/j.yrtph.2016.11.003.
- . 2017. Biomarker of Exposure Level Data Set in Smokers Switching from Conventional Cigarettes to Tobacco Heating System 2.2, Continuing Smoking or Abstaining from Smoking for 5 Days. *Data in Brief* 10: 283–293. doi: 10.1016/j.dib.2016.11.047
- Kärkelä T, Ebinger J-C, Tapper U, Robyr O, Jalanti T (2021). "Investigation into the Presence or Absence of Solid Particles Generated from Thermal Processes in the Aerosol from an Electrically Heated Tobacco Product with and without Filter Elements." *Aerosol and Air Quality Research* 21: 200667. doi: 10.4209/aaqr.200667.
- Kärkelä T, Tapper U, Kajolinna T (2022). Comparison of 3R4F Cigarette Smoke and IQOS Heated Tobacco Product Aerosol Emissions. *Environmental Science and Pollution Research International* 29: 27051–27069. doi: 10.1007/s11356-021-18032-x
- Lüdicke F, Picavet P, Baker G, Haziza C, Poux V, Lama N, Weitkunat R (2018). "Effects of Switching to the Tobacco Heating System 2.2 Menthol, Smoking Abstinence, or Continued Cigarette Smoking on Biomarkers of Exposure: A Randomized, Controlled, Open-Label, Multicenter Study in Sequential Confinement and Ambulatory Settings (Part 1)." *Nicotine & Tobacco Research* 20: 161–172. doi: 10.1093/ntr/ntw287.
- Maeder S, Peitsch MC (2017). Perplexing Conclusions Concerning Heat-Not-Burn Tobacco Cigarettes. *JAMA Internal Medicine* 177: 1698–1699. doi: 10.1001/jamainternmed.2017.5840
- Mallock N, Böss L, Burk R, Danziger M, Welsch T, Hahn H, Trieu H-L *et al.* 2018. Levels of Selected Analytes in the Emissions of 'Heat Not Burn' Tobacco Products That Are Relevant to Assess Human Health Risks. *Archives of Toxicology* 92: 2145–2149. doi: 10.1007/s00204-018-2215-y.
- McNeill A, Brose LS, Calder R, Bauld L, Robson D (2018). *Evidence Review of E-Cigarettes and Heated Tobacco Products 2018. A Report Commissioned by Public Health England.* London: Public Health England. https://assets.publishing.service.gov.uk/media/5a981c6740f0b67aa27253cc/cigarettes_and_heated_tobacco_products_2018.pdf
- Poget L, Goujon C, Kleinhans S, Maeder S, Schaller J-P (2021). Robustness of HPHC Reduction in THS 2.2 Aerosol Relative to 3R4F Reference Cigarette Smoke under Extreme Climatic Conditions. *Contributions to Tobacco & Nicotine Research* 30: 109–126. doi: 10.2478/cttr-2021-0008
- Pratte P, Cosandey S, Goujon C (2017). Investigation of Solid Particles in the Mainstream Aerosol of the Tobacco Heating System THS2.2 and Mainstream Smoke of a 3R4F Reference Cigarette. *Human & Experimental Toxicology* 36: 1115–1120. doi: 10.1177/0960327116681653
- Rodgman A, Green CR (2003). Toxic chemicals in cigarette mainstream smoke—hazard and hoopla. *Beitr Tabakforschung/Int Contrib Tob Res.* 2003;20(8):481–545.

- Rodgman A, Perfetti TA (2013). *Chemical Components of Tobacco and Tobacco Smoke, Second Edition*. Taylor & Francis Group.
- Ruprecht AA, De Marco C, Saffari A, Pozzi P, Mazza R, Veronese C, Angellotti G, et al. (2017). Environmental Pollution and Emission Factors of Electronic Cigarettes, Heat-Not-Burn Tobacco Products, and Conventional Cigarettes. *Aerosol Science and Technology* 51: 674–684. doi: [10.1080/02786826.2017.1300231](https://doi.org/10.1080/02786826.2017.1300231)
- Schaller J-P, Keller D, Poget L, Pratte P, Kaelin E, McHugh D, Cudazzo G et al. (2016). Evaluation of the Tobacco Heating System 2.2. Part 2: Chemical Composition, Genotoxicity, Cytotoxicity, and Physical Properties of the Aerosol. *Regulatory Toxicology and Pharmacology* 81 Suppl 2: S27–S47. doi: [10.1016/j.yrtph.2016.10.001](https://doi.org/10.1016/j.yrtph.2016.10.001)
- Sussman RA, Sipala F, Emma R, Ronsisvalle S (2023). Aerosol Emissions from Heated Tobacco Products: A Review Focusing on Carbonyls, Analytical Methods, and Experimental Quality. *Toxics* 11: 947. doi: [10.3390/toxics11120947](https://doi.org/10.3390/toxics11120947)
- Tobacco Advisory Group (2016). *Nicotine without smoke: tobacco harm reduction*. Royal College of Physicians: London. www.rcplondon.ac.uk/sites/default/files/media/Documents/Nicotinewithouts
- Uguna CN, Snape CE (2022). “Should IQOS Emissions Be Considered as Smoke and Harmful to Health? A Review of the Chemical Evidence.” *ACS Omega* 7: 22111–22124. doi: [10.1021/acsomega.2c01527](https://doi.org/10.1021/acsomega.2c01527)
- [VG Braunschweig, 23.09.2021 – 4. A 23/19 – Rauchlos; Tabakerzeugnisse; Verbrennungsprozess | Niedersächsisches Vorschrifteninformationssystem \(NI-VORIS\) \(wolterskluwer-online.de\)](https://www.wolterskluwer-online.de)
- Bundesgerichtshof [BGH] [Federal Court of Justice] Apr. 7, 2003, 154 Entscheidungen des Bundesgerichtshofes in Zivilsachen [BGHZ] 370, 371 (Ger.).
- Znyk M, Jurewicz J, Kaleta D (2021). Exposure to Heated Tobacco Products and Adverse Health Effects, a Systematic Review. *International Journal of Environmental Research and Public Health* 18: 6651. doi: [10.3390/ijerph18126651](https://doi.org/10.3390/ijerph18126651).

Declarations

Funding: The authors are employees of Philip Morris International.

Potential competing interests: The authors are employees of Philip Morris International.