

Long-term beneficial effect of faecal microbiota transplantation on colonisation of multidrug-resistant bacteria and resistome abundance in patients with recurrent *Clostridioides difficile* infection

Sam Nooij, Karuna E. W. Vendrik, Romy D. Zwitterink, Quinten R. Ducarmon, Josbert J. Keller, Ed J. Kuijper, & Elisabeth M. Terveer



Preprint v1

Sep 12, 2023

<https://doi.org/10.32388/1CZVNS>

1 Long-term beneficial effect of faecal microbiota transplantation on
2 colonisation of multidrug-resistant bacteria and resistome abundance in
3 patients with recurrent *Clostridioides difficile* infection

4 Sam Nooij^{1,2}, Karuna E. W. Vendrik^{1,2*}, Romy D. Zwitterink^{1*}, Quinten R. Ducarmon¹, Josbert J. Keller^{2,3},
5 Ed J. Kuijper^{1,2}, & Elisabeth M. Terveer^{1,2}, on behalf of the Netherlands Donor Feces Bank study group

6 ¹Center for Microbiome Analyses and Therapeutics, department of Medical Microbiology, Leiden
7 University Medical Center, Leiden, the Netherlands

8 ²Netherlands Donor Feces Bank, Leiden University Medical Center, Leiden, the Netherlands

9 ³Haaglanden Medical Center, department of gastroenterology, The Hague, the Netherlands

10 *: current affiliation Centre for Infectious Disease Control, Netherlands Institute for Public Health and
11 the Environment, Bilthoven, the Netherlands

12 Abstract

13 Multidrug-resistant (MDR) bacteria are a growing global threat, especially in healthcare facilities.
14 Faecal microbiota transplantation (FMT) is an effective prevention strategy for recurrences of
15 *Clostridioides difficile* infections (rCDI) and can also be useful for other microbiota-related diseases.
16 We study the effect of FMT in patients with rCDI on colonisation with MDR bacteria and antibiotic
17 resistance genes (ARG) on the short (3 weeks) and long term (1-3 years), combining culture methods
18 and faecal metagenomics. Based on MDR culture (n=87 patients), we see a decrease in the
19 colonisation rate of MDR bacteria after FMT (20/87 before FMT = 23%, 10/87 three weeks after FMT
20 = 11.5%). Metagenomic sequencing (n=63 patients) shows a reduction in relative abundances of
21 ARGs in faeces, while the number of different resistance genes in patients remained higher
22 compared to healthy donors (n=11 donors). Furthermore, plasmid predictions in metagenomic data
23 indicate that rCDI patients harboured increased levels of resistance plasmids, which appear
24 unaffected by FMT. In the long-term (n=22 patients), the recipients' resistomes became more donor-
25 like, suggesting that microbiota restoration continues after 3 weeks post-FMT. Taken together, we
26 hypothesise that FMT restores the gut microbiota to a composition that is similar to healthy donors,
27 and potential pathogens are either lost or forced to very low abundances. This process, however,
28 does not end in the days following FMT. It may take months for the gut microbiome to re-establish a
29 balanced state. Even though a reservoir of resistance genes remains, FMT may lead to a more stable
30 and resilient microbiota composition.

31 Introduction

32 The discovery of antibiotics altered the natural course of infectious diseases and saved millions of
33 lives. Antibiotics might be the most significant development in modern medicine, but there are
34 important trade-offs to their use. Antibiotic resistant bacteria have emerged that are unaffected by
35 standard therapies, which threatens effective prevention and treatment of infections. Antibiotic
36 resistance is now considered a major threat to public health [1, 2]. Besides, broad spectrum
37 antibiotic therapy disrupts the human microbiota, paradoxically resulting in an increased
38 susceptibility to infections, for example by *Clostridioides difficile* [3-5].

39 *C. difficile* can asymptotically reside in the gut but thrives in an antibiotic-affected microbiota. *C.*
40 *difficile* causes an infection (CDI) varying from self-limiting and mild diarrhoea to life-threatening
41 pseudomembranous colitis. The disruption of the gut microbiota is essential in maintaining the
42 recurrent nature of CDI, which is supported by the observation that replenishing the gut microbiota
43 by faecal microbiota transplantation (FMT) results in prompt resolution of CDI recurrence (rCDI) [6,
44 7]. It is thought that FMT restores the gut microbiota diversity after antibiotic treatment, thus
45 preventing outgrowth of *C. difficile* spores [8], and possibly decreasing the risk of other infections as
46 well. FMT has been mentioned in treatment guidelines for rCDI for years [9-11] and rCDI is currently
47 the only disease that is routinely treated with FMT.

48 A gut microbiota disrupted by antibiotics is also more susceptible to colonisation with multidrug-
49 resistant (MDR) bacteria [12], which in turn increases the risk of infection in critically ill patients [13].
50 A prominent and problematic group of MDR bacteria are extended spectrum beta-lactamase-
51 producing (ESBL) Enterobacterales. Most infections with ESBL-producing Enterobacterales have high
52 morbidity and mortality and are preceded by intestinal colonisation [14-16]. Hence, the prevention
53 and eradication of ESBL-producing Enterobacterales from the intestinal tract is of global interest.
54 Spontaneous decolonisation depends on comorbidities and type of species [17, 18], and innovative
55 strategies to promote decolonisation of MDR bacteria are desired. So far, there is no recommended
56 decolonisation method [19]. However, Millan et al., found that FMT in patients with rCDI decreased
57 the number and diversity of antimicrobial resistance genes in their faeces [20]. This observation was
58 followed by various case reports of patients colonised with ESBL-producing Enterobacterales who
59 were successfully treated with FMT [21-30]. Only a single, underpowered RCT has been conducted
60 (n=39 patients) to assess decolonisation of MDR Enterobacterales by treatment with oral non-
61 absorbable antibiotics and FMT [31]. No statistically significant advantage of FMT was found,
62 although colonisation rates were slightly lower in FMT-treated patients compared to untreated

63 control patients. Subsequently, questions were raised about the efficacy of FMT against MDR
64 bacteria and experiments were suggested to further assess this [32].

65 To further explore the effects of FMT in rCDI patients on antibiotic resistance of the gut microbiota,
66 we assess colonisation with MDR bacteria with both culture and metagenomics. We pay special
67 attention to the resistome (collection of all antibiotic resistance genes (ARG) present). Additionally,
68 we study the long-term effects on the microbiota up to three years after FMT.

69 Methods

70 In this cohort study, we use stool samples of rCDI patients treated with FMT provided by the
71 Netherlands Donor Feces Bank (NDFB, Leiden, the Netherlands) to assess the presence of MDR
72 bacteria and the resistome. The NDFB uses standardised procedures for the collection, screening,
73 preparation and storage of donor faecal suspensions and treatment and follow-up of rCDI patients
74 as described previously [33, 34]. In short, patients are first treated with antibiotics against *C. difficile*
75 for at least four days until 24 hours before FMT. The day before FMT, patients receive a bowel
76 lavage with macrogol solution [7]. Pre-FMT samples are collected during or after antibiotic
77 treatment and before bowel lavage. Approximately three weeks after FMT a short-term post-FMT
78 sample is requested. Pre- and short-term post-FMT stool samples of rCDI patients and their
79 corresponding donors were collected between May 2016 and March 2021. Additionally, in February
80 2021 we contacted 53 patients to obtain clinical information and request a long-term follow-up
81 (LTFU), or long-term post-FMT, stool sample. (~2 years after FMT.) Clinical data, including recurrence
82 of CDI after FMT, were recorded for further investigation. Stool samples were stored at -80°C until
83 DNA extraction for metagenomics sequencing or stored in an end concentration of 10% glycerol until
84 MDR culture testing. Based on availability, stool samples from the cohort were included for culture
85 and/or sequencing.

86 Written informed consent was obtained from all patients and donors for use of their faecal samples
87 and follow-up data. Ethical approval was granted for the protocols and practice of the NDFB by the
88 local medical ethics committee at the Leiden University Medical Center (reference P15.145, and
89 long-term follow-up: B21.49).

90 *Definition of multidrug-resistant bacteria*

91 Definitions and testing methods were used as described previously [35]. Multidrug-resistant (MDR)
92 bacteria were defined according to the definitions of the Dutch Working Group on Infection
93 Prevention [36]. This includes ESBL-producing Enterobacterales; Enterobacterales and *Acinetobacter*
94 spp. that are resistant to both fluoroquinolones and an aminoglycoside or produce carbapenemases;

95 *Pseudomonas aeruginosa* that produces carbapenemase or is resistant to at least three of the
96 following antibiotic classes or agents: fluoroquinolones, aminoglycosides, ceftazidime or piperacillin,
97 and carbapenems; co-trimoxazole resistant *Stenotrophomonas maltophilia*; penicillin and
98 vancomycin-resistant *Enterococcus faecium* (VRE); or methicillin-resistant *Staphylococcus aureus*
99 (MRSA).

100 *Culture and antimicrobial susceptibility testing of multidrug-resistant bacteria*

101 To identify MDR bacteria in stool, an inoculating loop was used to scrape 10 µL faeces from frozen
102 faeces aliquots (containing 10% glycerol). The faeces was enriched in 15 mL of tryptic soy broth and
103 incubated for 18h at 35°C prior to plating on ChromID ESBL, ChromID OXA-48 agar, MacConkey
104 tobramycin (8 mg/L) plus ciprofloxacin (0.5 mg/L) agar and VRE agar (bioMérieux, Marcy l'Etoile,
105 France). For MRSA detection a separate brain heart infusion enrichment broth was used which was
106 supplemented with 2.5 sodium chloride and 10 mg/L colistin sulphate and inoculation on MRSA-ID
107 agar plate. All suspected MDR colonies were identified by matrix-assisted laser desorption
108 ionisation-time of flight mass spectrometry (MALDI-TOF) Biotyper (Bruker Daltonik; Bremen,
109 Germany). Antibiotic susceptibility was evaluated by VITEK2 (Card N199, bioMérieux) adhering to
110 the European Committee of Antimicrobial Susceptibility Testing (EUCAST) breakpoints version 11.0
111 [37]. ESBL production was confirmed using the double disk method. Isolates with a meropenem
112 minimum inhibitory concentration > 0.25 mg/L (Etest, bioMérieux) were investigated for
113 carbapenemase production with a carbapenem inactivation method (CIM) test and an in-house
114 multiplex PCR to detect *KPC*, *VIM*, *NDM*, *OXA-48* and *IMP* genes. VRE were confirmed by an in-house
115 PCR targeting the *vanA* and *vanB* genes, and MRSA with the BD MAX assay targeting the *MREJ*,
116 *mecA/mecC* and *Nuc* genes (BD, New Jersey, USA). Six known MDR bacteria-positive and seven MDR
117 bacteria-negative defrosted faeces aliquots (also stored in 10% glycerol) of the NDFB donor
118 screening served as positive and negative controls.

119 *Whole-genome sequencing of multidrug-resistant isolates*

120 To assess the antibiotic resistance genotype of MDR isolates and persistence after FMT, 24 out of 30
121 cultured MDR bacteria (16 / 20 from pre-FMT stool samples, 8 / 10 from short-term post-FMT;
122 Table 1) were subjected to whole-genome sequencing. Only isolates of which DNA was available in
123 January 2021 were included for sequencing. DNA was isolated using the QIASymphony DSP
124 Virus/Pathogen Midi Kit (Qiagen, Hilden, Germany) and sent to GenomeScan B.V. (Leiden,
125 Netherlands) to sequence on the Illumina NovaSeq6000 platform (Illumina, Inc., San Diego,
126 California, USA) generating 150 bp paired-end reads. (Reads per bacterial isolate: 780k [258k-1.64M])

127 (median [range].) The raw sequencing reads were cleared of human-derived reads by mapping to
128 the GRCh38 genome [38] using bowtie2 (version 2.4.2, option '--very-sensitive-local') [39] and
129 samtools (version 1.11) [40] before adapter and low-complexity read removal and quality-trimming
130 using fastp (version 0.20.1, parameters '--cut_right --cut_window_size 4 --cut_mean_quality 20 -l 50
131 --detect_adapter_for_pe -y') [41]. High-quality reads were assembled using SPAdes (version 3.15.2,
132 option '--isolate') [42]. All scaffolds were screened for antibiotic resistance genes using ABRicate
133 (version 0.8.13, <https://github.com/tseemann/abricate>) with both the CARD (from 25 March 2021)
134 [43] and ResFinder (from 25 March 2021) [44] databases, only retaining hits of full-length genes
135 (100% coverage) with at least 97% identity. These cut-offs were used to keep the method consistent
136 with and comparable to the resistome analyses (see below). Furthermore, assembled genomes were
137 taxonomically classified using GTDB-Tk (version 2.1.0) [45]. These classifications were used to verify
138 or further specify classifications made by MALDI-TOF Biotyper as described above and are used as
139 species identification for sequenced isolates. Sequence data have been deposited in the European
140 Nucleotide Archive (ENA) under project number PRJEB64622.

141 *Shotgun metagenomic sequencing*

142 Samples collected before 2021 were stored and prepared for sequencing as previously described
143 [46]. This resulted in metagenomes of 49 patients pre- and short-term post-FMT and 56 donor
144 samples of 8 donors that have been deposited in the ENA under project number PRJEB44737. An
145 additional 22 sets of patient pre-, short-term and now including long-term post-FMT samples, of
146 which 7 were sequenced earlier, as well as 14 donor samples from 8 donors were sequenced at
147 GenomeScan B.V. (Leiden, Netherlands) using the Illumina NovaSeq6000 platform generating a
148 median of 42.6M 150bp paired-end reads per sample. Raw reads, excluding human-derived reads
149 (see below), have been deposited in the ENA under project number PRJEB64621. DNA was extracted
150 from 100 mg of unprocessed patient and donor faeces using the Quick-DNA Fecal/Soil Microbe
151 Miniprep Kit (ZymoResearch, Irvine, California, USA), with bead beating step on a Precellys 24 tissue
152 homogeniser (Bertin Technologies, Montigny-le-Bretonneux, France) at 5.5 m/s for three times 1
153 min with short intervals, as described previously [47]. Libraries were constructed using the NEBNext
154 Ultra II FS DNA kit and NEBNext Ultra II Ligation kit (New England Biolabs, Ipswich, Massachusetts,
155 USA), producing DNA fragments of approximately 500-700 bp. Besides, control samples were
156 included to verify successful DNA isolation and sequencing. These include blank (water) controls, and
157 ZymoBiomics Community Standard (ZymoResearch). Negative controls returned no sequencing
158 reads, while positive controls contained reads of all species present in the communities.

159 *Metagenomic read processing*

160 Human-derived reads were removed from raw metagenomic reads by mapping reads to the human
161 reference genome (GRCh38, NCBI accession ID GCF_000001405.26) using bowtie2 (version 2.4.2,
162 option '--very-sensitive-local') and samtools (version 1.11). Remaining non-human reads were then
163 processed by fastp (version 0.20.1) to trim low-quality 3'-ends (parameters: '--cut_right --
164 cut_window_size 4 --cut_mean_quality 20'), remove low-complexity sequences (parameter: '-y'),
165 remove remaining adapter sequences (parameter: '--detect_adapter_for_pe') and remove reads
166 shorter than 50 bases (parameter: '-l 50'). The resulting high-quality metagenomic reads were used
167 in read-based taxonomic profiling and assembly-based ARG profiling.

168 *Quantification of multidrug-resistant isolates in metagenomes*

169 To identify and quantify cultured and whole-genome sequenced MDR bacteria, we mapped
170 metagenomic reads derived from the same stool sample to the respective assembled whole genome
171 using BWA-MEM (version 0.7.17) [48]. Mapped reads were counted and coverage was quantified
172 using samtools coverage (version 1.10).

173 *Taxonomic profiling*

174 Taxonomic microbiota profiles were determined using MetaPhlan (version 4.0.3) [49], which maps
175 reads to its custom marker database. Resulting taxonomic profiles quantified as percentages of the
176 total microbiota were imported as R phyloseq object to facilitate visualisation and statistical
177 comparisons [50].

178 *Resistome analysis*

179 ARGs were detected using an assembly-based approach. Quality-trimmed reads were assembled
180 into scaffolds using metaSPAdes (version 3.15.4, default parameters) [51]. Next, resistance genes
181 were identified with ABRicate (version 0.8.13) using both the CARD (from 25 March 2021) and
182 ResFinder (from 25 March 2021) databases, only retaining hits of full-length genes (100% coverage)
183 with at least 97% identity. These criteria were selected based on visual inspection of the BLAST hits
184 to balance high specificity and adequate sensitivity. As a control, we repeated the analyses using a
185 coverage cut-off of 50% to include partial genes, which yielded equivalent results. ARGs were
186 annotated with their respective target antibiotic and antibiotic class using the respective databases'
187 annotation files. Scaffolds are quantified by mapping the metagenomic reads back to the scaffolds
188 using BWA-MEM (version 0.7.17) and samtools (version 1.10). Quantifications were normalised to
189 reads per kilobase per million (RPKM) by dividing the number of reads mapped to each contig by the
190 length of the contig and the number of high-quality reads used for the assembly, multiplied by 1,000
191 * 1,000,000. To annotate scaffolds with additional information, they were taxonomically classified

192 using the Genome Taxonomy Database Toolkit (GTDB-Tk; version 2.1.0) and the Contig Annotation
193 Tool (CAT, version 5.2.3, parameters: '-r 10 -f 0.5', [52] – which uses Prodigal version 2.6.3 [53];
194 DIAMOND version 2.0.6 [54]; and the NCBI BLAST nr database from 7 January 2021,
195 <https://ftp.ncbi.nlm.nih.gov/blast/db/>), using CAT as primary annotation and filling in gaps in
196 classification using the result of GTDB-Tk. The genomic origin of scaffolds with ARGs (chromosome or
197 plasmid) was estimated using viralVerify (version 1.1, option '-p',
198 <https://github.com/ablab/viralVerify>). Finally, all the scaffold annotation data was loaded into R
199 (version 4.0.2; <https://www.R-project.org/>) for further analyses. The source code of these analyses is
200 available online at: <https://doi.org/10.5281/zenodo.8256351>.

201 *Statistical analyses*

202 The colonisation rate of MDR bacteria among patients was compared between pre- and short-term
203 post-FMT and short-term and the long-term post-FMT using McNemar's chi-square test for paired
204 data. Depth of coverage of MDR bacteria in metagenomic data was compared between pre- and
205 post-FMT with a paired t-test on log-transformed coverage values.

206 For comparing taxonomic compositions of metagenomes and resistomes between donors and
207 patients, we selected one value for each donor as representative. For principal component analyses
208 (PCA), we picked the middle sample for each donor based on donation date (number of samples / 2,
209 rounded up). Aitchison distance was used as metric to describe differences between microbiota or
210 resistome compositions, with resistomes we used a pseudocount of 0.001. For comparisons of
211 diversity metrics using boxplots, we selected the median value for each donor. In PCA, donors and
212 patients are compared using PERMANOVA and PERMDISP tests, considering the repeated measures
213 in patients. Diversity metrics (richness, total abundance, Shannon index, and Simpson evenness) are
214 compared between donors and patients using t-tests, or Wilcoxon rank sum tests when values were
215 visually not normally distributed. Abundance values were log-transformed. Within patients, all pre-
216 and short-term post-FMT measures are compared using a paired t-test, while within the subgroup of
217 22 patients of whom we have collected long-term post-FMT samples values are first compared using
218 repeated measures ANOVA. If $p < 0.05$, paired t-tests were used as post-hoc test to determine
219 differences between pre-FMT and long-term post-FMT and between short and long-term post-FMT.

220 To evaluate if antibiotic (vancomycin) treatment duration before FMT influenced the resistome, we
221 compared the pre-treatment duration of patients (n=52) with their resistome richness (number of
222 different ARGs), total abundance, Shannon diversity and Simpson evenness using Spearman
223 correlation.

224 Colonisation rate of patients by Enterobacterales was determined using MetaPhlan4. If the order
225 Enterobacterales had abundance > 0%, we count it as present and otherwise absent. Colonisation
226 rates between timepoints are compared using McNemar's test. Total abundances were compared
227 using repeated measures ANOVA, followed by pairwise t-tests.

228 All statistical tests were done in R version 4.0.2, using the base, rstatix, vegan, and pairwiseAdonis
229 packages. When multiple tests were conducted simultaneously, p-values were adjusted using Holm's
230 method. A p-value below 0.05 was considered significant.

231 *Data availability*

232 Sequencing reads generated for this study are available in the European Nucleotide Archive under
233 project numbers PRJEB64622, PRJEB44737 and PRJEB64621.

234 Results

235 *Donor and patient population characteristics*

236 During the sample collection period the NDFB provided faecal suspensions for 208 FMT treatments
237 of 187 rCDI patients. Eighty-seven pairs of patient stool samples (median age: 73, interquartile range
238 (IQR): 64-81 years, 56 females (64%)) from pre- and short-term post-FMT in 10% glycerol were
239 available for testing for MDR bacteria by culture (

240 Table 1). Sixty-three pairs of raw frozen patient stool samples (median age: 73 years, interquartile
241 range (IQR) 65-81 years; 40 females (63%)) were available for shotgun metagenomic deep
242 sequencing (

243 Table 1). For 42 patients, samples were available for both culture and metagenomics. Twenty-two
244 patients provided a long-term post-FMT sample (median age: 73, IQR 64-78 years; 14 females
245 (64%)). Furthermore, a total of 70 donor stool samples from 11 different donors (median age: 31
246 years, IQR 27-42 years; 6 females (55%)) were included for metagenomics sequencing (

247 Table 1). The resistome analysis includes only complete sample triads and sample tetrads with long-
248 term post-FMT if both pre- and short-term post-FMT samples were available. That is, patients' pre-
249 FMT and short-term post-FMT, and corresponding donor stool sample (63 patients short-term and
250 21 long-term post-FMT, and 52 donor samples from 11 donors). The median sampling times for
251 patients are: 1 day pre-FMT (IQR 1-3 days), 27 days post-FMT (IQR 20-48 days; short-term), and 801
252 days post-FMT (IQR 447-1114 days; long-term).


253 **Table 1. Overview of sample numbers and applied techniques.**

Sample source	Faecal samples tested by culture for MDR bacteria	MDR isolates whole-genome sequenced / total cultured	Faecal samples used in metagenomic sequencing	Samples with both culture and metagenome data
Donor	76 (15 donors)	0 / 0	70 (11 donors)	43 (8 donors)
Pre-FMT	87	16 / 20	63	42
Short-term post-FMT (~3 wks)	87	8 / 10	63	44
Long-term post-FMT (~1-3 yrs)	22	0 / 2	22	22

254 FMT: faecal microbiota transplantation; MDR: multidrug-resistant.

255 *Prevalence of multidrug-resistant bacteria decreases after FMT*

256 Stool samples of 87 patients were selectively cultured to assess carriage of MDR bacteria. Before
257 FMT, 20/87 (23.0%) of patients carried an MDR bacterium (figure 1A, Table 2). Three weeks after
258 FMT, the colonisation rate decreased to 10/87 (11.5%; $p < 0.0001$), of which 7 MDR bacteria were
259 also cultured from stool samples of the same patient before the FMT. In the long-term, the
260 colonisation rate decreased further to 2/22 (9.1%; $p = 0.0008$ compared to short-term post-FMT).
261 Both were ESBL-producing *E. coli* also detected in the short-term post-FMT samples, thereby they
262 appear to be long-term persisters. Within the subgroup of patients that provided long-term samples,
263 colonisation rates and the shift after FMT were similar as in all 87 patients (figure 1B; pre-FMT 5/22
264 = 22.7%, post-FMT 4/22 = 18.2%; $p = 0.009$).



266

267 **Figure 1. Effect of faecal microbiota transplantation on prevalence and abundance of cultured**
 268 **multidrug resistant bacteria.** Stool samples of recurrent *C. difficile* infected (rCDI) patients were
 269 cultured to assess the prevalence of multidrug resistant (MDR) bacteria before and after faecal
 270 microbiota transplantation (FMT). For a subgroup of patients, long-term follow-up samples were
 271 obtained and tested. Cultured isolates were subjected to whole-genome sequencing and
 272 metagenomic sequencing data from the same stool samples were mapped to the assembled
 273 genomes to quantify the bacteria in the metagenomes. A) Prevalence of MDR bacteria in 87 rCDI
 274 patients. Before FMT, 20/87 patients were colonised by a MDR bacterium, after FMT 10/87 were
 275 colonised. The colonisation rate after FMT is significantly lower (McNemar's chi-square, $p < 0.0001$).
 276 B) Colonisation rates in 22 patients of whom long-term follow-up (~1-3 years after FMT) samples
 277 were collected. 5/22 had a MDR bacterium before FMT, 4/22 were colonized 3 weeks after FMT, and
 278 2/22 were still colonized a few years later. In both time intervals, colonisation rates dropped ($p =$
 279 0.01 and $p = 0.005$, respectively). C) To compare assay sensitivity, we mapped metagenomics reads
 280 to the assemblies of whole-genome sequenced cultured MDR isolates. In general, MDR bacteria had
 281 higher abundance in patients before FMT compared to 3 weeks after FMT (t-test, $p = 0.016$). D)
 282 Breadth of coverage and relative abundance of MDR bacteria in metagenomic sequencing data per
 283 species. Abundance is expressed as mean depth of coverage (nucleotides per position) as reported
 284 by samtools coverage. Asterisks indicate statistically significant differences, *: $p < 0.05$; **: $p < 0.01$;
 285 ****: $p < 0.0001$.
 286 MDR: multidrug resistant, FMT: faecal microbiota transplantation, LTFU: long-term follow-up.

287 **Table 2. Overview of cultured multidrug-resistant bacteria with genotype and phenotype.**

Patient	Sample timepoint	Species	Resistance phenotype	Genotype based on WGS	Detected in metagenome
P22	Post-FMT	<i>E. coli</i>	Aminoglycoside, fluoroquinolone, ampC	<i>APH(3')-Ia, APH(6)-Id, APH(3'')-Ib, ANT(2'')-Ia, acrD, ampC, QnrB5, emrR</i>	Yes
P30	Post-FMT	<i>E. coli</i>	Aminoglycoside, fluoroquinolone	<i>acrD, emrR, emrD</i>	Yes
P31	Pre-FMT	<i>E. coli</i>	Aminoglycoside, fluoroquinolone, ceftazidime	<i>acdD, emrR, emrB, ampC</i>	Yes
P33	Pre-FMT	<i>K. pneumoniae*</i>	Aminoglycoside, fluoroquinolone, ESBL	<i>aadA2, aadA16, AAC(3-IId, TEM-1, SHV-119, CTX-M-14</i>	Yes
P33	Post-FMT	<i>K. pneumoniae*</i>	Aminoglycoside, fluoroquinolone, ESBL	<i>aadA2, aadA16, AAC(3-IId, TEM-1, SHV-119, CTX-M-14</i>	Yes
P38	Pre-FMT	<i>E. coli*</i>	Fluoroquinolone, ESBL	<i>CTX-M-27, ermR, emrB</i>	Yes
P38	Post-FMT	<i>E. coli*</i>	Fluoroquinolone, ESBL	<i>CTX-M-27, ermR, emrB</i>	Yes
P39	Pre-FMT	<i>E. coli</i>	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-15, OXA-1, acrD, AAC(3)-Ile, emrA, emrB, emrR</i>	Yes
P44	Pre-FMT	<i>C. freundii*</i>	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-15, OXA-1, AAC(3)-Ile, AAC(6')-Ib-cr, APH(6)-Id, APH(3'')-Ib, QnrB6</i>	Yes
P44	Post-FMT	<i>C. freundii* + E. coli</i>	Aminoglycoside, fluoroquinolone, ESBL + ESBL	<i>CTX-M-15, OXA-1, AAC(3)-Ile, AAC(6')-Ib-cr, APH(6)-Id, APH(3'')-Ib, QnrB17</i>	Yes
P44	LTFU (3yr)	<i>E. coli</i>	ESBL	NA	NA


P51	Pre-FMT	<i>C. freundii</i>	ESBL	<i>CTX-M-9</i>	Yes
P58	Pre-FMT	<i>E. coli</i>	Aminoglycoside, fluoroquinolone	<i>acrD, emrR</i>	Yes
P59	Pre-FMT	<i>E. coli</i> *	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-14, acrD, AAC(3)-Ile, APH(3'')-Ib, APH(6)-Id, emrR</i>	Yes
P59	Post-FMT	<i>E. coli</i> *	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-14, acrD, AAC(3)-Ile, APH(3'')-Ib, APH(6)-Id, emrR</i>	Yes
P59	LTFU (1yr)	<i>E. coli</i> *	Aminoglycoside, fluoroquinolone, ESBL	NA	NA
P64	Pre-FMT	<i>E. coli</i> *	Aminoglycoside, fluoroquinolone	<i>AAC(6')-Ib-cr, emrA, emrB, emrR</i>	NA
P64	Post-FMT	<i>E. coli</i> *	Aminoglycoside, fluoroquinolone	<i>acrD, APH(3'')-Ib, APH(6)-Id, ampC, ampH, emrA, emrB, emrR</i>	NA
P65	Pre-FMT	<i>E. hormaechei_A (cloacae)</i>	Aminoglycoside, fluoroquinolone, ESBL	<i>ACT-27, CTX-M-15, OXA-1, TEM-1, AAC(3)-Ile, APH(6)-Id, APH(3'')-Ib, AAC(6')-Ib-cr, QnrB6</i>	NA
P66	Pre-FMT	<i>M. morgani</i>	ESBL	<i>DHA-18</i>	NA
P67	Pre-FMT	<i>P. mirabilis</i>	ESBL	<i>CTX-M-1</i>	NA
P68	Pre-FMT	<i>P. mirabilis_B (vulgaris/mirabilis)</i>	ESBL	(none)**	NA
P69	Pre-FMT	<i>C. freundii</i> *	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-15, TEM-1, OXA-1, AAC(3)-Ile, APH(3'')-Ib, APH(6')-Id, QnrB6</i>	NA
P69	Post-FMT	<i>C. freundii</i> *	Aminoglycoside, fluoroquinolone, ESBL	<i>CTX-M-15, TEM-1, OXA-1, AAC(3)-Ile, APH(3'')-Ib, APH(6')-Id, QnrB17</i>	NA
P70	Pre-FMT	<i>E. coli</i> *	ESBL	<i>ampC, ampH, SHV-134</i>	NA
P70	Post-FMT	<i>E. coli</i> *	ESBL	<i>ampC, ampH, SHV-134</i>	NA

P71	Pre-FMT	<i>K. pneumoniae</i>	ESBL	NA	NA
P72	Pre-FMT	<i>P. hauseri</i>	ESBL	NA	NA
P73	Pre-FMT	<i>C. freundii</i>	ESBL	NA	NA
P74	Pre-FMT	<i>E. cloacae</i>	ESBL	NA	NA
P75	Pre-FMT	<i>E. cloacae</i> *	ESBL	NA	NA
P75	Post-FMT	<i>E. cloacae</i> *	ESBL	NA	NA

288 *: same species before and after FMT, persistence is likely based on resistance genotype (supplementary figures 1-3) when available. **: no
289 antibiotic resistance genes were detected in the genome sequence data. Species names are listed as in the Genome Taxonomy Database
290 (GTDB), and the alias known by the National Center for Biotechnology Information (NCBI) is given in parentheses when different. When
291 multiple multidrug-resistant bacteria were cultured from the same stool, isolate characteristics are separated by a plus ('+') sign.
292 FMT: faecal microbiota transplantation, LTFU: long-term follow-up, ESBL: extended-spectrum beta-lactamase, NA: data not available (because
293 the isolate and/or the metagenome were not sequenced)

294 *Whole-genome sequencing of multidrug-resistant and comparison with metagenomics*

295 Twenty-four cultured isolates of multidrug-resistant bacteria were subjected to whole-genome
296 sequencing to study the ARGs and possible persistence after FMT. In all but one genome we were
297 able to detect ARGs associated with the resistance phenotype; e.g., ESBL genes in isolates classified
298 as ESBL-producing (table 2; supplementary figures 1-3). Furthermore, we have both cultured an MDR
299 bacterium and sequenced total DNA from 16 patients' stool samples using shotgun metagenomics.
300 We mapped metagenomic reads to the assembled isolate genome to compare assay sensitivity and
301 determine relative abundances in the microbiota. As expected in patients pre-treated with
302 antibiotics, we found that MDR bacteria had higher relative abundances in rCDI patients before FMT
303 than after FMT (figure 1C; $p = 0.006$). We detected near-complete genomes of MDR isolates in the
304 metagenomes, except one *Citrobacter freundii* (43%; figure 1D). We also compared resistance genes
305 detected in the WGS data to those detected in metagenomic data to estimate the sensitivity of
306 metagenomic sequencing compared to culturing. We found the relevant MDR bacterial resistance
307 genes of cultured isolates back in their respective metagenomes (table 2). Besides, metagenomic
308 data from P44 suggested the presence of an ESBL-producing *E. coli* in the pre-FMT sample, while
309 culture only picked it up in the post-FMT faeces. These data suggest that combining bacterial culture
310 with metagenomic sequencing can be used synergistically and provide more detailed results than
311 either method alone.




312
313 **Figure 2. Prevalence and abundance of Enterobacteriales in faecal donors and faecal microbiota**
314 **transplantation recipients.** A) Relative abundances of Enterobacteriales in metagenomes as
315 determined by MetaPhlan4. B) Total abundances of Enterobacteriales in stool donors and rCDI
316 patients treated with FMT sampled one day before (Pre) FMT, three weeks after (Post) FMT and 1-3
317 years after FMT (long-term follow-up, LTFU). Statistically significant differences are indicated by
318 asterisks, *: $p < 0.05$; ****: $p < 0.0001$.
319 FMT: faecal microbiota transplantation, LTFU: long-term follow-up.

320

321 *Effect of FMT on gut microbiota composition and diversity*

322 We used MetaPhlan4 to study the gut microbiota compositions of faeces from healthy donors and
323 FMT recipients before and after FMT. Donors had a stable microbiota, dominated by Firmicutes,
324 Bacteroidetes and Actinobacteria (supplementary figure 4A-B). Enterobacterales were present in
325 26/70 donor stools (37%). In rCDI patients, that underwent anti-CDI treatment prior to FMT (53 x
326 vancomycin, 6 x fidaxomicin, 1 x metronidazole 1 x metronidazole+vancomycin, 2 unknown),
327 Actinobacteria and Bacteroidetes were much less present, while Proteobacteria (mostly *Escherichia*
328 *coli* or *Klebsiella pneumoniae*) were often dominant (>50% abundance in 31/63 patients = 49%).
329 Enterobacterales were present in all pre-FMT patient stools. After FMT, the patients' microbiota
330 were mixed with their donors' and profiles were more donor-like. Enterobacterales prevalence
331 dropped in the weeks after FMT (58/63 = 92%; $p < 0.0001$; figure 3A) and decreased further in the
332 long-term (18/21 = 86%; $o = 0.012$). Abundance of Enterobacterales also decreased shortly after
333 FMT ($p < 0.0001$; figure 3B) and continued lowering in the long-term ($p = 0.025$) to levels no longer
334 different from those seen in donors ($p = 0.09$). We also compared the alpha and beta diversity
335 between species profiles of donor and patient metagenomes to quantify differences (figure 3). PCA
336 of species profiles showed differences between donors and patients (figure 3A; PERMANOVA, $p =$
337 0.001 ; PERMDISP, $p < 0.0001$). Differences in microbiota profiles were most pronounced between
338 donors and rCDI patients before FMT ($p = 0.003$). While still different, the microbiota profiles after
339 FMT were slightly more donor-like ($p = 0.014$ for both short and long-term compared to donors).
340 When comparing taxonomic profiles at the species rank, we see that richness and alpha diversity
341 (Shannon index) were higher in donors than in rCDI patients (figure 3B-C; $p < 0.0001$) and increased
342 dramatically in patients after FMT ($p < 0.0001$) to levels as seen in donors ($p > 0.1$). Richness and
343 Shannon index remained high at the long-term. The Simpson evenness, also known as inverse
344 Simpson index or Simpson's dominance, was not different between donors and patients (figure 3D; p
345 > 0.3). Overall, our data show the expected pattern of lower diversity in rCDI patients, high diversity
346 in FMT donors, and increased diversity in patients after FMT.



347


348 **Figure 3. Comparison of gut microbiota composition and diversity.** Stool samples of healthy donors
 349 and recipients of faecal microbiota transplantation (FMT) were sequenced using shotgun
 350 metagenomics. We used MetaPhlan4 to determine the taxonomic composition of metagenomes
 351 and compared alpha and beta diversity at the species rank between donors and recipients before
 352 and after FMT, and at a long-term follow-up moment, roughly between 1-3 years after FMT. A) Beta
 353 diversity expressed as Aitchison distances in a principal component analysis (PCA). Percentages on
 354 the X- and Y-axis represent the variance explained by the first two components. B-D) Species
 355 richness, Shannon index and Simpson evenness compared between donors and recipients,
 356 respectively. Asterisks indicate statistically significant differences, ****: $p < 0.0001$.
 357 FMT: faecal microbiota transplantation, LTFU: long-term follow-up.

358

359 *FMT decreases resistance gene abundance, not diversity*

360 Using the metagenomic sequencing data of stool samples from 52 donor stools from their respective
 361 11 donors, we determined the resistome of 63 patients before and after FMT and (supplementary
 362 figures 5-6). Next, we quantified differences in resistome composition between donors and patients
 363 using PCA (figure 4A). Donors had similar resistomes and often had the same ARGs for
 364 aminoglycoside, diaminopyrimidine and tetracycline resistance (supplementary figure 5), while rCDI
 365 patients had a very different resistome (PERMANOVA, $p = 0.003$; PERMDISP, $p < 0.0001$), in which

366 different ARGs for beta-lactam and fluoroquinolone resistance as well as multidrug efflux pumps
 367 were prevalent (supplementary figures 5-6). After FMT, a shift in the patients' resistome toward a
 368 donor-like composition is visible, although it remained different from the donors' ($p = 0.003$). At
 369 long-term follow-up (~1-3 years), the resistome appears visually more similar to the donor's, but was
 370 still statistically different ($p = 0.012$). To gain more insight in resistome changes after FMT, we
 371 visualised relative abundances of resistance genes by antibiotic class (supplementary figure 7A), and
 372 distances from each patient resistome to all donor resistomes (supplementary figure 7B). Both
 373 methods suggest that after FMT resistomes of patients become more like the donors'.



374

375 **Figure 4. Overview of resistomes of faeces donors and faecal microbiota transplantation**
376 **recipients.** Following metagenomics sequencing of stool samples, we determined the resistome of
377 faecal donors and faecal microbiota transplantation (FMT) recipients and looked for differences in
378 these groups and between before and after FMT. A) Principal component analysis (PCA) of
379 resistomes, based on Aitchison distances. Percentages on the X- and Y-axis represent the variance
380 explained by the first two components. B-E) Antibiotic gene richness, total abundance, Shannon
381 index and Simpson evenness compared between groups and between recipient timepoints,
382 respectively. Asterisks indicate statistically significant differences, *: $p < 0.05$; **: $p < 0.01$; ***: $p <$
383 0.001 ; ****: $p < 0.0001$.
384 FMT: faecal microbiota transplantation, LTFU: long-term follow-up.

385

386 We find that patients before FMT had more different resistance genes (higher resistome richness) in
387 their faecal metagenomes than donors ($p < 0.0001$; figure 4B). The duration of vancomycin pre-
388 treatment did not significantly influence the resistome (supplementary figure 8). After FMT,
389 resistome richness in patients did not change ($p > 0.1$) and remained higher than in donors (short-
390 term post-FMT: $p < 0.0001$; long-term: $p = 0.0002$). The total abundance of resistance genes was also
391 higher in patients pre-FMT than in donors (figure 4C; $p < 0.0001$), but in contrast to the resistome
392 richness, abundance decreased in patients shortly after FMT ($p = 0.0003$). In the long term, the
393 abundance lowered further ($p = 0.02$), although abundances remained higher than in donors ($p =$
394 0.02). The Shannon index combines richness and abundance and likewise showed a higher resistome
395 diversity in rCDI patients compared to donors, and a decrease after FMT (figure 4D). The Simpson
396 evenness shows no statistical difference between donors and patients (figure 4E; $p > 0.1$), but
397 indicates a decrease of resistome diversity in patients after FMT ($p = 0.017$). In summary, FMT
398 appears to alter the diversity of the resistome in recipients by lowering relative abundances of ARGs.


399 We observed different prevalence and abundance patterns of ARGs from different antibiotic classes
400 (supplementary figures 5-6). To explore this further, we selected classes of which genes were
401 present in both donors and patients and divided them in two groups. One group (beta-lactamase,
402 fluoroquinolone, and multidrug efflux pump) consists of genes that are rare in donors and common
403 and abundant in patients (supplementary figure 9 A-C and G-I). The abundance of genes in this group
404 decreased shortly after FMT, while the resistome richness decreased only in the long-term. The
405 second group (aminoglycoside, diaminopyrimidine and tetracycline) is common in donors
406 (supplementary figure 9 D-F and J-L). Genes from this donor-associated group may have been
407 transferred to the recipients, resulting in greater resistome richness after FMT and their abundance
408 did not decrease after FMT. These results highlight that the effects of FMT on the resistome vary
409 depending on type of antibiotic and the taxa that carry the genes.

410 *Remarkable resistances*

411 We found a number of ESBL genes in our resistome data, also in donor faeces. Furthermore, we
 412 found carbapenamase genes and one colistin resistance gene (mrc-10_1, predicted to be on a
 413 plasmid) only in rCDI patients before FMT. Vancomycin genes were detected by metagenomics in 7
 414 out of 63 patients before FMT (11.1%) and 11 / 63 after FMT (17.5%; supplementary figure 10).
 415 Besides, our cultures picked up an vancomycin-resistant *Enterococcus faecalis* from short-term post-
 416 FMT stool, which is not listed as MDRO. These resistances as well as those predicted to be on
 417 plasmids are discussed in more detail in the supplementary results and supplementary figures 10-12.

418 *Predicted plasmid-mediated antibiotic resistance remains high*

419 Using the plasmid prediction algorithm from viralVerify, we assessed which contigs with resistance
 420 genes were likely to derive from chromosomes and which from plasmids. Most (4,567 / 6,662 or
 421 68.6%) of the resistance genes were predicted to derive from chromosomes and 400 (6%) likely
 422 derived from plasmids. The remaining 1,695 (25.4%) contigs with ARGs could not confidently be
 423 classified to either plasmid or chromosome. Unlike chromosomal resistances, which follow the
 424 general resistome pattern and decrease after FMT (figure 5A-C), we find that the resistome richness,
 425 abundance, and diversity of ARGs derived from plasmids were higher in rCDI patients than in donors
 426 and stayed higher after FMT ($p \leq 0.01$; figure 5D-F). This effect persisted up to three years after the
 427 FMT, suggesting that FMT may not significantly influence plasmid-mediated antibiotic resistance.



428

429 **Figure 5. Resistome comparisons for chromosomal resistance genes and plasmid-associated**
 430 **resistance genes.** To assess possible differences between antibiotic resistance genes encoded on the
 431 chromosome and plasmid-mediated resistance, we predicted the origin of assembled contigs using

432 viralVerify. We then compared the parameters richness (number of resistance genes) total
433 abundance and Shannon index between faecal donors and FMT recipients separately for resistances
434 genes that are predicted to be on chromosomes (A-C) and genes that are predicted to derive from
435 plasmids (D-F). Richness (A,D) is compared using Wilcoxon rank sum tests, while abundance (B,E)
436 and Shannon index (C,F) are compared using t-tests. Statistically significant differences are indicated
437 by asterisks, *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p < 0.0001$.
438 FMT: faecal microbiota transplantation, LTFU: long-term follow-up.

439

440 Discussion

441 Our current study leverages the strengths of validated culture techniques, metagenomic deep
442 sequencing and long-term follow-up samples to provide the most detailed study on the effects of
443 FMT on antibiotic resistance thus far. The FMT recipients had a more diverse and donor-like
444 resistome and microbiota profile after FMT. The 1–3-year follow-up and plasmid predictions provide
445 novel insights in microbiota dynamics and antibiotic resistance after FMT. We find strong
446 correlations between culture and sequencing data, while each has its advantages. The plasmid
447 predictions give insight into which resistances may be more easily transferred between bacteria,
448 posing a risk when potentially pathogenic species are present.

449 We find effects of FMT on MDR bacteria and the resistome of rCDI patients that last for up to 3
450 years. The effect of FMT is more pronounced on the bacterial species composition than on ARGs.
451 Already three weeks after FMT the recipients' taxonomic composition is similar to the donors', while
452 the resistomes remain different even in the long-term. As most patients were pre-treated with
453 vancomycin, Gram-negative bacteria such as Enterobacterales could thrive. Indeed, we see high
454 prevalence and abundance of Enterobacterales before FMT and a significant decrease after FMT.
455 Enterobacterales are well-studied and many ARGs and plasmids have been described in this group,
456 leading to possible biases in reference databases. The high prevalence of Enterobacterales after FMT
457 correlates with the persistently high resistome richness, while the abundance of both is reduced.
458 This raises the hypothesis that FMT after antibiotic treatment primarily balances the bacterial
459 species composition, lowering abundances of MDR bacteria. However, these bacteria are not
460 eradicated and their ARGs remain present in low relative abundances. The outcome could then be
461 that both *C. difficile* and MDR bacteria cause no symptoms after FMT, but as long as they are present
462 in the patient, there is still a risk of infection [13-15].

463 There are vast differences in resistome composition between healthy donors and rCDI patients,
464 which may relate to medical history and use of antibiotics [55]. More generally, it has been reported
465 that anaerobic gut commensals frequently have aminoglycoside and tetracycline resistance genes

466 [56], which we found to be common in donors and patients, although there were differences in the
467 specific genes. These anaerobic and commensal bacteria may be transferred during FMT and cause a
468 shift in the resistome composition. Consequently, the resistome of FMT recipients is complemented
469 with a distinct donor resistome and this leads to different effects of FMT on different antibiotic
470 classes, as was also shown before [57]. Furthermore, we find that most antibiotic resistance genes in
471 the gut microbiota are chromosomally encoded and may not be as easily transferred as plasmids.
472 Therefore, we hypothesise that most antibiotic resistances are bound to their host species.
473 However, we also see that although the microbiota profile is restored after FMT, the resistome of
474 recipients remains different, largely due to persistence of resistance genes and predicted plasmids.
475 This may be associated with age and comorbidities. To further elucidate this discrepancy, techniques
476 are needed that can link ARGs to their host organism, such as Meta-HiC [56]. This should provide
477 clinicians extra information on the possible risk of infection and treatment options.

478 None of the patients in our cohort reported an infection with an MDR bacterium after FMT and we
479 included no control group without FMT. Therefore, we have no data on risk of infections with MDR
480 bacteria post-FMT. In other patient groups it has been found that FMT decreases the risk of infection
481 [58, 59]. In addition, FMT may reduce the number of infections with MDR bacteria even if the
482 patients' guts are not decolonised with MDR bacteria [60]. The hypothesised mechanism is that the
483 gut microbiota is restored by FMT to a balanced state that is resilient to MDR bacteria [58]. This
484 situation may be described as reduced infection susceptibility or infection resistance. Our data give
485 new details on the involved microbiota processes that play a role in this.

486 Microbiota therapies for the management of MDR bacteria are still in an experimental phase. To
487 better assess the possible benefits, we need larger (randomised controlled) trials and multi-omics
488 studies combined with classical microbiological methods that can link ARGs to bacterial taxa, and to
489 the host's gut ecosystem. Additionally, the use of international registries for FMT can help collect
490 long-term data to assess infection risks in different patient populations. And finally, studies with
491 control patients and more diverse patients are needed to explain the resistome differences and
492 obtain more generalisable results. Then we can evaluate the feasibility of FMT to control antibiotic
493 resistance in infection-susceptible patients.

494 Acknowledgements

495 We wish to express our gratitude to Eric K.L. Berssenbrugge and Ingrid M.G.J. Sanders from the
496 Experimental Bacteriology group at the LUMC for culturing the MDR bacteria. We thank Prof. Dr.
497 Hein W. Verspaget of the Netherlands Donor Feces Bank for continuous support and supervision.
498 Also, our thanks go to GenomeScan B.V. for providing the DNA sequencing. Finally, we thank the

499 Experimental Bacteriology group and the Center for Microbiome Analyses and Therapeutics of the
500 LUMC, in particular Dr. Wiep Klaas Smits, for fruitful work discussions.

501 Author contributions

502 SN, KEWV, RDZ, EJK and EMT conceptualised and designed the study. QRD, EJK, JJK and EMT
503 supervised the study. KEWV, JJK, EJK and EMT supervised treatment of patients. KEWV collected
504 clinical and microbiological data and performed analyses. SN performed genomics, metagenomics
505 and statistical analyses and drafted the manuscript. All authors reviewed and edited the manuscript,
506 and approved the final version.

507 References

- 508 1. **The new EU One Health action plan against antimicrobial resistance**
509 [[https://ec.europa.eu/health/sites/health/files/antimicrobial_resistance/docs/amr_2017_su
511 mary-action-plan.pdf](https://ec.europa.eu/health/sites/health/files/antimicrobial_resistance/docs/amr_2017_su
510 mmary-action-plan.pdf)]
- 512 2. Antimicrobial Resistance C: **Global burden of bacterial antimicrobial resistance in 2019: a
513 systematic analysis**. *Lancet* 2022, **399**(10325):629-655.
- 514 3. Blaser MJ: **Antibiotic use and its consequences for the normal microbiome**. *Science* 2016,
515 **352**(6285):544-545.
- 516 4. Goossens H, Ferech M, Vander Stichele R, Elseviers M, Group EP: **Outpatient antibiotic use
517 in Europe and association with resistance: a cross-national database study**. *Lancet* 2005,
518 **365**(9459):579-587.
- 519 5. Karanika S, Karantanos T, Arvanitis M, Grigoras C, Mylonakis E: **Fecal Colonization With
520 Extended-spectrum Beta-lactamase-Producing Enterobacteriaceae and Risk Factors Among
521 Healthy Individuals: A Systematic Review and Metaanalysis**. *Clin Infect Dis* 2016, **63**(3):310-
522 318.
- 523 6. Quraishi MN, Widlak M, Bhala N, Moore D, Price M, Sharma N, Iqbal TH: **Systematic review
524 with meta-analysis: the efficacy of faecal microbiota transplantation for the treatment of
525 recurrent and refractory Clostridium difficile infection**. *Aliment Pharmacol Ther* 2017,
526 **46**(5):479-493.
- 527 7. van Nood E, Vrieze A, Nieuwdorp M, Fuentes S, Zoetendal EG, de Vos WM, Visser CE, Kuisper
528 EJ, Bartelsman JF, Tijssen JG *et al*: **Duodenal infusion of donor feces for recurrent
529 Clostridium difficile**. *N Engl J Med* 2013, **368**(5):407-415.
- 530 8. Khoruts A, Sadowsky MJ: **Understanding the mechanisms of faecal microbiota
531 transplantation**. *Nat Rev Gastroenterol Hepatol* 2016, **13**(9):508-516.
- 532 9. McDonald LC, Gerding DN, Johnson S, Bakken JS, Carroll KC, Coffin SE, Dubberke ER, Garey
533 KW, Gould CV, Kelly C *et al*: **Clinical Practice Guidelines for Clostridium difficile Infection in
534 Adults and Children: 2017 Update by the Infectious Diseases Society of America (IDSA) and
535 Society for Healthcare Epidemiology of America (SHEA)**. *Clin Infect Dis* 2018, **66**(7):987-994.
- 536 10. Ooijevaar RE, van Beurden YH, Terveer EM, Goorhuis A, Bauer MP, Keller JJ, Mulder CJJ,
537 Kuisper EJ: **Update of treatment algorithms for Clostridium difficile infection**. *Clin Microbiol
538 Infect* 2018, **24**(5):452-462.
- 539 11. van Prehn J, Reigadas E, Vogelzang EH, Bouza E, Hristea A, Guery B, Krutova M, Noren T,
540 Allerberger F, Coia JE *et al*: **European Society of Clinical Microbiology and Infectious
541 Diseases: 2021 update on the treatment guidance document for Clostridioides difficile
infection in adults**. *Clin Microbiol Infect* 2021, **27** Suppl 2:S1-S21.

- 542 12. Isles NS, Mu A, Kwong JC, Howden BP, Stinear TP: **Gut microbiome signatures and host**
543 **colonization with multidrug-resistant bacteria.** *Trends in Microbiology* 2022, **30**(9):853-865.
- 544 13. Dickstein Y, Edelman R, Dror T, Hussein K, Bar-Lavie Y, Paul M: **Carbapenem-resistant**
545 **Enterobacteriaceae colonization and infection in critically ill patients: a retrospective**
546 **matched cohort comparison with non-carriers.** *J Hosp Infect* 2016, **94**(1):54-59.
- 547 14. Carlet J: **The gut is the epicentre of antibiotic resistance.** *Antimicrob Resist Infect Control*
548 2012, **1**(1):39.
- 549 15. Gorrie CL, Mirceta M, Wick RR, Judd LM, Wyres KL, Thomson NR, Strugnell RA, Pratt NF,
550 Garlick JS, Watson KM *et al*: **Antimicrobial-Resistant *Klebsiella pneumoniae* Carriage and**
551 **Infection in Specialized Geriatric Care Wards Linked to Acquisition in the Referring**
552 **Hospital.** *Clin Infect Dis* 2018, **67**(2):161-170.
- 553 16. Tillotson GS, Zinner SH: **Burden of antimicrobial resistance in an era of decreasing**
554 **susceptibility.** *Expert Rev Anti Infect Ther* 2017, **15**(7):663-676.
- 555 17. Weterings V, van den Bijllaardt W, Bootsma M, Hendriks Y, Kilsdonk L, Mulders A, Kluytmans
556 J: **Duration of rectal colonization with extended-spectrum beta-lactamase-producing**
557 ***Escherichia coli*: results of an open, dynamic cohort study in Dutch nursing home residents**
558 **(2013-2019).** *Antimicrob Resist Infect Control* 2022, **11**(1):98.
- 559 18. van Weerlee C, van der Vorm ER, Nolles L, Meeuws-van den Ende S, van der Bij AK: **Duration**
560 **of carriage of multidrug resistant Enterobacterales in discharged hospital and general**
561 **practice patients and factors associated with clearance.** *Infect Prev Pract* 2020,
562 **2**(3):100066.
- 563 19. Tacconelli E, Mazzaferri F, de Smet AM, Bragantini D, Eggimann P, Huttner BD, Kuijper EJ,
564 Lucet JC, Mutters NT, Sanguinetti M *et al*: **ESCMID-EUCIC clinical guidelines on**
565 **decolonization of multidrug-resistant Gram-negative bacteria carriers.** *Clin Microbiol Infect*
566 2019, **25**(7):807-817.
- 567 20. Millan B, Park H, Hotte N, Mathieu O, Burguiere P, Tompkins TA, Kao D, Madsen KL: **Fecal**
568 **Microbial Transplants Reduce Antibiotic-resistant Genes in Patients With Recurrent**
569 ***Clostridium difficile* Infection.** *Clin Infect Dis* 2016, **62**(12):1479-1486.
- 570 21. Bilinski J, Grzesiowski P, Sorensen N, Madry K, Muszynski J, Robak K, Wroblewska M,
571 Dzieciatkowski T, Dulny G, Dwilewicz-Trojaczek J *et al*: **Fecal Microbiota Transplantation in**
572 **Patients With Blood Disorders Inhibits Gut Colonization With Antibiotic-Resistant Bacteria:**
573 **Results of a Prospective, Single-Center Study.** *Clin Infect Dis* 2017, **65**(3):364-370.
- 574 22. Crum-Cianflone NF, Sullivan E, Ballon-Landa G: **Fecal microbiota transplantation and**
575 **successful resolution of multidrug-resistant-organism colonization.** *J Clin Microbiol* 2015,
576 **53**(6):1986-1989.
- 577 23. Davido B, Batista R, Michelon H, Lepointeur M, Bouchand F, Lepeule R, Salomon J, Vittecoq
578 D, Duran C, Escaut L *et al*: **Is faecal microbiota transplantation an option to eradicate highly**
579 **drug-resistant enteric bacteria carriage?** *J Hosp Infect* 2017, **95**(4):433-437.
- 580 24. Dinh A, Fessi H, Duran C, Batista R, Michelon H, Bouchand F, Lepeule R, Vittecoq D, Escaut L,
581 Sobhani I *et al*: **Clearance of carbapenem-resistant Enterobacteriaceae vs vancomycin-**
582 **resistant enterococci carriage after faecal microbiota transplant: a prospective**
583 **comparative study.** *J Hosp Infect* 2018, **99**(4):481-486.
- 584 25. Huttner BD, Galperine T, Kapel N, Harbarth S: **'A five-day course of oral antibiotics followed**
585 **by faecal transplantation to eradicate carriage of multidrug-resistant Enterobacteriaceae' -**
586 **Author's reply.** *Clin Microbiol Infect* 2019, **25**(7):914-915.
- 587 26. Lagier JC, Million M, Fournier PE, Brouqui P, Raoult D: **Faecal microbiota transplantation for**
588 **stool decolonization of OXA-48 carbapenemase-producing *Klebsiella pneumoniae*.** *J Hosp*
589 *Infect* 2015, **90**(2):173-174.
- 590 27. Manges AR, Steiner TS, Wright AJ: **Faecal microbiota transplantation for the intestinal**
591 **decolonization of extensively antimicrobial-resistant opportunistic pathogens: a review.**
592 *Infect Dis (Lond)* 2016, **48**(8):587-592.

- 593 28. Singh R, Nieuwdorp M, ten Berge IJ, Bemelman FJ, Geerlings SE: **The potential beneficial**
594 **role of faecal microbiota transplantation in diseases other than Clostridium difficile**
595 **infection.** *Clin Microbiol Infect* 2014, **20**(11):1119-1125.
- 596 29. Singh R, de Groot PF, Geerlings SE, Hodiament CJ, Belzer C, Berge I, de Vos WM, Bemelman
597 FJ, Nieuwdorp M: **Faecal microbiota transplantation against intestinal colonization by**
598 **extended spectrum beta-lactamase producing Enterobacteriaceae: a proof of principle**
599 **study.** *BMC Res Notes* 2018, **11**(1):190.
- 600 30. Stalenhoef JE, Terveer EM, Knetsch CW, Van't Hof PJ, Vlasveld IN, Keller JJ, Visser LG, Kuijper
601 EJ: **Faecal Microbiota Transfer for Multidrug-Resistant Gram-Negatives: A Clinical Success**
602 **Combined With Microbiological Failure.** *Open Forum Infect Dis* 2017, **4**(2):ofx047.
- 603 31. Huttner BD, de Lastours V, Wassenberg M, Maharshak N, Mauris A, Galperine T, Zanichelli V,
604 Kapel N, Bellanger A, Olearo F *et al*: **A 5-day course of oral antibiotics followed by faecal**
605 **transplantation to eradicate carriage of multidrug-resistant Enterobacteriaceae: a**
606 **randomized clinical trial.** *Clin Microbiol Infect* 2019, **25**(7):830-838.
- 607 32. Kuijper EJ, Vendrik KEW, Vehreschild M: **Manipulation of the microbiota to eradicate**
608 **multidrug-resistant Enterobacteriaceae from the human intestinal tract.** *Clin Microbiol*
609 *Infect* 2019, **25**(7):786-789.
- 610 33. Terveer EM, van Beurden YH, Goorhuis A, Seegers J, Bauer MP, van Nood E, Dijkgraaf MGW,
611 Mulder CJJ, Vandenbroucke-Grauls C, Verspaget HW *et al*: **How to: Establish and run a stool**
612 **bank.** *Clin Microbiol Infect* 2017, **23**(12):924-930.
- 613 34. Terveer EM, Vendrik KE, Ooijevaar RE, Lingen EV, Boeije-Koppenol E, Nood EV, Goorhuis A,
614 Bauer MP, van Beurden YH, Dijkgraaf MG *et al*: **Faecal microbiota transplantation for**
615 **Clostridioides difficile infection: Four years' experience of the Netherlands Donor Feces**
616 **Bank.** *United European Gastroenterol J* 2020, **8**(10):1236-1247.
- 617 35. Vendrik KEW, Terveer EM, Kuijper EJ, Nooij S, Boeije-Koppenol E, Sanders I, van Lingen E,
618 Verspaget HW, Berssenbrugge EKL, Keller JJ *et al*: **Periodic screening of donor faeces with a**
619 **quarantine period to prevent transmission of multidrug-resistant organisms during faecal**
620 **microbiota transplantation: a retrospective cohort study.** *Lancet Infect Dis* 2021, **21**(5):711-
621 721.
- 622 36. **Bijzonder resistente micro-organismen (BRMO) (in Dutch)**
623 [<https://www.rivm.nl/sites/default/files/2018-11/130424%20BRMO.pdf>]
- 624 37. **Breakpoint tables for interpretation of MICs and zone diameters. Version 11.0, 2021.**
625 [<http://www.eucast.org>]
- 626 38. **Genome assembly GRCh38**
627 [https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_000001405.26/]
- 628 39. Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat Methods* 2012,
629 **9**(4):357-359.
- 630 40. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
631 Genome Project Data Processing S: **The Sequence Alignment/Map format and SAMtools.**
632 *Bioinformatics* 2009, **25**(16):2078-2079.
- 633 41. Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ preprocessor.**
634 *Bioinformatics* 2018, **34**(17):i884-i890.
- 635 42. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A: **Using SPAdes De Novo**
636 **Assembler.** *Curr Protoc Bioinformatics* 2020, **70**(1):e102.
- 637 43. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA, Dave BM, Pereira S,
638 Sharma AN *et al*: **CARD 2017: expansion and model-centric curation of the comprehensive**
639 **antibiotic resistance database.** *Nucleic Acids Res* 2017, **45**(D1):D566-D573.
- 640 44. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM,
641 Larsen MV: **Identification of acquired antimicrobial resistance genes.** *J Antimicrob*
642 *Chemother* 2012, **67**(11):2640-2644.

- 643 45. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk v2: memory friendly**
644 **classification with the genome taxonomy database.** *Bioinformatics* 2022, **38**(23):5315-5316.
- 645 46. Nooij S, Ducarmon QR, Laros JFJ, Zwittink RD, Norman JM, Smits WK, Verspaget HW, Keller
646 JJ, Terveer EM, Kuijper EJ *et al*: **Fecal Microbiota Transplantation Influences**
647 **Procarcinogenic Escherichia coli in Recipient Recurrent Clostridioides difficile Patients.**
648 *Gastroenterology* 2021, **161**(4):1218-1228 e1215.
- 649 47. Ducarmon QR, Hornung BVH, Geelen AR, Kuijper EJ, Zwittink RD: **Toward Standards in**
650 **Clinical Microbiota Studies: Comparison of Three DNA Extraction Methods and Two**
651 **Bioinformatic Pipelines.** *mSystems* 2020, **5**(1).
- 652 48. Li H: **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.** In:
653 *arXiv: 13033997 [q-bioGN]*. 2013.
- 654 49. Blanco-Miguez A, Beghini F, Cumbo F, McIver LJ, Thompson KN, Zolfo M, Manghi P, Dubois L,
655 Huang KD, Thomas AM *et al*: **Extending and improving metagenomic taxonomic profiling**
656 **with uncharacterized species using MetaPhlAn 4.** *Nat Biotechnol* 2023.
- 657 50. McMurdie PJ, Holmes S: **phyloseq: an R package for reproducible interactive analysis and**
658 **graphics of microbiome census data.** *PLoS One* 2013, **8**(4):e61217.
- 659 51. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA: **metaSPAdes: a new versatile**
660 **metagenomic assembler.** *Genome Res* 2017, **27**(5):824-834.
- 661 52. von Meijenfildt FAB, Arkhipova K, Cambuy DD, Coutinho FH, Dutilh BE: **Robust taxonomic**
662 **classification of uncharted microbial sequences and bins with CAT and BAT.** *Genome Biol*
663 2019, **20**(1):217.
- 664 53. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ: **Prodigal: prokaryotic gene**
665 **recognition and translation initiation site identification.** *BMC Bioinformatics* 2010, **11**:119.
- 666 54. Buchfink B, Reuter K, Drost HG: **Sensitive protein alignments at tree-of-life scale using**
667 **DIAMOND.** *Nat Methods* 2021, **18**(4):366-368.
- 668 55. Fredriksen S, de Warle S, van Baarlen P, Boekhorst J, Wells JM: **Resistome expansion in**
669 **disease-associated human gut microbiomes.** *Microbiome* 2023, **11**(1):166.
- 670 56. McCallum GE, Rossiter AE, Quraishi MN, Iqbal TH, Kuehne SA, Schaik Wv: **Noise reduction**
671 **strategies in metagenomic chromosome confirmation capture to link antibiotic resistance**
672 **genes to microbial hosts.** *bioRxiv* 2022:2022.2011.2005.514866.
- 673 57. Langdon A, Schwartz DJ, Bulow C, Sun X, Hink T, Reske KA, Jones C, Burnham CD, Dubberke
674 ER, Dantas G *et al*: **Microbiota restoration reduces antibiotic-resistant bacteria gut**
675 **colonization in patients with recurrent Clostridioides difficile infection from the open-label**
676 **PUNCH CD study.** *Genome Med* 2021, **13**(1):28.
- 677 58. Ghani R, Mullish BH, Davies FJ, Marchesi JR: **How to adapt an intestinal microbiota**
678 **transplantation programme to reduce the risk of invasive multidrug-resistant infection.**
679 *Clin Microbiol Infect* 2022, **28**(4):502-512.
- 680 59. Tariq R, Pardi DS, Tosh PK, Walker RC, Razonable RR, Khanna S: **Fecal Microbiota**
681 **Transplantation for Recurrent Clostridium difficile Infection Reduces Recurrent Urinary**
682 **Tract Infection Frequency.** *Clin Infect Dis* 2017, **65**(10):1745-1747.
- 683 60. Bilsen MP, Lambregts MMC, van Prehn J, Kuijper EJ: **Faecal microbiota replacement to**
684 **eradicate antimicrobial resistant bacteria in the intestinal tract - a systematic review.** *Curr*
685 *Opin Gastroenterol* 2022, **38**(1):15-25.

686