

A Systematic Review of Antibiotic Use in Humans in Nigeria and Its Potential Contribution to Rising Antimicrobial Resistance

Abiodun Egwuenu¹, Adebola Olayinka, Busayo Olayinka, Yonni Apeji², Olufemi Abayomi, Biodun Ogunniyi, Hamzat Omotayo, Joshua Obasanya, Oladipo Aboderin, Iruka Okeke, Chikwe Ihekweazu

¹ Nigeria Centre for Disease Control

² Ahmadu Bello University

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Introduction: The value of medicines is lost when these resources are not used rationally. Out of 12 developing countries, Nigeria has the third highest percentage of antibiotic prescriptions, at 48%. Antibiotic misuse results in limited efficacy of antibiotics, which can lead to the emergence of antimicrobial resistance. We conducted systematic review to synthesise the evidence on antibiotic use in humans in Nigeria.

Methods: We conducted a systematic review of medicine use behaviour by patients and prescription behaviour by health workers, which were searched for in articles published in English from 2000 to 2017. Data was entered into purpose-built templates. Key quantitative indicators were extracted and summarised as frequencies and proportions, while free-text responses were synthesised.

Results: The systematic review determined that the median prevalence of persons using antibiotics without prescription is 46.7%. The drivers of irrational antibiotic use included poor regulation of medicines and premises, a chaotic medicine distribution system, limited licensed medicine prescribers, over-the-counter (OTC) sales of antibiotics, patients' demand for antibiotics, and access to health insurance.

Discussion: Irrational antibiotic use is widespread in humans and animals. We recommend that the government enforce regulations on antibiotic sales to humans and animals and increase awareness of irrational antibiotic use and AMR in Nigerian communities through a whole-of-society approach. The collated information was used to develop a National Action Plan on AMR in 2017.

Abiodun Egwuenu¹, Adebola Olayinka², Busayo Olayinka³, Yonni Apeji³, Olufemi Abayomi⁴, Biodun Ogunniyi¹, Hamzat Omotayo², Joshua Obasanya⁶, Oladipo Aboderin⁷, Iruka Okeke⁸ and Chikwe Ihekweazu²

¹ Nigeria Centre for Disease Control and Prevention, Federal Capital Territory, Nigeria

² World Health Organization

³ Ahmadu Bello University, Kaduna, Nigeria

⁴ Federal Ministry of Defense, Federal Capital Territory, Nigeria

⁵ Africa Centres for Disease Control and Prevention

⁶ Obafemi Awolowo University Teaching Hospital Complex, Osun, Nigeria

⁷ University of Ibadan, Oyo, Nigeria

Introduction

Since the advent of the first antibiotic (penicillin), 20 new classes of antibiotics have been introduced or approved for use between 1935 and 2012 (Figure 1)^[1]. Concomitantly, there has been an increase in antibiotic use by animal and human populations. Global antibiotic consumption grew by over 30% in 71 countries between 2000 and 2010, with substantial increases recorded for low- and middle-income countries (LMICs)^[2]. Notably, Nigeria contributes 60% of the health products consumed in the Economic Community of West African States (ECOWAS) sub-region, which has a population of 600 million^[3].

Rational use of medicines (antibiotics), an important component of good clinical practice, requires that "patients receive medicines (antibiotics) appropriate to their clinical needs and at the lowest cost"^{[4][5]}. In many countries, 80% of antibiotics used occur outside healthcare facility settings; more than half are unnecessary and inappropriate^[6]. Out of 12 developing countries, Nigeria had the highest average number of medicines prescribed to patients per encounter (3.8 medicines/encounter) and the third highest percentage of antibiotic prescriptions (48%) during a field test^[7].

While some patients are prescribed unnecessary courses of antibiotics, access remains an issue for others. In LMICs, 40% of antibiotics prescribed are insufficient doses, and half of those on therapy adhere to their regimens. Barriers to antibiotic access may be due to limited access to healthcare facilities, the unavailability of antibiotics in local markets, the cost of medicines, often in the face of high out-of-pocket spending, or ineffective medicine supply chain systems^{[8][9][10]}. Patients may also be consuming counterfeit antimicrobials, which often contain suboptimal doses of the agent^{[11][12]}.

In May 2015, the 68th World Health Assembly (WHA) recognised AMR as a threat to global health and requested that member States participate in an integrated global programme for surveillance of AMR and adopt the global action plan. Nigeria committed to establishing a national AMR surveillance system in 2016, in conformity with the global action plan on AMR.

To provide baseline information to guide the proper implementation of the national AMR response, we conducted a systematic review to synthesise the evidence on antibiotic use in humans in Nigeria.

Methods

The protocol for the systematic review was developed by an AMR Working Group set up by the Nigeria Center for Disease Control (NCDC). We searched for articles published in English between January 2000 and January 2017 using Medline via PubMed and African Journals Online (AJOL) databases. The following search terms were used: "rational use", "antibiotic consumption", "antimicrobials", "antibiotic", "Nigeria", "purchase of antibiotics" and "over the counter".




Figure 1. Classes of antibiotics introduced or approved from 1935 to 2012

The titles and abstracts of all potential papers were assessed independently by three of the authors (A.E., Y.E., O.A.), and the lists were compared to ensure that all studies had been identified and duplicates removed. Titles and abstracts of compiled studies were thereafter screened and followed by a full-text review. Articles that were not published in Nigeria were excluded ab initio. We extracted the relevant data using a database that listed the variables: author, year of study, location of study, study design, study setting and demographics, sample size and sampling method, exclusion criteria, commonest antibiotic used/prescribed, quantitative indicators and possible drivers of irrational antibiotic use. We extracted data and reported it as outlined by the reviewers without any alteration using the PRISMA statement, the preferred reporting system for systematic review and meta-analysis [13]. Any inconsistencies were discussed and finalised by the team members. Following our search, 849 articles were identified (Figure 2). Duplicates were removed, and 770 records were excluded based on incompatible titles and abstracts. Other articles were excluded because the article's outcome was not relevant to our objectives (3), only abstract data was available (11), three were reports (no primary data), and one was a field test. From the systematic reviews, drivers of irrational medicine were also compiled.

Data management

Two dimensions of irrational antibiotic use were searched: prescription behaviour by prescribers (supply) and medicine use behaviour by patients (demand). The possible drivers of irrational antibiotic use were summarised. Data was entered into purpose-built Word and Excel templates. The data was analysed as frequencies and proportions, while free-text responses were synthesised.




Figure 2. Summary of the systematic review process

Results

In total, 11 articles were included in the final data synthesis (Figure 2). All articles included used a cross-sectional design, and

the majority (90%) were conducted in clinical settings. Three (27%) of the studies were from the southeast zone, four (36%) from the southwest, and two (18%) each were from south-south and north-central parts of Nigeria. No compatible articles were identified from the north-west or north-eastern zones. The study population in two (18%) of the identified studies comprised entirely of medical doctors.

Regarding antibiotic prescription, four articles (36%) reported the percentage of antibiotics prescribed per patient encounter. Among these articles, the percentage of antibiotics prescribed per encounter ranges from 26.8% among persons ≥ 15 years to 71.1% in children < 5 years and with a median of 44.7%^{[14][15][16]}. One study (9%) found that 49.4% of patients attending NHIS clinics were prescribed antibiotics compared with 33.6% attending general outpatient (non-NHIS) clinics^[17]. From three studies (27%), the percentage of antibiotics of all prescribed medicines among children under five years ranged from 13.0% to 53.7%^{[14][16][18]}. Unnecessary prescription of antibiotics by medical doctors was also reported, with 56.4% of prescribers providing antibiotics empirically to children less than five years old visiting four clinics in one study^[19]. With respect to underuse of antibiotics, 25.5% of house officers were not confident in prescribing antibiotics to patients.

Concerning antibiotic use, three studies (27%) reported using antibiotics without a prescription. The proportion of persons using antibiotics without prescription ranged from 31.7% to 71.7%, with a median of 46.7% and the highest proportion reported among children < 5 years^{[20][21][22]}. One study (9%) found that only 42% of adults completed the recommended antibiotic course^[22]. Eight (73%) articles, as outlined in Figure 3, reported the commonest antibiotic used or prescribed, which were Metronidazole for diarrhoeal diseases (71.4%), Penicillins (71.2%), Amoxiclav (70.9%), and Cotrimoxazole for sore throat (53.1%)^{[20][21][22][14][15][16][18][19]}.

From these studies, poor regulation of medicines, premises and practitioners, volume of counterfeit medicines, chaotic drug distribution system and over-reliance on imported medicines were some health system factors that may be contributory to irrational antibiotic and medicine use in humans and animals^{[23][24]}. Non-client-friendly opening hours by licensed prescribers/health facilities and geographical accessibility to OTC premises were also contributory^[25]. Prescribers with longer years of practice who were non-specialised and older were more likely to give antibiotics empirically to children. In comparison, doctors with shorter years of practice were not confident in prescribing antibiotics to patients without supervision^{[19][26]}. Patients on the National Health Insurance Scheme and demand for antibiotics by patients or their caregivers were also highlighted as some patient-related factors predisposing Nigerian communities to higher antibiotic prescriptions^{[17][27]}. Furthermore, having a caregiver with a lower education level, older age of caregivers and being a child contributed to higher chances of antibiotic use or prescription^{[20][21][27]}.




Figure 3. Antibiotics used by or prescribed to humans in Nigeria, 2000-2017 (n=11)

Discussion

Overall, one in two persons took unprescribed antibiotics or left the health facility with an antibiotic prescription, with higher proportions reported among children. At the health facility, irrational antibiotic prescriptions may be attributable to clinicians' lack of diagnostic support and tools and the prohibitive cost of healthcare to the majority of the population, worsened by limited health insurance coverage^{[28][29]}. The national standard treatment guidelines (STG) aid empiric medicine prescriptions in Nigeria. However, the guidelines are not widely accessible, and AMR surveillance data do not guide the antibiotic recommendations^[30]. Increasing access to diagnostic assays and tools for the healthcare provider and the patient (including financial access) are important interventions that would enhance appropriate antibiotic use and improve AMR detection in LMICs such as Nigeria, especially for children who appear to be more at risk^[31].

The most common antibiotic used or prescribed was the penicillin group. According to a study on global antibiotic consumption, the most consumed antibiotics were cephalosporins and broad-spectrum penicillins, at 55% of the total standard units, with both medicines accounting for the largest absolute increases in consumption between 2000 and 2010 worldwide^[30]. The implication is that the selective pressure on this class of antibiotics remains high and may account for the reduction in the therapeutic effect of these antibiotics, which may worsen if no action is taken. The occurrence of AMR in humans results from selection pressure on human microbiota following constant exposure to antibiotic residues or acquisition of resistant microbial genes from animals and/or the environment^{[31][32][33]}. As such, a One Health approach to AMR surveillance is necessary to break the chain of transmission in the tripartite sectors, and quantifying the antibiotics sold for use in animals is a critical step in defining a strategy for AMR prevention in humans and animals.

Concerning drivers of irrational antibiotic use in this review, enrollment in the National Health Insurance Scheme (NHIS) clinic resulted in higher chances of being prescribed antibiotics^[17]. The NHIS is a social security mechanism aimed at improving access to health services, encouraging the use of generic medicines, including antibiotics, and patients' contributions to co-payments are sometimes as low as 10% of total costs^{[34][35]}. However, delays in reimbursing premiums to the healthcare facilities, combined with shortfalls in the amounts disbursed, as the government-set prices for health services are lower than the real costs, result in healthcare providers prescribing antibiotics empirically to minimise financial loss^[36].

Another driver was patients' demand for antibiotics, which may be real or perceived by the healthcare provider^{[36][37]}. Patients' beliefs that antibiotics are medications that can cure or prevent any illness often increase their expectations of antibiotic therapy^{[38][39][40]}. At other times, healthcare providers may feel that patients would prefer an antibiotic prescription; the patient takes the medication because they feel the health provider is more knowledgeable^[41]. This interaction becomes a vicious cycle of demand and supply for unnecessary antibiotics during patient-health provider interactions. This highlights the need for patient and health provider education as key to reducing antibiotic misuse in Nigerian communities and health facilities.




Figure 4. Factors contributing to irrational antibiotic prescription and or use

We also found that over-the-counter access to antibiotics was a driver for irrational antibiotic use. The sale of antibiotics without prescription is widespread and poorly controlled^[29]. By law, medical doctors in tertiary, secondary and private hospitals are authorised to prescribe medicines, including antibiotics and pharmacists are licensed to dispense antibiotics^{[42][43]}. However, a

high patient-to-doctor ratio has been reported in Nigeria, far below the 100 per 100,000 population recommended by the WHO, and these data is likely to be an underestimate given that the numbers reported do not account for deaths for doctors that are no longer practising or those practising outside the country [41][44][45]. The patient-to-pharmacist ratio is even higher. The paucity of licensed healthcare workers contributes significantly to increased work pressure and longer waiting hours in the clinic, thus limiting healthcare worker-patient interaction. In combination with other factors such as medicine stockouts at healthcare facilities, sparseness of licensed retail pharmacies and geographical accessibility of OTC premises, clients are “pushed” to seek care outside the healthcare facility [38][39]. These OTC premises often stock counterfeit medicines and are closer to poorer, hard-to-reach communities where regulatory oversight by the Food and Drug Administration agencies is limited. The National Drug Policy, Essential Medicines List and the National Drug Distribution Guidelines are some documents that have been developed as part of the country’s efforts to streamline medicine distribution [37][40][41].

We acknowledge that the data search ended in 2017 when the situation analysis was conducted in Nigeria. For the systematic review, we only searched PubMed and AJOL for literature; some journals are not indexed on these databases. We did not explore possible grey literature by contacting known experts for potential articles that are yet to be published. However, the evidence provided by this review represents a baseline assessment of the situation on antimicrobial use, which provides a comparator for future assessments. Understanding the situation in Nigeria is an excellent case study of the challenges low-income countries face in detecting and preventing AMR.

In conclusion, Nigeria has many legislations, albeit outdated, the workforce for health remains low and irrational antibiotic use is widespread in humans and animals, driven by uncontrolled access to antibiotics. Therefore, educating the public on irrational antibiotic use and AMR is crucial, and regulations on antibiotic prescription and sales need to be enforced. Universal health coverage and surveillance of resistant microorganisms should be prioritised in Nigeria. This will entail advocacy and training of the stakeholders and a whole-of-society approach. The gaps identified from these reviews formed part of the formative data to develop the first National Action Plan for AMR in 2017.

Tables

Table 1. Characterization of eligible studies on rational/irrational antibiotic use in Nigeria

	Author(s)	Year	Location	Study design	Demographics	Study setting, Sample size	Sampling method	Exclusion criteria	Commonest antibiotics used/prescribed	Quantitative indicator(s)	Drivers of irrational antibiotic use or prescription
1	Ekwochi	2014	Enugu, South	Prospective	Mother-child ≤5	Health facility	Convenience		Cotrimoxazole	% patients using antibiotics without prescriptions	Availability of the drugs over-the-counter in 163 (42.4%) cases Children older than 11 months (2–5.96%)

1	et al	2014	South east	observational	years pairs	facility, 423	Convenience		(171,53.1%)	prescriptions: 71.7%	($\chi^2=0.00$, $p=0.051$) Mothers with secondary education compared with those with post-secondary education (OR= 1.36, CI= 0.71–2.60)
2	Ekwochi et al	2012	Enugu, South east	Prospective observational	Mother-child ≤ 5 years pairs	Health facility, 210			Metronidazole (70,71.4%)	% patients using antibiotics without prescriptions: 46.7%	Efficaciousness of antibiotic 31(31.6%). Availability of the drugs over-the-counter 28(28.6%). Decreasing level of maternal education ($\chi^2= 10.068$, $p=0.018$)
3	Fadare et al	2015	Ekiti, South west	Cross-sectional, medical records review	Children <5 years	Health facility, 526	Systematic		Penicillins (71.2%)	% of patients/drug encounters with antibiotic prescribed: 374,71.1% % antibiotics of all prescribed medicines: 28.2%	
4	Sanya et al.	2013	Oyo, South west	Cross-sectional	Non-medical undergraduate students	University, 400	Purposive	Postgraduate and undergraduate students from the Faculties of Pharmacy, Veterinary Medicine and	Ampiclox (38,29.9%)	% patients using antibiotics adhering to treatment regimen: 168,42.0% % patients using antibiotics without	Reasons: financial incapability to purchase full dose (73; 18.3%), long duration of treatment (70; 17.5%), side effects experienced (60;15.0%). Commonest reason for buying antibiotics without prescription was for treatment of boils 28, 20.7%.

								College of Medicine		prescriptions: 273,31.7%	Response on what was done with unfinished doses of antibiotics: 187 (46.8%) kept it for future use, 111 (27.8%) threw it away with refuse, 30 (7.5%) flushed it down the toilet while 25 (6.3%) gave it to other people who may have similar complaints.
5	Sadoh et al	2009	Edo, South south		Medical doctors	Health facility, 55			Amoxiclav (39,70.9%)	% prescribers providing antibiotic empirically: 31, 56.4%	Speciality: Paediatricians vs non-paediatricians (45.2 % vs 54.8 %) gave empiric treatment. Paediatricians vs non-paediatricians (88.0 % vs 60.0 %) examined the throats of children most of the time p= 0.032 Years of practice: practicing for over five years Age: Doctors older than 30 years
6	Rasaq et al	2013	Oyo, South west	Cross-sectional	Patients ≥15 years, health workers	Health facility, 400			Amoxicillin (77,25.4%)	% of patients/drugs encounter with antibiotic prescribed: 26.8%	
7	Fadare et al	2014	Ekiti, South west	Cross-sectional	Patients ≥18 years	Health facility, 454	Systematic			% of patients/drugs encounter with antibiotic prescribed: 49.4% vs 33.6% (NHIS vs general clinic)	Attending NHIS clinic resulted in higher chances of antibiotic prescriptions.
										% prescribers not	

8	Adetutu et al	2012	FCT, North central	Cross-sectional	Medical doctors (interns)	Health facility, 30				% prescribers not confident prescribing antibiotics (without supervision): 25.5%	
9	Akande et al	2007	Ilorin (Kwara), North Central	Cross-sectional, prescription record review	Patients	Health facility, 303	Systematic		Amoxicillin (25.0%)	% of patients/drug encounters with antibiotic prescribed: 135, 44.7%	
10	Nwolisa et al	2004	Imo, South east	Prospective observational, medical record review	Children <5 years	Health facility, 709	Total	Patients who had no drugs prescribed	Cephalexin (132,31.1%)	% antibiotics of all prescribed medicines: 53.7%	Age group 13 to 24 months accounted for the highest number of antibiotic prescriptions.
11	Sadoh et al	2013	Edo, South south		Children	Health facility, 320	Convenience			% patients/caregivers requesting for antibiotics: 174,57.2%	<p>29(42.2%) will not be satisfied with a doctor who does not prescribe an antibiotic for their child's sore-throat.</p> <p>70(22.7%) respondents would change their doctors for not prescribing an antibiotic.</p> <p>Respondents (caregivers) with secondary and tertiary levels of education 59.0% and 56.6% respectively compared to 20.6% with primary level of education would not request for antibiotic.</p> <p>p <0.001</p> <p>Older caregivers 71(52.2%) in the 30 – 39 years' age group and 22(57.9%) ≥40 years compared to younger</p>

References

1. ^a Nirjana SE, Sambath Kumar R, Sudha M, Vwnkateswaramurthy N. Review on clinically developing antibiotics. *International Journal of Applied Pharmaceutics*. 2018;10(3):13–8.
2. ^a Van Boeckel TP, Gandra S, Ashok A, Caudron Q, Grenfell BT, Levin SA, et al. Global antibiotic consumption 2000 to 2010: An analysis of national pharmaceutical sales data. *The Lancet Infectious Diseases*. 2014;14(8):742–50.
3. ^a Nigeria Center for Disease Control. *Antimicrobial use and resistance in Nigeria: situation analysis and recommendations*. 2017.
4. ^a World Health Organisation. WHO | Rational use of medicines [Internet]. World Health Organization; 2015 [cited 2017 Mar 8]. Available from: http://www.who.int/medicines/areas/rational_use/en/
5. ^a Sartelli M, Weber DG, Ruppé E, Bassetti M, Wright BJ, Ansaloni L, et al. Antimicrobials: a global alliance for optimizing their rational use in intra-abdominal infections (AGORA). *World J Emerg Surg*. 2016;11:33.
6. ^a Wise R, Hart T, Cars O, Streulens M, Helmuth R, Huovinen P, et al. Antimicrobial resistance is a major threat to public health. *BMJ*. 1998;317:609–10.
7. ^a Hogerzeil HV, Bimo, Ross-Degnan D, Laing RO, Ofori-Adjei D, Santoso B, et al. Field tests for rational drug use in twelve developing countries. *Lancet*. 1993;342:1408–10.
8. ^a Miller-Petrie M, Gelband H. Socioeconomics, antimicrobial use and antimicrobial resistance | AMR control. *Economics and Innovation*. 8 2017;23:812–8.
9. ^a Cox JA, Vlieghe E, Mendelson M, Wertheim H, Ndegwa L, Villegas MV, et al. Antibiotic stewardship in low- and middle-income countries: the same but different? *Clinical Microbiology and Infection*. 2017 Nov 1;23(11):812–8.
10. ^a Laxminarayan R, Duse A, Wattal C, Zaidi AKM, Wertheim HFL, Sumpradit N, et al. Antibiotic resistance - the need for global solutions. *Lancet Infect Dis*. 2013;13:1057–98.
11. ^a Komolafe OO. Antibiotic resistance in bacteria - an emerging public health problem. *Malawi Medical Journal*. 2004 Feb 6;15(2):63.
12. ^a Ayukekbong JA, Ntemgwa M, Atabe AN. The threat of antimicrobial resistance in developing countries: Causes and control strategies. *Antimicrob Resist Infect Control*. 5 2017;6:47.
13. ^a Moher D, Liberati A, Tetzlaff J, Altman DG, Grp P. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med*. 2009;6:e1000097.
14. ^{a, b, c} Fadare J, Olatunya O, Oluwayemi O, Ogundare O. Drug prescribing pattern for under-fives in a paediatric clinic in South-Western, Nigeria. *Ethiop J Health Sci*. 2015;25:73–8.
15. ^{a, b} Adisa R, Fakeye TO, Aindero VO. Evaluation of prescription pattern and patients' opinion on healthcare practices in selected primary healthcare facilities in Ibadan, South-Western Nigeria. *Afr Health Sci*. 2015;15:1318–29.
16. ^{a, b, c} Akande TM, Ologe MO. Prescription pattern at a secondary health care facility in Ilorin, Nigeria. *Ann Afr Med*. 2007;6:186–9.
17. ^{a, b, c} Fadare JO, Adeoti AO, Aina F, Solomon OA, Ijalana JO. The influence of health insurance scheme on the drug prescribing pattern in a Nigerian tertiary healthcare facility. *Niger Med J*. 2015;56:344–8.
18. ^{a, b} Nwolisa CE, Erinaugha EU, Ofoleta SI. Prescribing practices of doctors attending to under fives in a children's outpatient clinic in Owerri, Nigeria. *J Trop Pediatr*. 2005;52:197–200.
19. ^{a, b, c} Sadoh WE, Akinsete AM. Physicians management of sore throat in children in Benin city, Nigeria. *Niger J Clin Pract*.

- 2009;12:407–11.
20. ^{a, b, c}Ekwochi U, Chinawa JM, Osuorah CDI, Odetunde OI, Obu HA, Agwu S. *The Use of Unprescribed Antibiotics in Management of Upper Respiratory Tract Infection in Children in Enugu, South East Nigeria.* *J Trop Pediatr.* 2014;60:249–52.
 21. ^{a, b, c}Ekwochi U, Chinawa JM, Obi I, Obu HA, Agwu S. *Use and/or Misuse of Antibiotics in Management of Diarrhea Among Children in Enugu, Southeast Nigeria.* *J Trop Pediatr.* 2013;59:314–6.
 22. ^{a, b, c}Sanya TE, Titilayo OF, Adisa R, Segun JS. *Use of antibiotics among non-medical students in a Nigerian university.* *Afr Health Sci.* 2013;13:1149–55.
 23. [^]UNIDO. *Pharmaceutical Sector Profile: Nigeria Global.* Vienna; 2011.
 24. [^]West African Health Organisation. *The Economic Community of West African States (ECOWAS) Regional Pharmaceutical Plan (ERPP). Bobo-Dioulasso: WAHO Essential Medicines and Vaccines programme; 2014.*
 25. [^]Kombe G, Fleisher L, Kariisa E, Arur A, Sanajan P, Paina L. *Nigeria Health System Assessment.* Washington, DC: Abt Associates Inc; 2008.
 26. [^]Ajemigbitse AA, Omole MK, Ezike NC, Erhun WO. *Assessment of the knowledge and attitudes of intern doctors to medication prescribing errors in a Nigeria tertiary hospital.* *Journal of basic and clinical pharmacy.* 2013;5:7–14.
 27. ^{a, b}Sadoh WE, Sadoh AE, Eki-Udoko FE. *Parental contribution to over prescription of antibiotics for sore throat in children.* *Niger J Paediatr.* 2015;42:98–102.
 28. [^]Federal Ministry of Health Nigeria. *Standard treatment guidelines Nigeria.* 2008.
 29. ^{a, b}World Health Organization. *WHO | Density of medical doctors (total number per 10 000 population, latest available year) [Internet].* World Health Organization; [cited 2020 Jun 29]. Available from: http://www.who.int/gho/health_workforce/doctors_density/en/
 30. ^{a, b}Bebell LM, Muiru AN. *Antibiotic use and emerging resistance: How can resource-limited countries turn the tide?* *Glob Heart.* 2014;9:347–58.
 31. ^{a, b}Nigeria National Population Commission, ICF International. *Nigeria Demographic and Health Survey.* 2013.
 32. [^]National Health Insurance Scheme. *Operational Guidelines [Internet].* [cited 2017 Mar 15]. Available from: http://www.nhis.gov.ng/file/repository/NHIS_OPERATIONAL_GUIDELINES.pdf
 33. [^]Academy of Managed Care Pharmacy. *Formulary Management.* 2009.
 34. [^]World Health Organization, Harvard Medical School and Harvard Pilgrim Health. *Medicines use in primary care in developing and transitional countries: fact book summarizing results from studies reported between 1990 and 2006.* *Bulletin of the World Health Organization [Internet].* 2009;87(10). Available from: <http://dx.doi.org/10.2471/09-070417.doc>
 35. [^]Holloway KA. *Promoting the rational use of antibiotics.* *Regional Health Forum.* 2011;15:122–30.
 36. ^{a, b}Okeke IN, Lamikanra A, Edelman R. *Socioeconomic and behavioral factors leading to acquired bacterial resistance to antibiotics in developing countries.* *Emerging infectious diseases.* 1999;5(1):18–27.
 37. ^{a, b}Nordberg P, Stålsby-lundborg C, Tomson G. *Consumers and providers – Could they make better use of antibiotics?* *Int J Risk Saf Med.* 2005;17:117–25.
 38. ^{a, b}Olusegun A, Adeniyi A. *Irrational use and non-prescription sale of antibiotics in Nigeria: A need for change.* *Journal of Scientific and Innovative Research.* 2014;3:251–7.
 39. ^{a, b}Global Antibiotic Resistance Partnership-India (GARP-India). *Situation Analysis: Antibiotic Use and Resistance in India.*

2011.

40. ^{a, b} Morgan DJ, Okeke IN, Laxminarayan R, Perencevich EN. *Non-prescription antimicrobials use worldwide: a systematic review. NIH Public Access. 2011;11:692–701.*
41. ^{a, b, c} *National Drug Policy. 2005.*
42. [^] Kombe G, Fleisher L, Kariisa E, Arur A, Sanjana P, Paina L, et al. *Nigeria Health System Assessment 2008. In 2009. p. 1–135.*
43. [^] *Federal Ministry of Health. National Strategic Health Development Plan (NSHDP) 2010-2015 [Internet]. 2010. Available from: [http://dx.doi.org/10.1016/S1474-4422\(11\)70006-0](http://dx.doi.org/10.1016/S1474-4422(11)70006-0)*
44. [^] *Federal Ministry of Health. National Drug Distribution Guidelines. 2nd ed. Abuja; 2012.*
45. [^] *Federal Ministry of Health, World Health Organization, European Commission. Essential medicines list. 5th ed. Abuja; 2010.*