

[Open Peer Review on Qeios](#)

Maternal Misconceptions Against Infant Sunlight Exposure Are Still Bottlenecks in Northwest Ethiopia, by 2022

Besfat Erega¹¹ Debre Tabor University**Funding:** No specific funding was received for this work.**Potential competing interests:** No potential competing interests to declare.

Abstract

Background: Sunlight has been used for therapeutic purposes for neonatal jaundice. Exposure to sunlight is vital for the synthesis of vitamin D. Lack of exposure to sunlight is the single most important cause of rickets. Rickets is a major public health problem in many countries of the world, especially in Sub-Saharan Africa. Vitamin D deficiency due to low exposure to sunlight, suboptimal exclusive breastfeeding, and low vitamin D intake is one of the health problems in Ethiopia. The implementation of the strategy has remained inconsistent and lacked focus on factors that influence maternal practice, excluding infants from getting adequate sunlight.

Objective: To assess knowledge, practices, and factors affecting sunlight exposure of infants among women attending Debre Tabor Comprehensive Specialized Hospital, Ethiopia, 2022.

Methods: An institutional-based cross-sectional study was conducted through patient interviews at Debre Tabor Comprehensive Specialized Hospital from May 2022 to July 2022. Study participants were selected using a systematic random sampling method. A pretest was conducted on 5% (18) of participants before the actual data collection began. The data was collected by five BSc midwives. Data was entered into EpiData version 4.6.2 and exported to SPSS Software version 25 for further analysis. Bivariate and multivariate logistic regression analyses were carried out to test the association between each independent variable and the dependent variable. At 95% CI, variables with a p-value < 0.05 in the multivariate logistic regression model were considered as associated factors.

Results: About 67.5% and 62.1% of the mothers had good knowledge and good practice about adequate sunlight exposure, respectively. Mother's age, mother's educational status, marital status, and family size were the factors associated with sunlight exposure in multinomial logistic regression.

Conclusion and Recommendations: According to our study, participants did not have good knowledge and practice regarding sunlight exposure of infants. Therefore, health education focusing on the importance of sunlight exposure is important to improve the knowledge and practice of mothers towards sunlight exposure of their infants.

Besfat Berihun Erega (BBE)**Department of Midwifery, College of Medicine and Health Sciences, Debre Tabor University, Ethiopia*

*Correspondence: berihunbesfat@gmail.com

Keywords: Sunlight, Mothers, Debre Tabor, Ethiopia.

Background

For years, sunlight has been used for therapeutic purposes, which dates back to ancient Rome and Greece^[1]. In the second half of the 19th century, it was realized that sunshine could have a bactericidal effect as well as a therapeutic role in rickets^{[2][3][4][5]}. In the 1950s, sunlight was first used for neonatal jaundice^{[2][3][4][5][6][7]}. Placing a child in a room where sunlight enters through window panes (not in direct sunlight) for 10 minutes twice a day was often used to help treat mild neonatal jaundice^[8]. Currently, exposure to sunlight has two interests: beneficial and adverse effects on human health.

In a study conducted at Aleta Wondo Health Center, SNNPR, Ethiopia, more than half, 196 (62.8%), of the study participants were knowledgeable about sunlight exposure^[9].

Another study conducted in Debre Markos and Dejen District, Amhara Region, Northwest Ethiopia, found that 433 (50%) had poor knowledge status regarding infant sunlight exposure. The majority of mothers (852, 98.4%) knew that infant sunlight exposure was essential for the infant, and 373 (43.1%) had answered that infant sunlight exposure is for the development, growth, and strength of bones, and 779 (90%) of mothers knew relatively the safest period of infant sunlight exposure^{[10][11]}.

A study done in Farta District, South Gondar Zone, Northwest Ethiopia showed that nearly half (50.1%) of the respondents had knowledge of sunlight exposure for their infant. Of 183 respondents, 84 (46%) had knowledge of sunlight exposure every day, and 39 (11.4%) two times per week^[12].

In a study done at St. Paul's Hospital, Ethiopia, about 40% of the mothers had poor practice regarding sunlight exposure. Out of 346 respondents, 318 (91.1%) of mothers exposed their babies to sunlight^[13].

A study conducted in Debre Berhan Town, North Shewa Zone, Amhara Region, Ethiopia revealed that 65.7% of them had good practice in sunning their infants. Almost all (525; 99.1%) respondents intentionally expose their infants to direct sunlight. Four hundred and fifty (85.7%) of the mothers start to expose their infants before they are 15 days old^[14].

A study done in Debre Markos Town, Ethiopia found that 60% of mothers had poor knowledge about sunlight exposure of infants^[10].

A study in Ethiopia also found that a significant association was observed between maternal age and the knowledge of mothers; mothers in the age group of above 33 years were 8.67 times more likely to be knowledgeable than mothers in the age group of 15-20 years. The educational status of mothers was also associated with knowledge; mothers who have

a diploma and above were 3.24 times more knowledgeable than mothers who were unable to read and write ^[11]. A family size of 4-6 was 3.88 times more knowledgeable than a family size of 1-3. Regarding sunlight exposure practice, there was a statistically significant association between maternal age and practice. A study conducted in Debre Berhan Town, North Shewa Zone, Amhara Region, Ethiopia showed that maternal age, occupational status, perceived benefits to strengthen infant bones, perceived to result in a healthier infant, and source of information from health-care professionals were found to be significantly associated with maternal practice of infant sunlight exposure ^{[12][13][14][15][16][17][18][19]}.

A study carried out in Debre Markos Town, Ethiopia, found that there was a significant association between maternal age, maternal education, family size, husband's educational status, and knowledge about sunlight exposure ^[10].

Methods and Materials

Study Area and Period

The study was conducted at Debre Tabor Comprehensive Specialized Hospital, Northwest Ethiopia. Debre Tabor is the capital city of the South Gondar zone, located about 666 kilometers from Addis Ababa (the capital city of Ethiopia) and 100 kilometers from Bahir Dar. This town has a latitude and longitude of 11°51'N 38°1'E with an elevation of 2,706 meters (8,878 ft) above sea level. The hospital was established by Italian missionaries in 1928. Currently, the hospital serves around 2.7 million people in the zone. Based on the facility-based report of the 2010 Ethiopian fiscal year, 1,268 children were delivered at Debre Tabor Comprehensive Specialized Hospital. The study was carried out from May 2022 to July 2022.

Study Design

A cross-sectional institutional-based study was conducted.

Source Population

The source populations were all mothers with infants attending PNC and immunization clinics at Debre Tabor Comprehensive Specialized Hospital.

Study Population

The study population were mothers with infants attending PNC and immunization clinics at Debre Tabor Comprehensive Specialized Hospital during the data collection period and who fulfilled the inclusion criteria.

Eligibility Criteria

Inclusion criteria

- Mothers of infants aged < 1 year attending PNC and immunization clinics at DTCSH during the study period.

Exclusion criteria

- Mothers with infants who were seriously ill, mentally incompetent, and unable to communicate.
- Those mothers with infants who were not willing to participate in the study.

Sample Size Determination

The sample size was calculated using the single population proportion formula by assuming p = prevalence of mothers' knowledge and practice on sunlight exposure in Debre Markos town = 44.6%(37), $Z = 1.96$, and D = precision (marginal error) = 5%. The sample size was calculated as follows:

$$n = \frac{(Z_{\alpha/2})^2 P(1 - P)}{d^2} = \frac{(1.96)^2(0.446)(0.554)}{(0.05)^2} = 379.679 \sim 380$$

$$n = 380$$

Where: - n = sample size

The total number of infants in Debre Tabor Town is 2,277. Since this figure is below 10,000, we use the following adjustment formula for the sample size: $n = n / (1 + n/N)$

Where,

- n = sample size for a population of size above 10,000
- N = number of infants in Debre Tabor Town Therefore,
- $n = 380 / (1 + 380/2,277) n \approx 341$

Taking a 10% (34) non-response rate, the final sample size was **375**.

Sampling Procedure

Study subjects were selected using systematic random sampling. The sampling interval (K) was obtained by dividing the total number of mothers with their infants by the sample size at the vaccinated clinic in Debre Tabor Hospital, so that $K = N/n =$ every 3rd study subject was interviewed until we reached 375 study subjects. The first study subject was selected by the lottery method from the sampling interval (K). Each study participant was selected using a systematic sampling technique in which every third client was interviewed in the Hospital.

Study Variables

Dependent variables

- Knowledge about sunlight exposure
- Practice of sunlight exposure

Independent Variables

Socio-demographic factors (age, religion, ethnicity, marital status, educational status, occupation)

Source of information (physician, nurse/midwife, TV/Radio, neighbours/elder people)

Other factors for sunlight exposure: Mothers' fear (Fear of Sickness, evil eye, Cold, Pneumonia)

Operational Definitions

Knowledge - The theoretical understanding of mothers about sunlight exposure of infants.

Good knowledge - Those mothers who responded to knowledge questions and scored above the median value of 4.

Poor knowledge - Those mothers who responded to knowledge questions and scored below the median value of 4.

Practice - Mothers' activity or behavioural experience in relation to sunlight exposure of infants.

Good practice - Mothers who responded to practice questions and scored above the median value of 6.

Poor practice - Mothers who responded to practice questions and scored 6 and below the median value.

Data Collection Tools and Procedure

A structured questionnaire was designed by reviewing different literature to include all the relevant variables to meet the objectives.

After the informed consent was obtained, data were collected through face-to-face interviews using a prepared standard checklist and structured questionnaire. Three nurses and two midwives who were working at the immunization center were considered as data collectors.

Data Quality Assurance

The questionnaire was sent to two senior staff who are academicians and researchers for face validity; approval was obtained, and modifications were made. The questionnaire was prepared in the English version and translated into Amharic and back to English to check its consistency. A 5% pretest was done at Hidar 11 health center outside the study area. The collected data were checked for completeness and the overall quality of data collection was monitored by the principal investigators.

Data Analysis Procedure

Data were entered into EpiData version 4.6.2 and exported to SPSS Software version 25 for further analysis. Mean, standard deviation, frequencies, percent, and odds ratio were calculated. Binary and multiple logistic regression analyses were conducted to assess the association between independent and dependent variables. The strength of the statistical association was measured by odds ratio and 95% confidence intervals, and statistical significance was considered at $P < 0.05$.

Results

Socio-Demographic Characteristics of Mothers

A total of 375 mothers with a response rate of 100% were involved. Of these, 176 (46.9%) of them were in the age range of 21-26 years and 361 (96.3%) of them were married (**Table 1**).

Knowledge of Mothers About Sunlight Exposure of Their Infants

About 253 (67.5%) of the mothers had good knowledge about sunlight exposure. The overall knowledge of mothers about the effect of sunlight exposure was 77.8% (**Table 2**).

Practice of Mothers About Sunlight Exposure of Their Infants

Regarding the practice of mothers about the exposure of infants to sunlight, 233 (62.1%) of them exposed their babies to sunlight and 36.9% of the mothers started exposing their infants at the age of 45 days and above (**Table 3**).

Factors Associated with Sunlight Exposure

Mother's age (21-26 (AOR=2.862 (1.862, 4.399)), 27-32 (AOR= 0.255 (0.154, 0.422)), and 33+ (AOR=1.909 (1.300, 2.803))), mother's educational status (grade 7-10 (COR= 0.486 (1.253-4.933)), grade 11-12 (AOR= 1.531 (1.531-3.214)), certificate and above (AOR=1.498 (1.909-2.471))), marital status (divorced (AOR=1.411 (0.144-0.177))), and family size (4—6 (AOR=3.28 (2.189-4.903))) were the factors associated with sunlight exposure in multinomial logistic regression (**Table 4**).

List of Tables

Table 1. Distribution of socio-demographic characteristics of mothers in Debre Tabor Comprehensive Specialized Hospital, 2022 (N=375).

| Variables | Frequency | Percentage |
|-----------|-----------|------------|
| | | |

| | | |
|------------------------------|-----|------|
| Mother's age | | |
| 15-20 | 33 | 8.8 |
| 21-26 | 176 | 46.9 |
| 27-32 | 106 | 28.3 |
| 33+ | 60 | 16 |
| Marital status | | |
| Single | 11 | 2.9 |
| Married | 361 | 96.3 |
| Divorced | 3 | 0.8 |
| Mother's educational status | | |
| Unable to read & write | 37 | 9.9 |
| Able to read & write | 54 | 14.4 |
| Grade1-6 | 111 | 29.6 |
| Grade7-10 | 66 | 17.6 |
| Grade11-12 | 15 | 4 |
| Certificate and above | 92 | 24.5 |
| Husband's educational status | | |
| Unable to read & write | 19 | 5 |
| Able to read & write | 42 | 11.2 |
| Grade1-6 | 11 | 2.9 |
| Grade7-10 | 78 | 20.8 |
| Grade11-12 | 64 | 17.1 |
| Certificate and above | 161 | 42.9 |
| Mother's occupation | | |
| Student | 8 | 2.1 |
| House wife | 198 | 52.8 |
| Government employee | 98 | 26.1 |
| Private employee | 16 | 4.3 |
| Daily laborer | 19 | 5 |
| Merchant | 36 | 9.6 |
| Family size | | |

| | | |
|-----|-----|------|
| 1-3 | 236 | 62.9 |
| 4-6 | 127 | 33.9 |
| >6 | 12 | 3.2 |

Table 2. Knowledge of mothers about sunlight exposure of their infants in Debre Tabor Comprehensive Specialized Hospital, 2022 (N=375).

| Variables | Frequency | Percentage |
|---|-----------|------------|
| Had information about sunlight exposure (N=354) | | |
| Yes | 344 | 91.7 |
| No | 31 | 8.3 |
| Source of information about sunlight exposure | | |
| Health professionals | 281 | 74.9 |
| Television/radio | 45 | 12 |
| Neighbors/elder people | 49 | 13.1 |
| Sunlight exposure is beneficial (N=375) | | |
| Yes | 354 | 94.4 |
| No | 21 | 5.6 |
| Sunlight exposure strengthens teeth (N=354) | | |
| Yes | 197 | 55.6 |
| No | 157 | 44.4 |
| Sunlight exposure keeps child warm (N=354) | | |
| Yes | 98 | 27.7 |
| No | 256 | 72.3 |
| Sunlight exposure stimulates vitamin D production (N=354) | | |
| Yes | 112 | 31.6 |
| No | 242 | 68.4 |
| Sunlight exposure strengthens body (N=354) | | |
| Yes | 222 | 62.7 |
| No | 132 | 37.3 |

| | | |
|---|-----|------|
| Sunlight exposure has harmful effect (N=375) | | |
| Yes | 54 | 14.4 |
| No | 321 | 85.6 |
| Skin cancer is a harmful effect of sunlight exposure (N=54) | | |
| Yes | 42 | 77.8 |
| No | 12 | 22.2 |
| Blindness is a harmful effect of sunlight exposure (N=54) | | |
| Yes | 40 | 74 |
| No | 14 | 26 |
| Sterility is a harmful effect of sunlight exposure (N=54) | | |
| Yes | 2 | 3.7 |
| No | 52 | 96.3 |
| Knowledge | | |
| Good | 253 | 67.5 |
| Poor | 122 | 32.5 |

Table 3. Practice of mothers regarding sunlight exposure of their infants in Debre Tabor Comprehensive Specialized Hospital, 2022 (N=375).

| Variables | Frequency | Percentage |
|--|-----------|------------|
| Expose your baby on sunlight (N=375) | | |
| Yes | 233 | 62.1 |
| No | 142 | 38.9 |
| Age infant start sunlight exposure (N=233) | | |
| 0-15 days | 38 | 16.3 |
| 16-30 days | 44 | 18.9 |
| 31-45 days | 65 | 27.9 |
| 45 days and above | 86 | 36.9 |
| Frequently do you expose (N=233) | | |

| | | |
|---|-----|------|
| Daily | 170 | 73.4 |
| Sometimes | 62 | 26.6 |
| Place to expose your baby on sunlight (N=233) | | |
| Outdoor | 207 | 88.8 |
| Indoor | 26 | 11.2 |
| Time of the day do you expose your baby outdoors (N=233) | | |
| Morning 8-10 AM | 214 | 91.8 |
| Mid day 11AM-1 PM | 13 | 5.6 |
| Afternoon 2-4 PM | 6 | 2.6 |
| Condition of clothing during exposure (N=233) | | |
| Unclothed | 88 | 37.8 |
| With diapers and eye protection only | 28 | 12 |
| Partly covered | 77 | 33 |
| Completely covered | 40 | 17.2 |
| Minutes you expose your baby on sunlight (N=233) | | |
| < 15 minute | 83 | 35.6 |
| 15-20 minute | 57 | 24.5 |
| 20-1hr | 88 | 37.8 |
| Above 1hr | 5 | 2.1 |
| Apply lubricants on your baby's body during sunlight exposure (N=233) | | |
| Yes | 145 | 62.2 |
| No | 88 | 37.8 |
| Time to apply lubricants on your baby's body during sunlight (N=233) | | |
| Before exposure | 42 | 18 |
| During exposure | 158 | 67.8 |
| After exposure | 33 | 14.2 |
| Things do you apply (N=233) | | |

| | | |
|---------------|-----|------|
| Baby vaseline | 190 | 81.5 |
| Baby lotion | 18 | 7.7 |
| Butter | 25 | 10.7 |

Table 4. Associated factors with sunlight exposure among mothers in Debre Tabor Comprehensive Specialized Hospital, 2022 (N=375).

| Variables | OR at 95 % CI | |
|-----------------------------|------------------------|------------------------|
| | COR(95% CI) | AOR(95% CI) |
| Mother's age | 1 | 1 |
| 15-20 | 0.432 (0.285, 0.654)** | 2.862 (1.862, 4.399)** |
| 21-26 | 0.255 (0.154, 0.422)** | 4.338 (2.362, 7.964)** |
| 27-32 | 1.909 (1.300, 2.803)** | 1.432 (0.285, 0.654)** |
| 33+ | | |
| Mother's educational status | | |
| Unable to read & write | 1 | 1 |
| Able to read & write | 0.321(0.171-1.600) | 2.666 (0.282-1.282) |
| Grade1-6 | 0.316(0.181-2.554) | 3.271 (0.790-5.978) |
| Grade7-10 | 0.486(1.253-4.933)* | 2.122 (0.048-4.293) |
| Grade11-12 | 0.533(1.265- 2.069)** | 1.531 (1.531-3.214)** |
| Certificate and above | 0.583(0.364-0.934)** | 1.498 (1.909-2.471)** |
| Marital status | | |
| Single | 1 | 1 |
| Married | 3.472(0.137-10.602) | 1.010(0. 1.01-1.772) |
| Divorced | 0.426 (0..108-1.673)** | 1.411(0.144-0.177)** |
| Family size | | |
| 1-3 | 1 | 1 |
| 4-6 | 0.386(0.268-0.557)** | 3.28(2.189-4.903)** |
| >6 | 0.459(0.310-0.679)* | 2.051 (0.337-3.147) |

NB: * = Association in binomial logistic regression, ** = Association in both binomial and multinomial logistic regression

Discussion

This study revealed that about 67.5% of the mothers had good knowledge about sunlight exposure. The result of this study is higher when compared with the results of the studies conducted in Aleta Wondo Health Center, Dejen District,

Farta District, and Debre Markos Town where 62.8%, 50%, 50.1%, and 60% of the respondents had good knowledge of sunlight exposure for their infants, respectively [9][10][11][12]. Such discrepancies might have resulted either from cultural influences or poor awareness. However, this finding is lower when compared with the result found in a study conducted at St. Paul's Hospital, Ethiopia, which showed that 86.1% of the mothers had good knowledge towards sunlight exposure of their infants [20]. The possible reason may be differences in study period, study area, and cultural differences. In addition to this possible justification for the variation, it might be due to differences in the awareness level of mothers concerning the importance of infants' sunlight exposure.

The current study found that the majority (91.7%) of the mothers had information about the need for sunlight exposure. This result is similar to the finding of the study done at Aleta Wondo Health Center, SNNPR, Ethiopia, where 92.3% of the mothers had information about the need for sunlight exposure for their infants [9].

Regarding the practice of mothers, 62.1% of the mothers had good practice regarding adequate sunlight exposure. This finding is comparable with the results of studies conducted at Aleta Wondo Health Center (58%) and St. Paul's Hospital (60%) [11][12][13][14][15]. This finding is higher than the results obtained from the studies conducted in Dejen District, Amhara Region, and Farta District, South Gondar Zone, where 44% and 46% of the mothers had good practice towards sunlight exposure of infants, respectively [10][11][12]. The possible reason for this variation could be differences in the awareness level of mothers concerning the importance of infants' sunlight exposure.

Concerning the factors associated with sunlight exposure of mothers, mother's age, mother's educational status, marital status, and family size were the factors associated with sunlight exposure. Mother's age is significantly associated with sunlight exposure. Mothers in the age group above 33 years were 1.43 times more likely to be knowledgeable than mothers in the age group of 15-20 years. This finding is supported by the study conducted at St. Paul's Hospital, which showed that mothers in the age group above 33 years were 8.67 times more likely to be knowledgeable than mothers in the age group of 15-20 years [20]. This finding is also supported by the studies conducted in Debre Berhan Town and Debre Markos Town, which found that mother's age is significantly associated with sunlight exposure [10][14].

The educational status of mothers and family size are also associated with the knowledge of mothers towards sunlight exposure. This result is supported by the studies conducted at St. Paul's Hospital [20], Debre Berhan Town [14], Debre Markos Town [10], and Dejen District, Amhara Region [11].

Conclusion and Recommendation

According to our study, participants did not have good knowledge and practice regarding sunlight exposure of infants. Mother's age, mother's educational status, marital status, and family size were the factors associated with sunlight exposure.

Therefore, health education focusing on the importance of sunlight exposure is important to improve the knowledge and practice of mothers towards sunlight exposure of their infants.

Abbreviations and Acronyms

| Abbreviation | Full Form |
|------------------|--|
| 25 (OH) D | 25-Hydroxy Vitamin D |
| CI | Confidence Interval |
| DCSH | Debre Tabor Comprehensive Specialized Hospital |
| HF | Health Facility |
| IU | International Unit |
| KM | Kilometre |
| OR | Odds Ratio |
| USA | United States of America |
| UV | Ultraviolet Radiation |
| VDD | Vitamin D Deficiency |
| WHO | World Health Organization |

Statements and Declarations

Availability of Data and Materials

All data included in this manuscript can be accessed from the corresponding author upon reasonable request.

Competing Interests

The authors declare that they have no competing interests.

Funding

No funding.

Consent to Participate

Written informed consent from the respondents was obtained after thoroughly explaining the aim of the study to each respondent.

Consent for Publication

Not applicable.

Code Availability

All code for data cleaning and analysis associated with the current submission is available with the principal investigator and can be presented upon request.

Ethical Approval

Ethical clearance was obtained from the Institutional Review Board of Debre Tabor University, School of Midwifery, College of Medicine and Health Sciences (CMHS). A letter of permission was obtained from the clinical coordinator of each study hospital. A clear explanation about the purpose of the study was given along with the letter of support to all concerned bodies. Finally, written informed consent from the respondents was obtained after thoroughly explaining the aim of the study to each respondent. In addition, all methods were performed in accordance with the relevant guidelines and regulations.

Author's Contribution

BBE is the primary author, participated in the conceptualisation, design, analysis and interpretation of the data, and drafted the manuscript.

Acknowledgments

I am thankful to Debre Tabor University for granting us ethical clearance. We are also grateful to the data collectors, women who participated in the data collection, hospital managers, health professionals, and all individuals who were willing to support us in any kind. I am also thankful to the author of the thesis for availing his online sources.

Author's Information

Besfat Berihun Erega, BSc, MSc in Clinical Midwifery, Lecturer at Debre Tabor University, College of Medicine and Health Sciences, Department of Midwifery

References

- ^a Harrison, S. L., Buettner, P. G., & MacLennan, R. (2002). *Why do mothers still sun their infants? Journal of Paediatrics and Child Health*, 38(3), 296-299.
- ^{a, b} Roelandts, R. (2002). *The history of phototherapy: Something new under the sun? Journal of the American Academy of Dermatology*, 46(6), 926-930.
- ^{a, b} Albert, M. R., & Ostheimer, K. G. (2002). *The evolution of current medical and popular attitudes toward ultraviolet light exposure: Part 1. Journal of the American Academy of Dermatology*, 47(6), 930-937.
- ^{a, b} Newman, B. Y. (2000). *The suntan myth. Optometry*, 71(11), 688.

5. ^{a, b}Albert, M. R., & Ostheimer, K. G. (2003). *The evolution of current medical and popular attitudes toward ultraviolet light exposure: Part 2. Journal of the American Academy of Dermatology*, 48(6), 909-918.
6. [^]Yurdakok, M. (1995). *Hiperbilirubinemide Isık ve İlaç Tedavisi [Light and Drug Treatment in Hyperbilirubinemia]. Katki Pediatri Dergisi [Katki Pediatric Journal]*, 16(5), 722-744. [Article in Turkish]
7. [^]Johnston, R. V., Anderson, J. N., & Prentice, C. (2003). *Is sunlight an effective treatment for infants with jaundice? Medical Journal of Australia*, 178, 403.
8. [^]Emedicine. (2005, August 10). *Newborn Jaundice. Cited 2005, October 16.*
<http://www.emedicinehealth.com/articles/10101-6>.
9. ^{a, b, c}Getaneh, T., Assefa, A., & Tadesse, Z. (1998). *The knowledge and practice of exposure to sunlight in Jimma Town, South Western Ethiopia. Ethiopian Journal of Health Development*, 12, 29-32.
10. ^{a, b, c, d, e, f, g}Lulseged, S., & Fitwi, G. (1999). *Vitamin D Deficiency rickets: Socio-demographic and clinical risk factors in children seen at a referral hospital in Addis Ababa. East African Medical Journal*, 76, 457-461.
11. ^{a, b, c, d, e, f}Sileshi, L., & Catherine, D. (1999). *Behavioral intention and factors influencing intention of Ethiopian mothers to expose infants to sunshine.*
12. ^{a, b, c, d, e}Reddy, V., & Lamb, W. H. (1991). *Nutritional Rickets. In P. Stanfield, M. Brueton, M. Chan, M. Parkin, & T. Waterston (Eds.), Diseases of children in the subtropics & tropics (4th ed., pp. 376-379). Edward Arnold.*
13. ^{a, b, c}Eshetu, M. (1994). *Analysis of paediatric admissions to Jimma Hospital paediatrics ward: A three-year retrospective study. Bulletin of Jimma Institute of Health Sciences*, 4, 1-11.
14. ^{a, b, c, d, e}Tesfaye, G., Afework, A., & Zerihun, T. (1998). *Rickets and the knowledge and practice of exposure to sunlight in Jimma Town.*
15. ^{a, b}Asres, B., Melese, G., Bedilu, D., Mohammed, A., & Bereket, D. (2019). *Knowledge and practice on adequate sunlight exposure of infants among mothers attending EPI unit of Aleta Wondo Health Center, SNNPR, Ethiopia. BMC Research Notes*, 12, 183.
16. [^]Melika, G. (2020). *Knowledge, Attitude and Practice of Mothers on Sun Exposure of their Infants at St. Paul's Hospital, Ethiopia. International Journal of Advanced Studies in Computers, Finance and Engineering*, 6(1), 1-1.
17. [^]Wegayehu, Z., Leul, D., Tewodros, T., Abebe, M., Michael, A., Ababayehu, M., Kassa, M., & Mikyas, A. (2021). *Assessment of Practice and Factors associated with Sunlight Exposure of Infants among Mothers in Debre Berhan Town, North Shewa Zone, Amhara Region, Ethiopia. Pediatric Health, Medicine and Therapeutics*, 12, 507-517.
18. [^]Rajalakshmi, M., & Abebe, A. (2013). *Assessment of Knowledge and Factors Affecting Sunlight Exposure of Infants among Mothers Attending Governmental Health Facilities in Debre Markos Town, Ethiopia. International Journal of Science and Research*, 2(12), Article 2319-7064.
19. [^]Amare, B., Abebaw, M., Bayachew, A., Yidersal, H., & Habtamu, T. (2022). *Practice of sunlight exposure of infants and associated factors among infant coupled mothers at Dejen District, Amhara Region, Northwest Ethiopia 2021. Nutrition and Metabolic Insights*, 15, 1-7.
20. ^{a, b, c}Haileyesus, G., & Yilkal, T. (2019). *Assessment of knowledge, attitude, and practice of sunlight exposure of infants among mothers attending in governmental health facilities in Farta District, South Gondar Zone, North West Ethiopia, 2018. International Journal of Reproductive Medicine.* <https://doi.org/10.1155/2019/2638190>

