

Robin's criterion on superabundant numbers

Frank Vega

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Robin's criterion states that the Riemann hypothesis is true if and only if the inequality $\sigma(n) < e^\gamma \cdot n \cdot \log \log n$ holds for all natural numbers $n > 5040$, where $\sigma(n)$ is the sum-of-divisors function of n , $\gamma \approx 0.57721$ is the Euler-Mascheroni constant and \log is the natural logarithm. We require the properties of superabundant numbers, that is to say left to

right maxima of $n \mapsto \frac{\sigma(n)}{n}$. Let P_n be equal to $\prod_{q|n} \frac{q^{v_q(n)+2}-1}{6 \cdot q^{v_q(n)+2}}$ for a superabundant number $n > 5040$, where $v_p(n)$ is the p -adic order of n , q_k is the largest prime factor of n and $N_r = \prod_{i=1}^r q_i$ is the largest primorial number of order r such that $\frac{N_r}{6} < q_k^2$. In this note, we prove that the Riemann hypothesis is true when $P_n \geq Q$ holds for all large enough

$$\frac{1.2 \cdot (2 - \frac{1}{8}) \cdot (3 - \frac{1}{3})}{(2 - \frac{1}{2^{19}}) \cdot (3 - \frac{1}{3^{12}})}$$

superabundant numbers n , where $Q = \frac{1.2 \cdot (2 - \frac{1}{8}) \cdot (3 - \frac{1}{3})}{(2 - \frac{1}{2^{19}}) \cdot (3 - \frac{1}{3^{12}})} \approx 1.0000015809$. In particular, the inequality $P_n \geq Q$ holds when

$\sum_{q|m} \frac{m}{q^{\sigma(q^{v_q(n)+1})}} \geq \sigma(m) \cdot \log Q$ also holds such that $m = \prod_{q|n} N_r q^{v_q(n)+1}$ since $\sigma(\dots)$ is multiplicative.

Frank Vega

Research Department, NataSquad, 10 rue de la Paix, Paris, 75002, France

vega.frank@gmail.com

Keywords: Riemann hypothesis, Robin's inequality, Superabundant numbers, Sum-of-divisors function, Prime numbers.

MSC Classification: 11M26, 11A41, 11A25

1. Introduction

The hypothesis was proposed by Bernhard Riemann (1859). The Riemann hypothesis belongs to the Hilbert's eighth problem on David Hilbert's list of twenty-three unsolved problems. As usual $\sigma(n)$ is the sum-of-divisors function of n

$$\sum_{d|n} d,$$

where $d | n$ means the integer d divides n . Define $f(n)$ as $\frac{\sigma(n)}{n}$. We say that Robin(n) holds provided that

$$f(n) < e^\gamma \cdot \log \log n,$$

where $\gamma \approx 0.57721$ is the Euler-Mascheroni constant and \log is the natural logarithm. The Ramanujan's Theorem states that if the Riemann hypothesis is true, then the previous inequality holds for large enough n [1]. Next, we have the Robin's Theorem:

Proposition 1. Robin(n) holds for all natural numbers $n > 5040$ if and only if the Riemann hypothesis is true [2], Theorem 1 pp. 188].

In 1997, Ramanujan's old notes were published where it was defined the generalized highly composite numbers, which include the superabundant and colossally abundant numbers [1]. Let $q_1 = 2, q_2 = 3, \dots, q_k$ denote the first k consecutive primes, then an integer of the form $\prod_{i=1}^k q_i^{a_i}$ with $a_1 \geq a_2 \geq \dots \geq a_k \geq 1$ is called a Hardy-Ramanujan integer [3], pp. 367]. A natural number n is called superabundant precisely when, for all natural numbers $m < n$

$$f(m) < f(n).$$

We know the following properties for the superabundant numbers:

Proposition 2. If n is superabundant, then n is a Hardy-Ramanujan integer [4], Theorem 1 pp. 450].

Proposition 3. [4], Theorem 7 pp. 454]. Let n be a superabundant number such that p is the largest prime factor of n , then

$$p \sim \log n, \quad (n \rightarrow \infty).$$

Proposition 4. [4], Theorem 9 pp. 454]. The number of superabundant numbers less than x exceeds

$$\frac{c \cdot \log x \cdot \log \log x}{(\log \log \log x)^2}.$$

A number n is said to be colossally abundant if, for some $\epsilon > 0$,

$$\frac{\sigma(n)}{n^{1+\epsilon}} \geq \frac{\sigma(m)}{m^{1+\epsilon}} \quad \text{for } (m > 1).$$

There is a close relation between the superabundant and colossally abundant numbers

Proposition 5. Every colossally abundant number is superabundant [4], pp. 455].

Several analogues of the Riemann hypothesis have already been proved. Many authors expect (or at least hope) that it is true. However, there are some implications in case of the Riemann hypothesis might be false.

Proposition 6. If the Riemann hypothesis is false, then there are infinitely many colossally abundant numbers $n > 5040$

such that $\text{Robin}(n)$ fails (i.e. $\text{Robin}(n)$ does not hold) [2], Proposition pp. 204].

The following is a key Corollary.

Corollary 1. *If the Riemann hypothesis is false, then there are infinitely many superabundant numbers n such that $\text{Robin}(n)$ fails.*

Proof. This is a direct consequence of Propositions 1,5 and 6.□

In number theory, the p -adic order of an integer n is the exponent of the highest power of the prime number p that divides n . It is denoted $v_p(n)$. Equivalently, $v_p(n)$ is the exponent to which p appears in the prime factorization of n .

Proposition 7. $\text{Robin}(n)$ holds for all natural numbers $n > 5040$ such that $v_2(n) \leq 19$ and $v_3(n) \leq 12$ [5], Theorem 1 pp. 2, Theorem 2 pp. 2].

Proposition 8. [4], Theorem 5 pp. 452]. Let n be a superabundant number such that $v_q(n) = t$, p is the largest prime factor of n , $2 \leq q \leq p$ and $q < (\log p)^\alpha$, where α is a constant, then

$$\log \frac{q^{t+2} - 1}{q^{t+2} - q} < \frac{\log q}{p \cdot \log p} \cdot \left(1 + O\left(\frac{(\log \log p)^2}{\log p \cdot \log q} \right) \right).$$

This is the main insight.

Lemma 1. *Let n be a large enough superabundant number such that $p > 3$ is the largest prime factor of n , then*

$$p < 2^{v_2(n) - 19}$$

and

$$p < 3^{v_3(n) - 12}.$$

Let P_n be equal to $\prod_{q|n} \frac{q^{v_q(n)+2} - 1}{q^{v_q(n)+2} - q}$ for a superabundant number $n > 5040$, where q_k is the largest prime factor of n and

$N_r = \prod_{i=1}^r q_i$ is the largest primorial number of order r such that $\frac{N_r}{6} < q_k^2$. Putting all together yields the main theorem:

Theorem 1. *The Riemann hypothesis is true when $P_n \geq Q$ holds for all large enough superabundant numbers n , where*

$Q = \frac{1.2 \cdot (2 - \frac{1}{8}) \cdot (3 - \frac{1}{3})}{(2 - \frac{1}{2^{19}}) \cdot (3 - \frac{1}{3^{12}})} \approx 1.0000015809$. In particular, the inequality $P_n \geq Q$ holds when $\sum_{q|m} \sigma\left(\frac{m}{q^{v_q(n)+1}}\right) \gtrsim \sigma(m) \cdot \log Q$ also holds such that $m = \prod_{q|n} N_r q^{v_q(n)+1}$ since $\sigma(\dots)$ is multiplicative.

2. Proof of the Lemma 1

Proof. Let $q \in \{2, 3\}$ and $v_q(n) = t$. For every large enough superabundant number n , there is a constant α such that $q < (\log p)^\alpha$. For example, we can take $\alpha = 2.5$ since $(\log p)^{2.5} \geq (\log 5)^{2.5} > 3$. We will use the following inequality

$$\frac{u}{u+1} < \log(1+u), \quad (u > 0).$$

From the previous inequality, we notice that

$$\begin{aligned} \log \frac{q^{t+2} - 1}{q^{t+2} - q} &= \log \left(1 + \frac{q-1}{q^{t+2} - q} \right) \\ &> \frac{\frac{q-1}{q^{t+2} - q}}{\frac{q-1}{q^{t+2} - q} + 1} \\ &= \frac{q-1}{(q^{t+2} - q) \cdot \left(\frac{q-1}{q^{t+2} - q} + 1 \right)} \\ &= \frac{q-1}{(q-1) + (q^{t+2} - q)} \\ &= \frac{q-1}{q^{t+2} - 1} \\ &> \frac{1}{3 \cdot q^{t+1}}. \end{aligned}$$

Hence, there is a constant $C > 0$ such that

$$q^t > C \cdot \frac{p \cdot \log p}{\log q}$$

by Proposition 8. Putting $c = \frac{C}{\log q}$, then we obtain that

$$c \cdot p \cdot \log p < q^t,$$

where c is a positive constant. We deduce that

$$c \cdot \log p > 3^{12}$$

by Proposition 3 for large enough n . Therefore, the proof is done. \square

3. Proof of the Theorem 1

Proof. There are infinitely many superabundant numbers by Proposition 4. Let $n > 5040$ be a large enough superabundant number. Let $\prod_{i=1}^k q_i^{a_i}$ be the representation of this superabundant number n as the product of the first k consecutive

primes $q_1 < \dots < q_k$ with the natural numbers $a_1 \geq a_2 \geq \dots \geq a_k \geq 1$ as exponents, since n must be a Hardy-Ramanujan

integer by Proposition 2. Let P_n be equal to $\prod_{q|n} \frac{q^{v_q(n)+2}-1}{6 \cdot q^{v_q(n)+2-q}}$ for $n > 5040$, where $N_r = \prod_{i=1}^r q_i$ is the largest primorial

number of order r such that $\frac{N_r}{6} < q_k^2$. Suppose that Robin(n) fails and $P_n \geq Q$, where $Q = \frac{1.2 \cdot (2^{-\frac{1}{8}}) \cdot (3^{-\frac{1}{3}})}{(2^{-\frac{1}{2^{19}}}) \cdot (3^{-\frac{1}{3^{12}}})} \approx 1.0000015809$.
So,

$$f(n) \geq e^{\gamma} \cdot \log \log n.$$

We know that

$$\begin{aligned} f(n) &= f(2^{v_2(n)} \cdot 3^{v_3(n)}) \cdot f\left(\frac{n}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right) \\ &< 3 \cdot f\left(\frac{n}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right) \\ &= f(2^3 \cdot 3 \cdot 5) \cdot f\left(\frac{n}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right) \\ &\leq f\left(\frac{2^{19} \cdot 3^{12} \cdot n \cdot \frac{N_r}{6}}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right) \\ &= f\left(\frac{n \cdot \frac{N_r}{6}}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}}\right) \end{aligned}$$

since $P_n \geq Q$, $\frac{q_i}{q_i-1} > \frac{q_i^{a_i+1}-1}{q_i^{a_i} \cdot (q_i-1)} = f(q_i^{a_i})$ and $f(\dots)$ is multiplicative, where

$f(2^3 \cdot 3 \cdot 5) = 3 = 2 \cdot \frac{3}{2} > f(2^{v_2(n)}) \cdot f(3^{v_3(n)}) = f(2^{v_2(n)} \cdot 3^{v_3(n)})$. This is true because of

$$f(2^3 \cdot 3 \cdot 5) \cdot f\left(\frac{n}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right) \leq f\left(\frac{2^{19} \cdot 3^{12} \cdot n \cdot \frac{N_r}{6}}{2^{v_2(n)} \cdot 3^{v_3(n)}}\right)$$

is equivalent to say that

$$\frac{f(2^3 \cdot 3 \cdot 5)}{f(2^{19} \cdot 3^{12})} \leq q \mid \prod_{q \mid \frac{N_r}{6}} \frac{f(q^{v_q(n)+1})}{f(q^{v_q(n)})}$$

Certainly, we know that

$$\frac{f(2^3 \cdot 3 \cdot 5)}{f(2^{19} \cdot 3^{12})} = Q$$

and

$$\prod_{q \mid \frac{N_r}{6}} \frac{f(q^{v_q(n)+1})}{f(q^{v_q(n)})} = q \mid \prod_{q \mid \frac{N_r}{6}} \frac{q^{v_q(n)+2} - 1}{q^{v_q(n)+2} - q} = P_n$$

Consequently, that is true under the supposition that $P_n \geq Q$. We have

$$f \left(\frac{n \cdot \frac{N_r}{6}}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right) < e^Y \cdot \log \log \left(\frac{n \cdot \frac{N_r}{6}}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right)$$

by Proposition 7. Therefore, we obtain that

$$e^Y \cdot \log \log \left(\frac{n \cdot \frac{N_r}{6}}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right) > e^Y \cdot \log \log n$$

which is the same as

$$\left(\frac{n \cdot q_k^2}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right) > \left(\frac{n \cdot \frac{N_r}{6}}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right) > n$$

using the inequality $\frac{N_r}{6} < q_k^2$. However, we know that

$$2^{v_2(n)-19} > q_k$$

and

$$3^{v_3(n)-12} > q_k$$

by Lemma 1, due to n is large enough. So, we can see that necessarily,

$$\left(\frac{n \cdot q_k^2}{2^{v_2(n)-19} \cdot 3^{v_3(n)-12}} \right) < n.$$

In this way, we obtain a contradiction under the assumption that Robin(n) fails and $P_n \geq Q$, where

$Q = \frac{1.2 \cdot (2^{-\frac{1}{8}}) \cdot (3^{-\frac{1}{3}})}{(2^{-\frac{1}{2^{19}}}) \cdot (3^{-\frac{1}{3^{12}}})} \approx 1.0000015809$. To sum up, the study of this arbitrary large enough superabundant number n reveals that Robin(n) holds whenever $P_n \geq Q$. Accordingly, Robin(n) holds for all large enough superabundant numbers n when $P_n \geq Q$ holds. This contradicts the fact that there are infinitely many superabundant numbers n , such that Robin(n) fails when the Riemann hypothesis is false according to Corollary 1. By reductio ad absurdum, we prove that the Riemann hypothesis is true when $P_n \geq Q$ holds for all large enough superabundant numbers n . From the proof of the Lemma 1, we show that $\log \frac{q^{v_q(n)+2}-1}{q^{v_q(n)+2}-q} > \frac{q-1}{q^{v_q(n)+2}-1}$, where we know that $\frac{q-1}{q^{v_q(n)+2}-1} = \frac{1}{\sigma(q^{v_q(n)+1})}$. Thus, the inequality $P_n \geq Q$ holds when $\sum_{q|n} \frac{N_q}{6} \sigma\left(\frac{n'}{q^{v_q(n)+1}}\right) \geq \sigma(n') \cdot \log Q$ also holds such that $n' = \prod_{q|n} \frac{N_q}{6} q^{v_q(n)+1}$ since $\sigma(\dots)$ is multiplicative. However, the inequality

$$\sum_{q|n} \frac{N_q}{6} \frac{n'}{\sigma(q^{v_q(n)+1})} \geq \sigma(n') \cdot \log Q$$

is the same as

$$\sum_{q|m} \frac{N_q}{6} \frac{m}{\sigma(q^{v_q(n)+1})} \geq \sigma(m) \cdot \log Q$$

after multiplying both sides by $\sigma(2^{v_2(n)+1} \cdot 3^{v_3(n)+1})$ such that $m = n' \cdot 2^{v_2(n)+1} \cdot 3^{v_3(n)+1}$. That would be the same as

$$\begin{aligned} \sum_{q|m} \frac{m}{\sigma(q^{v_q(n)+1})} &\geq \sigma(m) \cdot \log Q + \frac{\sigma(m)}{\sigma(2^{v_2(n)+1})} + \frac{\sigma(m)}{\sigma(3^{v_3(n)+1})} \\ &= \sigma(m) \cdot \left(\log Q + \frac{1}{\sigma(2^{v_2(n)+1})} + \frac{1}{\sigma(3^{v_3(n)+1})} \right) \\ &\cong \sigma(m) \cdot \log Q \end{aligned}$$

and thus, the proof is done. \square

References

1. [a](#), [b](#) J.L. Nicolas, G. Robin, *Highly Composite Numbers by Srinivasa Ramanujan*. *The Ramanujan Journal* 1(2), 119–

- 153 (1997). <https://doi.org/10.1023/A:1009764017495>
2. ^{a, b}G. Robin, *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann*. *J. Math. pures appl* 63(2), 187–213 (1984)
 3. [^]Y. Choie, N. Lichiardopol, P. Moree, P. Solé, *On Robin's criterion for the Riemann hypothesis*. *Journal de Théorie des Nombres de Bordeaux* 19(2), 357–372 (2007). <https://doi.org/10.5802/jtnb.591>
 4. ^{a, b, c, d, e}L. Alaoglu, P. Erdős, *On Highly Composite and Similar Numbers*. *Transactions of the American Mathematical Society* 56(3), 448–469 (1944). <https://doi.org/10.2307/1990319>
 5. [^]A. Hertlein, *Robin's Inequality for New Families of Integers*. *Integers* 18 (2018)