

1 **Determination of Evapotranspiration and Crop Coefficients of Irrigated Legumes on**
2 **Different Soil Textures Using the FAO56 Approach**

3 *Sasanya, Blessing Funmbi¹, Udoh, Divine-Joy Imaobong¹, Sasanya, Rashida¹

4 blessing.env@gmail.com,

5 ¹Department of Crop and Soil Science, Faculty of Agriculture, University of Port Harcourt,
6 Port Harcourt, Nigeria.

7 *Corresponding Author

8 **Abstract**

9 Climates and locations specific crop coefficients (K_c) plays pivotal roles in effective water
10 management and crop yield optimization. Research on K_c for many staple crops remains limited
11 in many regions across Africa. This study examined the crop coefficients of two varieties of
12 cowpea (Sampea 14 and Sampea 17) and a variety of soybean (TGX 1465-1D) across different
13 soil textures for optimized water management and crop yield. This study ascertained the K_c of
14 Sampea14 (V1), Sampea17 (V2), and TGX 1465-1D (V3) cultivated on sandy, sandy clay loam,
15 and sandy loam soils, within Port Harcourt, Southern Nigeria. This formed a 3 by 3 factorial
16 experiment in three replicates. Changes in soil water content and crops' agronomic parameters
17 were monitored regularly. Meteorological data were obtained from the National Aeronautic and
18 Space Administration website and reference evapotranspiration were estimated from the
19 meteorological data using the FAO Penman Monteith's method. K_c were estimated from the
20 actual crop water use and reference crop evapotranspiration using the FAO56 method. During
21 the initial-growth stage, the average K_c for V1, V2, and V3 were computed as 0.55, 0.72, and
22 0.81, respectively. At the mid-growth stage, these coefficients increased to 1.48, 1.30, and 1.60
23 and at the late season, the average K_c were 0.80, 0.69, and 0.86, for the respective varieties. The
24 findings from this study hold the potential to significantly contribute to proper water
25 management practices in the studied area. The understanding of K_c for these specific legume
26 varieties across different soil textures offers valuable insights for optimizing irrigation
27 scheduling and ultimately enhancing crop yield and water productivity.

28 **Keywords:** Crop coefficients, Irrigation, Legumes, Soil textures, Water consumption.

29 **1.0 Introduction**

30 Legumes are cost-effective nutrient sources and potential income generators for
31 subsistent farmers lacking access to costly irrigation and fertilizers. Legumes serve as cover
32 crops to mitigate soil erosion and demonstrate remarkable resilience to both water stress and
33 waterlogging throughout their growth cycle (Poyen et al., 2016). Furthermore, their symbiotic
34 relationship with nitrogen-fixing Rhizopus in root nodules renders them valuable in crop
35 rotation (Kalidass and Mahapatra, 2014). According to National Agricultural Extension
36 Research Services (NAERLS) (2010) and the National Programme on Agriculture and Food
37 Security (NPAFS) (2010), the total production of cowpea and soybean were 1,561,964 and
38 491,504 t/ha, respectively, between 2007 and 2011. This implies that one of the foci of the
39 Nigerian nation in the agricultural sector is to engage in an intensified legume production with
40 optimal inputs and management.

1 Evapotranspiration is the total water loss from plants due to transpiration and soil
2 surface evaporation, influenced by factors such as temperature, humidity, sunlight, wind, and
3 vegetation cover (Nouri et al., 2016). Water regularly evaporates from various surfaces,
4 including lakes, rivers, pavements, soils, and wet vegetation (Dehghanisanij and Kosari, 2011).
5 Different crops also have distinct water requirements or transpiration requirement at different
6 growth stages, primarily determined by crop coefficients (K_c), which play a crucial role in
7 estimating crop water needs (Rallo et al., 2021; Drechsler, 2022). Crop coefficients are inherent
8 crop characteristics used to calculate crop evapotranspiration or water requirements (ET_c)
9 (Chandra and Kumari, 2021). They compare the evapotranspiration of the specific crop to a
10 well-calibrated reference crop under identical conditions. Crop Water Requirement (CWR)
11 (mm) measures the evapotranspiration when soil water is adequate due to precipitation or
12 irrigation, supporting unhindered plant growth and yield (Djaman et al., 2018). It also ensures
13 water remains within the root system's absorption capacity (Silva et al., 2017). According to
14 Sultana et al. (2022), CWR represents the water depth required to fulfil ET_c by a disease-free
15 crop under unrestricted soil conditions, including water and fertility, to achieve maximum
16 production potential in the given environment. Factors such as daily temperature, relative
17 humidity, and wind velocity influence the CWR (Khan et al., 2021; Sepaskhah and Razzaghi,
18 2022). Consequently, crops in hot, dry, windy, and sunny climates typically have the highest
19 water requirements (Djaman et al., 2018).

20 The crop coefficient (K_c) is typically determined through empirical experimentation
21 and encompasses the cumulative impact of multiple factors such as changes in leaf area, plant
22 height, crop attributes, irrigation methods, developmental rate, planting date, canopy cover,
23 canopy resistance, soil and climatic conditions, and agricultural management practices
24 (Shenkut et al., 2013). Each crop is associated with a specific set of K_c values, which project
25 varying water utilization across different crops during distinct growth stages (Pereira et al.,
26 2021). The duration of these stages is contingent on local climate, latitude, elevation, planting
27 date, crop type, and agricultural techniques. The most precise determination of a crop's growth
28 stage, and subsequent adjustment of empirical K_c values, is achieved through on-site
29 observations (Kenjabaev et al., 2020). According to Pereira et al. (2021), during the initial
30 stages of crop germination and establishment, the majority of evapotranspiration (ET_c)
31 primarily comprises soil surface evaporation, resulting in lower crop water use rates and
32 smaller K_c values (referred to as the K_c initial stage). As the crop matures, with its canopy
33 expanding to cover the soil surface, soil surface evaporation diminishes, and the
34 transpiration component of ET rises. Agronomical plants reach their maximum ET_c rate when

1 fully developed (referred to as the K_c mid-season), but as the season nears its end and the plant
2 reaches physiological maturity, the ET rate decreases (referred to as the K_c end season) (Pereira et
3 al., 2021).

4 According to Wang et al. (2018) and Fan et al. (2021), crop water consumption is a
5 crucial factor in assessing the viability of crop cultivation in any given region. In recent times,
6 crop coefficients of various crops have been investigated using various methods. For instance,
7 Hassan et al. (2022) utilized remote sensing and meteorological data to monitor plant
8 phenology and estimate crop coefficients and evapotranspiration. They reported a strong
9 coefficient of determination (0.98) between field crop coefficients derived from meteorological
10 data and those predicted from an NDVI value of 0.147. In another study, Dingre et al. (2020)
11 investigated the relationship between field water balance-derived crop coefficients and canopy
12 reflectance-based NDVI for irrigated sugarcane. This relationship was described by a second-
13 order polynomial regression equation. Similarly, Puig-Sirera et al. (2021) determined the
14 transpiration and water use of a traditional olive grove under irrigation using the sap flow
15 method and the FAO56 dual crop coefficient approach. Their findings indicated that the basal
16 crop coefficient (K_{cb}) for the mid-season of the olive6 grove ranged from 0.40 to 0.45, while
17 at the end of the season, K_{cb} ranged between 0.35 and 0.40.

18 In the context of Tangara da Serra, MT, Brazil, Bariviera et al. (2020) determined the
19 dual crop coefficient for an early cycle soybean cultivar using a precise lysimeter. They
20 reported K_{cb} and soil evaporation (K_e) values of 0.47, 1.15, 0.89 and 0.94, 0.14, 0.44 for the
21 early, mid, and late growth stages, respectively. Additionally, Gong et al. (2020) conducted
22 research on tomato cultivation in a solar greenhouse under both full and deficit irrigation
23 conditions. They observed maximum crop evapotranspiration ranging from 0.15 to 1.88 mm/h
24 under full irrigation and 0.15 to 0.89 mm/h under deficit irrigation. The average daily standard
25 evapotranspiration was 5.11 mm/day, with a seasonal evapotranspiration of 1814 mm. The crop
26 coefficients for citrus trees in the study of Jamshidi et al, 2020, ranged from 0.67 to 0.96. Lastly,
27 in the semi-arid climate of the Senegal River Valley, average crop coefficient values of 1.01,
28 1.31 and 1.12 were reported for rice during the crop development, mid-season, and late-season
29 growth stages (Djaman et al., 2019).

30 In many regions across Africa, comprehensive research on crop coefficients for key
31 staple crops remains limited due to the diverse climatic variations (Poyen et al., 2016). This
32 study was thus designed to examine the crop coefficients of specific legumes across different
33 soil textures for optimized water management and crop yield. The primary objective of this
34 study is to determine the crop coefficients (K_c) of three distinct legumes species—Sampea 14

1 (V1), Sampea 17 (V2) of cowpea, and TGX 1465-1D (V3) of soybean. The FAO56 approach
2 was employed to determine the K_c values from the directly measured actual crop water use and
3 estimated reference evapotranspiration. According to Cardova et al. (2015) the FAO56 method
4 is of great benefits, because it provides structures for standardizing potential
5 evapotranspiration. This research is conducted across various soil textures including sandy,
6 sandy clay loam, and sandy loam soils. The investigation spans multiple growth stages,
7 encompassing emergence (initial), flowering (development), pod-setting (mid), and maturity
8 (late), all within the prevailing climatic conditions of Port Harcourt, located in southern
9 Nigeria. Results from this study would enable irrigation planning and water management in the
10 cultivation of different varieties of soybean and cowpea on different soil textures.

11 **2.0 Materials and Methods**

12 **2.1 Study Area**

13 This study was conducted at the agricultural research station within the Department of
14 Crop and Soil Science, situated in the Faculty of Agriculture at the University of Port Harcourt.
15 This facility is located at coordinates 4.847°N latitude and 6.975°E longitude, as illustrated in
16 Figure 1. It is positioned in the Obio/Akpor Local Government Area of Rivers State, Nigeria,
17 at an elevation of approximately 15.85 meters above mean sea level. The climate in Port
18 Harcourt is predominantly humid, characterized by an average annual precipitation of 2,293
19 mm, and the mean annual temperature and relative humidity levels are approximately 28°C
20 and 75%, respectively.

21 **2.2 Experimental Design**

22 The study was designed as a 3 x 3 factorial experiment, with three replications within
23 a controlled environment. Twenty-seven experimental plots (each with 4 m² size) were used for
24 the experiment while considering an effective root zone depth of 0.5 m. The experimental
25 treatments consisted of three different varieties of leguminous crops, namely two cowpea types
26 (Sampea 14 and Sampea 17) and one type of soybean (TGX 1465-1D). The experiment also
27 considered three distinct soil textures, specifically sandy, sandy clay loam and sandy loam soils.
28 Detailed descriptions are shown on Figure 2. Some soil movements from other sites were done
29 to ensure planting beds were made from the specified soil textures.

30




Figure 1: Map of Rivers State, Delineating Obio-Akpor Local Government, Nigeria




Figure 2: Detailed Description of the Experimental Set-up

2.3 Physicochemical Properties of Experimental Soils

The bulk densities of the sand, sandy clay and sandy clay loam soils also indicated the level of compaction which influences the soil water holding capacity, water movement, plant water availability and crop rooting depth (Jabro et al., 2020). Investigation on bulk densities, hydraulic conductivities and particle size analysis were carried out on collected soil samples. The saturated hydraulic conductivity was investigated using the method specified by Reynolds et al., (2002). Soils' bulk densities were determined according to Blake and Hartage method (1986). Soil's particle size distributions were analysed to determine the relative proportion of sand, silt, and clay in the soils using the hydrometer method as described by Gavlack et al. (2005).

The investigation on total nitrogen, total phosphorus, cation exchange capacity, calcium, magnesium and potassium content were carried out on collected soil samples to determine the soil nutrient level and to inform the level of needed fertilizer application to the crops. The total nitrogen was determined according to the O'Dell (1993) method. The total phosphorous was determined using the colorimetric, ascorbic acid, two reagents (1978) method. The Cation Exchange Capacity (CEC) is a measure of the number of ions that can be adsorbed in an exchangeable fashion, on the negative charge sites of the soil (Bache, 1976). The CEC was determined according to Bache (1976) method which is known as the ammonium acetate extraction method. Calcium, magnesium and potassium were determined using the calcium chloride extraction method described by Houba et al. (2000).

2.4 Planting and Crop Management

Three different varieties of leguminous crops were planted on March 15, 2022. These varieties included Sampea 14 (V1), and Sampea 17 (V2), TGX 1465-1D (V3). The planting depth was 2cm below the soil surface. Each legume variety was planted on three distinct soil textures in three replicates, resulting in a total of 27 experimental plots with nine plots per replicate. The experiment lasted for 15 weeks and concluded on July 18, 2022. The experiment's duration was divided into specific growth stages (emergence, flowering, pod setting, and maturity) as defined by Allen et al. (1998). During the experiment, water use and crop coefficients were estimated at three stages: the initial (15 days for soybeans and 20 days for cowpea varieties), mid-season (55 days for soybeans and 60 days for cowpea varieties), and late stages (15 days for soybeans and 20 days for cowpea varieties) (Allen et al., 1998)

To enhance the nutrient content of the collected soil samples, granular water-soluble fertilizers with a ratio of N: P: K at 6: 25: 5 were applied to the soil surface before planting, following the recommended agronomic rate of 50kg/ha based on the soil chemical analysis.

1 Fertilizer application was repeated 21 days after planting to promote crops' healthy
 2 development. Weed control measures were implemented to maintain disease-free plants and
 3 prevent competition with the main crops for nutrients and water. Throughout the crops' growth
 4 period efforts were made to prevent water stress. Irrigation water was consistently applied
 5 manually using watering can, whenever the soil water content reduced to 40% of the soil's field
 6 capacity. Irrigation water was to return the soil back to 100% field capacity.

7 **2.5 Agrometeorological Data Collection**

8 Daily remotely sensed agrometeorological data for the experimental location (4.847 °N,
 9 6.975 °E) were obtained from the National Aeronautics and Space Administration's (NASA)
 10 website (<https://power.larc.nasa.gov/data-access-viewer/>). Data collected include daily
 11 minimum and maximum temperature, daily maximum and minimum relative humidity, daily
 12 average wind velocity and mean hours of daily sunlight. Collected data spanned from the 1st of
 13 March 1981 to 31st July 2022. These data were employed for the estimation of reference
 14 evapotranspiration throughout the growth period of planted crops.

15 **2.6 Estimation of Reference Evapotranspiration (ET_p)**

16 Among all methods, the Penman-Monteith equation has been recommended by the
 17 Food and Agriculture Organisation (FAO) as the standard method for the computation of ET_p
 18 especially under arid conditions (Allen et al., 1998). Adesogan and Sasanya (2023) also
 19 recommended the method, since the Reference Evapotranspiration (ET_p) values obtained from
 20 it were closely related to measured ET_p values from a class A evaporation pan.

21 The reference evapotranspiration (ET_p) at the growth stages of the crops was thus
 22 computed for the growing period by the Penman-Monteith Method. The FAO-56 recommended
 23 method was chosen for its accuracy in the estimation of ET_p from climatological data
 24 (Adesogan and Sasanya, 2023). Daily ET_p was estimated from Equations 1 to 11.

$$25 \quad ET_p = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T+273} U_2 (e_s - e_a)}{\Delta + \gamma (1 + 0.34 U_2)} \quad (1)$$

$$26 \quad R_s = \left(a_s + b_s \frac{n}{N} \right) R_a \quad (2)$$

$$27 \quad R_{so} = (0.75 + 2 z * 10^{-5}) R_a \quad (3)$$

$$28 \quad R_{ns} = (1 - \alpha) R_s \quad (4)$$

$$29 \quad R_{nl} = \sigma \left[\frac{(T_{maxk})^4 + (T_{mink})^4}{2} \right] (0.34 - 0.14 \sqrt{e_a}) \left(1.35 \frac{R_s}{R_{so}} - 0.35 \right) \quad (5)$$

$$30 \quad G = 0.14 (T_{MONTH i} - T_{MONTH i-1}) \quad (6)$$

1 Where: R_n = Average net radiation at the crop surface (MJ/m²/day), G is soil flux density
2 (MJ/m²/day), γ = Psychrometric constant (kPa/ °C) = 0.067, U_2 = Wind velocity at 2m height (m/s),
3 e_s = Saturation vapor pressure (kPa), e_a = Actual vapor pressure (kPa), $e_s - e_a$ = Saturation vapor
4 pressure deficit (kPa), T = air temperature at 2m height (°C), Δ = Slope vapor pressure curve (kPa/
5 °C).
6 R_s is the incoming solar radiation; obtained from the relationship between angstrom values $a_s = 0.25$,
7 $b_s = 0.5$ and R_a . (equation 1b). R_{so} is the clear sky solar radiation obtained from elevation (z m) and
8 R_a (equation 1c). R_{ns} is the shortwave radiation; calculated from albedo (α) and R_s (equation 1e). R_{nl}
9 was estimated from equation (1e). σ is Stefan Boltzmann constant = 4.903×10^{-9} MJ/ K⁴/m²/day,
10 T_{maxk} and T_{mink} are absolute values of monthly minimum and maximum temperatures. R_n was
11 calculated as the difference between the net long wave radiation (R_{nl}) and net shortwave radiation
12 (R_{ns}) measured in MJ/m²/day. The soil flux G from obtained from equation 1f. $T_{MONTH i}$ is the mean
13 temperature of the present month and $T_{MONTH i-1}$ is the mean temperature of the previous month in
14 °C. 0.408 is a constant which converts the unit MJ/m²/day to mm/day.

$$15 \quad e_s = e^o(T_{mean}) = 0.611 \exp\left(\frac{17.27T_{mean}}{T_{mean}+237.3}\right) \quad (7)$$

$$16 \quad \Delta = \frac{4098e_s}{(T+237.3)^2} \quad (8)$$

$$17 \quad e_{a1} = \frac{e^o(T_{min}) \times RH_{max}}{100} \quad (9)$$

$$18 \quad e_{a2} = \frac{e^o(T_{max}) \times RH_{min}}{100} \quad (10)$$

$$19 \quad e_a = \frac{e_{a1} + e_{a2}}{2} \quad (11)$$

20 Where: RH_{max} = Maximum relative humidity (%), RH_{min} = Minimum relative humidity (%).
21 T_{min}, T_{max} and T_{mean} are the monthly minimum, maximum and mean temperature in °C.

22 **2.7 Collection of Water Use and Agronomic Data**

23 Depletion in the soil water content and changes in the agronomic properties of the
24 planted crops were monitored and measured on a weekly basis. These enabled the monitoring
25 of evapotranspiration rate and changes in plant development.

26 **2.7.1 Monitoring Soil Moisture Depletion**

27 The soil moisture was monitored from the water balance Equation (Equation 12).
28 Depletion of soil moistures were monitored using the gravimetric method as employed by Alla
29 Jabow et al. (2015) and Djaman et al. (2019).

$$ET_c = I + R_a + G - R - D \mp \Delta S \quad (12)$$

Where: I is Irrigation, R_a is the rainfall, G is capillary water, R is runoff, D is deep drainage, ΔS is the change in soil moisture. 'I' was considered to be equal to zero before and after irrigation as at when needed, R_a is zero during the cultivation period since the experiment was done in a controlled environment, the groundwater table in the study area is not close to the soil surface, therefore capillary rise from groundwater G was zero. Runoff and deep drainage were zero, since the right amount of irrigation water was applied without permitting excesses.

The change in soil moisture (ΔS) were estimated by collecting soil samples from the effective root zone, weekly, in replicates, from each experimental plots, by means of the soil cores samplers. Soil samples in soil core samplers were oven dried for 24 hours at 105°C and the dry weight were determined afterwards. These were done before and after water application. Weekly depletion of soils was determined from Equation 13.

$$\Delta S = \frac{SWC1 - SWC2}{\Delta t} \quad (13)$$

Where: ΔS is the gravimetric weekly water depletion (g/g), SWC1= Soil water content at week 1 (g/g), SWC2 = Soil water content at week 2 (g/g), Δt is the change in time.

The gravimetric water content was converted to volumetric water content using Equation 14. The volumetric water depletion is equivalent to the actual evapotranspiration (ET_c) of water used by the crops. The seasonal ET_c was estimated from the total ET_c over the cultivation period from planting to harvesting.

$$\Delta d = 1000 \times \rho \times \Delta S \quad (14)$$

Where: Δd = change in soil water depth (mm/m), ρ = dry bulk density (g/cm³), ΔS = Change in gravimetric moisture content (g/g).

2.7.2 Crop Data Collection

Agronomic data including plant height, leaf length, leaf width, number of leaves and leaf area index were measured weekly starting from the 7th day after planting. The leaf area index (LAI) is the green leaf area of plants per unit area. The plant height (cm), leaf length (cm) and leaf width (cm) were measured weekly using a meter rule. The LAI was estimated from Equation 15.

$$LAI = \frac{a \times LL \times LW}{CA} \quad (15)$$

Where: a is the shape factor (2.018) (Richter et al., 2014), (LL= Leaf Length, LW= Leaf Width and CA= Cultivated Area

1 **2.8 Estimation of Crop Coefficients from the FAO56 Approach**

2 The crop coefficients (K_c) values represent the integrated effects of changes in leaf area,
3 plant height, crop characteristics, irrigation method, rate of crop development, crop planting
4 date, degree of canopy cover, canopy resistance, soil and climate conditions, and management
5 practices (Allen et al., 1998). Each crop will have a set of specific crop coefficient and will
6 predict different water use, for different crops for different growth stages (Irkman, 2008). Crop
7 coefficients were estimated for each growth stage from Equation 16.

$$8 \quad K_c = \frac{ET_c}{ET_p} \quad (16)$$

9 Where: K_c = Crop Coefficient, ET_c = Actual Crop Evapotranspiration (mm/week) ET_p =
10 Reference Evapotranspiration (mm/week)

11 **2.9 Data analysis**

12 Analysis of variance (ANOVA) was used to statistically determine the differences in
13 measured means and variances of all data. The doebio research statistical package in R Studio
14 4.1.3 (released 10th March, 2022) was used for the statistical analysis. Least significant
15 difference (LSD) was used to separate all means at the 95% level of confidence.

16 **3.0 Results and Discussions**

17 **3.1 Soil Physicochemical Properties**

18 Table 1 presents the mean and standard deviations of the soils' bulk densities, hydraulic
19 conductivities, nitrogen, phosphorus, potassium, calcium, magnesium contents and cation
20 exchange capacities. The bulk densities of the soil samples exhibited a range from 1.40 g/cm³
21 to 1.55 g/cm³. Notably, sandy soil displayed the highest bulk density, whereas loam and clay
22 soils exhibited similar bulk density values. In terms of saturated hydraulic conductivities, there
23 were no significant variations observed among the three soil textures, with values ranging from
24 a maximum of 0.11 cm/sec to a minimum of 0.09 cm/sec, as detailed in Table 1. The results
25 pertaining to saturated hydraulic conductivity indicated a close proximity between the values
26 for sandy clay loam and sandy loam soils, particularly. Specifically, the mean bulk density of
27 sand was found to be 1.55 g/cm³, which was significantly higher when compared to the values
28 obtained for both sandy clay loam (1.40 g/cm³) and sandy loam (1.45 g/cm³) at a significance
29 level of $p \leq 0.05$. The percentages of sand, silt and clay were used to separate the soil into their
30 respective textural classes as shown on Table 1, and applied throughout the study.

31 The chemical properties analysed gave clue to the nutrient status of the soils. The sandy
32 clay loam soil has the highest nitrogen and phosphorus contents as compared to the other soil
33 textures. The sandy soil however has the potassium contents while the calcium and magnesium

1 contents of the sandy loam soils are higher than the other two textures. The highest CEC was
 2 as well found in the sandy clay loam. Statistically, the chemical properties inherent in the three
 3 soil texture are significantly different from one texture too the other.

4 **Table 1: Mean and Standard Deviation of Soils' Physicochemical Properties for**
 5 **Different Soil Textural Class**

Physical Parameters	Sandy	Sandy Clay Loam	Sandy Loam	LSD (p < 0.05)
Bulk Density (g/cm ³)	1.55±0.11 ^a	1.40±0.12 ^b	1.45±0.09 ^b	0.12
Hydraulic Conductivity (cm/s)	0.11±0.02 ^a	0.09±0.03 ^a	0.10±0.03 ^a	0.02
Sand (%)	91.11±3.56 ^a	74.72±2.99 ^c	75.83±3.12 ^b	2.08E-14
Clay (%)	7.22±2.99 ^c	23.33±3.12 ^a	18.06±3.07 ^b	3.55E-14
Silt (%)	1.67±2.87 ^c	1.95±1.94 ^b	6.11±1.71 ^a	6.74E-14
Nitrogen (mg/kg)	0.048±0.001 ^b	0.058±0.001 ^a	0.044±0.001 ^c	0.002
Phosphorus (mg/kg)	6.13±0.01 ^c	8.23±0.01 ^a	6.93±0.01 ^b	0.020
Potassium (mg/kg)	21.86±0.01 ^a	17.28±0.01 ^b	15.20±0.02 ^c	0.116
Calcium (mg/kg)	1.12±0.01 ^b	0.50±0.01 ^c	1.26±0.01 ^a	0.020
Magnesium (mg/kg)	0.08±0.01 ^c	0.14±0.01 ^b	0.20±0.01 ^a	0.020
CEC (cmol/kg)	7.82±0.00 ^b	13.69±0.00 ^a	7.37±0.00 ^c	0.020

6 *Values with the same superscripts are not significantly different. LSD is Least Significant Difference

7 **3.2 Agronomic Growth Parameters**

8 **3.2.1 Plant Height**

9 Table 2 provides an overview of the mean plant heights and their standard deviations
 10 recorded throughout the growth stages of three different legume varieties. During the
 11 emergence stage after planting, the plant heights ranged from 18.83 cm to 36.17 cm across the
 12 various legume varieties grown on the three soil textures. When flowering occurred, plant
 13 heights ranged from 31.43 cm (V3 on sandy soil) to 59.40 cm (V2 on sandy soil). Notably,
 14 V2's height differed significantly from V3's height on sandy soil during this stage.
 15

16 Moving to the pod setting growth stage, a significant difference was observed between
 17 V1 on sandy soil and V2 on sandy clay loam. This trend persisted into the maturity stage.
 18 Interestingly, the performance of V2 on sandy clay loam soil exhibited the lowest plant height.
 19 The maximum plant height recorded at maturity was 115.50 cm, observed in V1 on sandy soils.
 20 Between the flowering and emergence stages, plant heights for varieties 1, 2, and 3 increased
 21 by average percentages of 40.29%, 31.24%, and 31.74%, respectively. However, between the
 22 pod setting and maturity stages, the growth rate slowed down, resulting in average increases of
 23 17.73 cm, 16.26 cm, and 0.60 cm for Varieties 1, 2, and 3, respectively.

24 Notably, legume varieties planted in sandy soil exhibited the highest range of plant
 25 heights (27.50 to 115.50 cm), whereas those planted on sandy clay loam had the lowest plant
 26 heights range (18.83 to 97.17 cm). This discrepancy may be attributed to root restriction due
 27 to the compacted nature of clay soil, which hinders the plants' ability to adequately absorb
 28 water from the soil (Eldeiry, 2005).

Table 2: Mean and Standard Deviation of Plant Height of Legume Varieties at Different Growth Stages (cm)

Growth Stages	Variety/Treatment	Sandy	Sandy Clay Loam	Sandy Loam
Emergence	V1	27.50± 1.22 ^{abc}	30.00±3.77 ^{ab}	26.17±5.10 ^{bc}
	V2	36.17±4.13 ^a	30.83±1.65 ^{ab}	33.17±4.90 ^{ab}
	V3	26.50±1.63 ^{bc}	32.67±8.99 ^{ab}	18.83±4.50 ^c
	LSD (p < 0.05)	9.44		
Flowering	V1	56.50± 9.39 ^a	42.00±5.72 ^{ab}	44.33±6.51 ^{ab}
	V2	59.40±22.34 ^a	42.93±2.67 ^{ab}	45.10±5.78 ^{ab}
	V3	31.43±14.70 ^b	44.17±11.26 ^{ab}	40.50±11.50 ^{ab}
	LSD (p < 0.05)	24.49		
Pod setting	V1	97.00±2.45 ^a	73.33±15.56 ^{ab}	72.17±9.72 ^{ab}
	V2	90.83±43.41 ^{ab}	55.17±8.37 ^b	56.33±5.24 ^{ab}
	V3	75.83±24.87 ^{ab}	92.00±15.58 ^{ab}	95.50±14.84 ^{ab}
	LSD (p < 0.05)	40.68		
Maturity	V1	115.50±1.22 ^a	100.50±3.27 ^{ab}	80.33±13.68 ^{ab}
	V2	99.00±42.43 ^{ab}	55.67±8.18 ^b	93.33±45.73 ^{ab}
	V3	75.90±11.64 ^{ab}	92.00±11.66 ^{ab}	97.17±15.97 ^{ab}
	LSD (p < 0.05)	46.74		

*Values with the same superscripts are not significantly different. V1 is variety 1 of cowpea (sampea 14), V2 is variety 2 of cowpea (sampea 17) and V3 is a variety of soybean (TGX 1465-1D), LSD is Least Significant Difference

3.2.2 Number of Leaves

The number of leaves generated by the various legume varieties at specific growth stages is a crucial parameter for assessing crop transpiration and water usage. The resulting leaf numbers for the planted legume varieties are presented in Table 3. During the emergence stage, the number of leaves observed across the three soil textures ranged from 4 to 9. Notably, V1 cultivated on sandy soil exhibited the highest leaf count (9), while the same variety on sandy clay loam soil had the lowest leaf count (4). Among the legume varieties, V2 produced the highest number of leaves across all three soil textures.

As the flowering stage approached, V1 on sandy loam soil displayed the highest leaf numbers (16), whereas V1 on sandy clay loam had the lowest leaf count (6). When the legumes entered the pod-setting stage, V3 on sandy loam soil bore the highest number of true leaves (30), whereas V1 and V2 on sandy clay loam exhibited the lowest leaf numbers (10). However, at maturity, V1 on sandy soil had the highest number of true leaves (34), while V2 on sandy clay loam had the lowest leaf count (14).

Between the flowering and emergence stages, there was a substantial increase in the number of leaves for each variety, with V1, V2, and V3 showing average percentage increases of 47.36%, 40.76%, and 33.33%, respectively. However, between the pod-setting and maturity stages, the rate of increase in leaf numbers slowed down to 27.68%, 25.00%, and 8.39%, respectively. This phenomenon can be attributed to physiological leaf deterioration, aging, and leaf shedding (Nuwamanya, et al., 2019).

1 Statistically, significant differences were observed at emergence in the number of leaves
 2 produced by V1 on sandy soil and V1 and V3 on sandy clay loam, as well as among the three
 3 varieties on sandy loam soils (Table 3). At the flowering stage, the leaf numbers of V1 on sandy
 4 soil differed significantly from those of V3 on sandy soil, V1 and V2 on sandy clay loam, and
 5 V3 on sandy loam. During the pod-setting stage, significant differences were noted in the leaf
 6 numbers produced by V3 planted on sandy loam soil compared to the numbers for V2 on sandy
 7 loam and V1 and V2 on sandy clay loam. Additionally, at maturity, significant differences in
 8 leaf numbers were observed between V2 on sandy clay loam and V1 and V3 on sandy soil, as
 9 well as V3 on sandy clay loam.

10 **Table 3: Mean and Standard Deviation of Number of Leaves of Legume Varieties at**
 11 **Different Growth Stages**

Growth Stages	Variety/Treatment	Sandy	Sandy Clay Loam	Sandy Loam
Emergence	V1	9.00±1.00 ^a	4.00±2.00 ^c	5.00±2.00 ^{bc}
	V2	8.00±0.00 ^{ab}	7.00±1.00 ^{abc}	6.00±1.00 ^{bc}
	V3	7.00±0.00 ^{abc}	6.00±1.00 ^{bc}	5.00±2.00 ^{bc}
	LSD (p < 0.05)	3.04		
Flowering	V1	15.00±3.00 ^a	6.00±0.00 ^c	16.00±5.00 ^a
	V2	13.00±3.00 ^{ab}	10.00±2.00 ^{bc}	13.00±1.00 ^{ab}
	V3	7.00±2.00 ^c	12.00±1.00 ^{ab}	10.00±2.00 ^{bc}
	LSD (p < 0.05)	4.57		
Pod setting	V1	26.00±4.00 ^{ab}	10.00±6.00 ^c	19.00±9.00 ^{abc}
	V2	22.00±12.00 ^{abc}	10.00±2.00 ^c	15.00±5.00 ^{bc}
	V3	26.00±7.00 ^{ab}	28.00±9.00 ^{ab}	30.00±1.00 ^a
	LSD (p < 0.05)	14.98		
Maturity	V1	34.00±5.00 ^a	20.00±4.00 ^{ab}	21.00±7.00 ^{ab}
	V2	22.00±8.00 ^{ab}	14.00±6.00 ^b	28.00±1.00 ^{ab}
	V3	31.00±4.00 ^a	32.00±13.00 ^a	29.00±11.00 ^{ab}
	LSD (p < 0.05)	28.17		

12 *Values with the same superscripts are not significantly different. V1 is variety 1 of cowpea (sampea 14), V2 is
 13 variety 2 of cowpea (sampea 17) and V3 is a variety of soybean (TGX 1465-1D), LSD is Least Significant
 14 Difference

16 3.2.3 Leaf Area Index

17 Leaf area is a critical growth parameter that serves as an indicator of both leaf canopies
 18 and the extent of soil coverage by leaves. It plays a significant role in influencing water
 19 evaporation from the soil surface and transpiration from leaf surfaces. Table 4 presents the
 20 estimated mean values of the leaf area indices (LAI) for the various legume varieties grown on
 21 three distinct soil textures.

22 During the emergence stage, the LAI of the legume varieties ranged from 0.015 to
 23 0.021, and these ranges did not exhibit statistically significant differences. Notably, V1
 24 cultivated on sandy clay loam soil displayed the highest LAI (0.021), while the same V1
 25 planted in sandy clay loam had the lowest LAI. As the flowering stage approached, the LAI
 26 range expanded from 0.015 (V3 on sandy clay loam) to 0.027 (V1 on sandy soil), with these

1 two values being significantly different. This suggests that, despite having a high number of
 2 leaves at this growth stage, these leaves had relatively smaller surface areas, resulting in smaller
 3 canopies and LAI. During the pod-setting phase, V 1 planted on sandy soil exhibited the highest
 4 LAI (0.108), while V1 planted on sandy clay loam had the lowest LAI (0.027). This significant
 5 difference in LAI was observed between V1 on sandy soil and V3 on sandy soil, as well as
 6 between V1 and V3 on sandy clay loam and V 3 on sandy loam soil.

7 At maturity, the highest LAI was recorded for V1 planted on sandy soil, whereas V3
 8 planted on sandy clay loam and sandy loam exhibited the lowest LAI values (Table 4).
 9 Consequently, the LAI of V1 on sandy soil differed significantly from that of all other legume
 10 varieties on different soil textures, except for V2 on sandy soil and sandy loam soil. In
 11 summary, it appears that legume varieties V1 and V2 had better LAI on sandy soil compared
 12 to the other soil textures.

13 **Table 4: Mean and Standard Deviation of Leaf area index of Legume Varieties at**
 14 **Different Growth Stages**

Growth stages	Variety/Treatment	Sandy	Sandy Clay Loam	Sandy Loam
Emergence	V1	0.021±0.01 ^a	0.015±0.005 ^a	0.018±0.01 ^a
	V2	0.025±0.01 ^a	0.017±0.002 ^a	0.016±0.004 ^a
	V3	0.023±0.01 ^a	0.018±0.004 ^a	0.018±0.001 ^a
	LSD (p < 0.05)	0.012		
Flowering	V1	0.027±0.002 ^a	0.017±0.000 ^{ab}	0.017±0.00 ^{ab}
	V2	0.015±0.010 ^b	0.021±0.004 ^{ab}	0.021±0.003 ^{ab}
	V3	0.018±0.010 ^{ab}	0.015±0.002 ^b	0.019±0.003 ^{ab}
	LSD (p < 0.05)	0.010		
Pod setting	V1	0.108±0.007 ^a	0.027±0.009 ^c	0.063±0.026 ^{abc}
	V2	0.096±0.085 ^{ab}	0.045±0.011 ^{abc}	0.063±0.036 ^{abc}
	V3	0.036±0.008 ^{bc}	0.031±0.0160 ^c	0.041±0.002 ^{bc}
	LSD (p < 0.05)	0.064		
Maturity	V1	0.131±0.016 ^a	0.040±0.002 ^{cd}	0.063±0.022 ^{bcd}
	V2	0.101±0.082 ^{ab}	0.042±0.020 ^{bcd}	0.100±0.005 ^{abc}
	V3	0.040±0.010 ^{bcd}	0.031±0.016 ^d	0.032±0.007 ^d
	LSD (p < 0.05)	0.061		

15 *Values with the same superscripts are not significantly different. V1 is variety 1 of cowpea (sampea 14), V2 is
 16 variety 2 of cowpea (sampea 17) and V3 is a variety of soybean (TGX 1465-1D), LSD is Least Significant
 17 Difference

18 3.3 Crop Water Use and Crop Coefficients

19 Tables 5 and Figure 3 present data on water consumption and resulting crop
 20 coefficients, respectively, for each legume variety. Detailed data on actual crop water use,
 21 reference evapotranspiration and crop coefficients are presented on Supplementary Table 1.
 22 During the initial season, it was observed that V3 planted on sandy loam soil had the highest
 23 water consumption, followed by V2 on sandy and V2 on sandy loam soils. In contrast, V2 on
 24 sandy clay loam exhibited lower water consumption compared to the other two legume
 25

1 varieties on the same soil. The water use for V1, V2, and V3 ranged from 47.7 mm/m to 583.55
 2 mm/m across the three soil textures.

3
 4 **Table 5: Mean and Standard Deviation of Actual Evapotranspiration by Legume**
 5 **Varieties at Different Growth Stages (mm)**

Growth stages	Variety/Treatment	Sandy	Sandy Clay Loam	Sandy Loam
Initial Season	V1	59.08±2.36 ^{abc}	67.70±1.98 ^{abc}	48.93±1.9.2 ^{bc}
	V2	90.03±2.56 ^{ab}	47.70±2.56 ^c	90.11±2.56 ^{ab}
	V3	57.17±0.51 ^{abc}	44.68±1.13 ^c	93.90±3.43 ^a
	LSD (p < 0.05)	41.44		
Mid-Season	V1	454.74±10.78 ^{abc}	583.55±10.90 ^a	346.42±5.63 ^c
	V2	352.04±2.80 ^{bc}	422.41±3.23 ^{abc}	442.23±8.28 ^{abc}
	V3	391.75±3.80 ^{abc}	445.11±14.07 ^{abc}	546.59.2±18.62 ^{ab}
	LSD (p < 0.05)	195.10		
Late Season	V1	90.75±2.21 ^a	59.55±1.23 ^a	90.72±0.13 ^a
	V2	70.08±3.88 ^a	46.00±2.63 ^a	93.83±4.97 ^a
	V3	84.80±2.01 ^a	68.27±2.97 ^a	41.47±1.34 ^a
	LSD (p < 0.05)	55.43		

6 *Values with the same superscripts are not significantly different. V1 is variety 1 of cowpea (sampea 14), V2 is
 7 variety 2 of cowpea (sampea 17) and V3 is a variety of soybean (TGX 1465-1D), LSD is Least Significant
 8 Difference

9
 10 At the initial season, legumes grown on sandy and sandy loam soils consumed more
 11 water than those on sandy clay loam soil. The larger pore spaces in sandy and sandy loam soils,
 12 compared to sandy clay loam, allowed for sufficient water release to the atmosphere. The
 13 permanent wilting point (PWP) of sandy soils is thus below the threshold PWP of other soil
 14 textural classes. Consequently, more water applications may be needed to meet the crop water
 15 requirement on sandy soils during the initial season when the soil surfaces were still bare and
 16 uncovered (Myneni et al., 2002; Somers et al., 2010; Vallejos, 2008; Dong et al., 2019).
 17 Statistically, significant differences were observed in water consumptions at the crops' initial
 18 stages, particularly between V3 on sandy loam soil and V2 on sandy clay loam soil, as well as
 19 between V3 on sandy loam soil and V3 on sandy clay loam soil (Table 5).

20 During the mid-season, a noticeable increase in water consumption was observed for
 21 all three legume varieties compared to the initial stages. The variety V1 planted on sandy clay
 22 loam exhibited the highest water consumption (583.55 mm/m), while the lowest water
 23 consumption was recorded for V1 on sandy loam soil (346.30 mm/m). At this stage, significant
 24 differences in water consumption were observed between V1 on sandy clay loam soil and the
 25 following plots: V3 on sandy soil and V1 on sandy loam soils.

26 During the late season, typically characterized by pod maturation, water consumption
 27 rates for all three varieties decreased compared to the mid-growth stage. In this late season, the
 28 crops experienced reduced transpiration, resulting in significantly lower water usage compared
 29 to the mid-season. This decrease in transpiration rates can be attributed to physiological leaf

1 deterioration, aging, and leaf shedding (Nuwamanya et al., 2019). Notably, the water
2 consumptions of legumes at this stage were not significantly different from one soil texture and
3 legume variety to the other.

4 The FAO-56 crop coefficient approach, commonly utilised for estimating crop water
5 requirements under well-watered conditions, stands as the standard method widely adopted for
6 irrigation management, particularly for irrigation scheduling (Allen et al., 1998). According to
7 this approach, K_c tends to increase as the crop develops, reaching maximum values when the
8 crops are fully matured (Abedinpour, 2016; López-Urrea et al., 2021). The findings of this
9 study also revealed a gradual increase in K_c values from the initial growth stage to the
10 developmental and mid-season stages, followed by a decrease in the late-season's K_c . This
11 pattern of increasing crop coefficients reflects the impact of crop growth, development, and
12 physiological processes on water consumption and evapotranspiration (Shenkut et al., 2013).

13 According to Allen et al. (1998), K_c values of 0.4, 1.15, and 0.55 were determined for
14 legumes during the initial, mid-season, and late-season, respectively. However, in this study,
15 the averages of K_c obtained are 0.69, 1.46, and 0.78 for the initial, mid-season, and late-seasons,
16 respectively. Specifically, legumes grown on sandy soils had an average K_c of 0.71, 1.32, and
17 0.90 for the initial, mid-season, and late-season stages, respectively. On sandy clay loam soils,
18 the average K_c s are 0.54, 1.59, and 0.65, and on sandy loam soils, they are 0.82, 1.47, and 0.79,
19 respectively (Figure 3). The slight variations in K_c observed can be attributed to differences in
20 climate and soil characteristics, as noted by Kullberg et al. (2017) and Jamshidi et al. (2020).

21 It's worth noting that the K_c obtained for V1, V2, and V3 closely align with those
22 estimated by Alla-Jabow et al. (2015) for faba bean, chickpea, and common bean in a semi-
23 desert climatic region with very hot conditions in Ed-Damer, Sudan. Alla-Jabow et al. (2015)
24 estimated values of 0.33, 0.26, and 0.52 for faba beans, chickpea, and common bean,
25 respectively, during the initial stages. During the mid-season, the reported K_c values were 1.08,
26 1.22, and 1.07 for faba beans, chickpea, and common beans, respectively. The K_c was observed
27 to have increased from the initial stage. However, during the late season, the K_c decreased to
28 0.60, 0.52, and 0.52, in the same order. During the initial stage, the averages of K_c for V1, V2,
29 and V3 are 0.55, 0.71, and 0.81, respectively. At the mid-growth stage, the averages of K_c are
30 1.48, 1.30, and 1.60 in the same order and during the late season, the averages of K_c are 0.79,
31 0.68, and 0.86, respectively (Figure 3). These coefficients depict variations in crop water
32 requirements and transpiration rates at different growth stages, on different soil textures, and
33 among different legume varieties.


1 In a similar study conducted by El-Noemani et al. (2015) on irrigated varieties of
2 *Phaseolus vulgaris L.* in the Nile Delta clay loam old alluvial soil, the initial K_c ranged from
3 0.63 to 0.64 for the bronco variety. The K_c ranges of 0.82 to 0.87, 0.99 to 1.09, and 0.80 to 0.95
4 were obtained for the flowering, pod-setting and maturity stages, respectively, at varying levels
5 of water applications (40%, 60%, and 80%). Furthermore, K_c values ranges of 0.59 to 0.61,
6 0.78 to 0.98, 1.07 to 1.19 and 0.73 to 0.88 were obtained for the contender variety at flowering,
7 pod-setting and maturity stages at varying levels of water application.

8 **4.0 Conclusions**

9 This study aimed to investigate water usage and crop coefficients of legumes cultivated
10 on three distinct soil textures in southern Nigeria, with the goal of improving crop and water
11 management practices. The findings indicated that legume water consumption was highest
12 during the mid-season, influenced by various factors such as climatic conditions, months, and
13 seasons of the year. Legume varieties planted on sandy clay loam soils exhibited the highest
14 average actual evapotranspiration, indicating greater water usage or soil water depletion on this
15 soil type. Notably, crop coefficients estimated for legumes grown on sandy loam were higher
16 than those obtained for sandy and sandy clay loam soils, particularly during the mid and late
17 seasons. The soybean variety V3 consistently displayed higher average K_c across the soil types
18 than the two cowpea varieties, V1 and V2, all growth stages.

19 Although, only one-time trial of this experiment was conducted however, the findings
20 provide major information on the K_c of legumes. It is recommended that the coefficients
21 obtained from this study be re-affirmed within and outside the study area, especially by
22 repeating the study during several planting seasons. Furthermore, the use of machine learning
23 models for the estimation of ET_p and ET_c , and thus K_c , is also recommended. Less human-
24 dependent methods should be employed to monitor changes in soil moisture content over time,
25 in such studies, in order to ensure the collection of error-free data.

26 The implications of this study are significant, especially for irrigation engineers,
27 agriculturists, soil scientists, and environmentalists. The findings can inform improved
28 irrigation scheduling, water management practices, and enhanced crop and water productivity
29 in agriculture, particularly in regions with a climate similar to southern Nigeria.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

REFERENCES

Abedinpour, M. (2016). Evaluation of growth-stage-specific crop coefficients of maize using weighing lysimeter. *Soil and Water Research*, 10(2): 99–104. <https://doi.org/10.17221/63/2014-swr>

Adesogan S. O. and Sasanya B. F. (2023). Efficiency of indirect and estimated evapotranspiration methods in South Western Nigeria. *International Journal of Hydrology Science and Technology*, 15(1): 64-77. <https://doi.org/10.1504/IJHST.2021.10041388>

Alla Jabow, M.K., Ibrahim, O.H. and Adam, H.S. (2015) Evapotranspiration and Crop Coefficients of Grain Legumes in Semi Desert Climatic Conditions, *African Journal of Crop Science*, 23(3):239 - 247

Allen, R. G., Pereira, L. S., Raes, D., Smith, M., & Ab, W. (1998). Crop Evapotranspiration - Guidelines for Computing Crop Water Requirements - *FAO Irrigation and Drainage Paper 56*.

Bache, B. W. (1976). The measurement of cation exchange capacity of soils. *Journal of the Science of Food and Agriculture*, 27(3): 273-280.

- 1 Bariviera, G. Dallacort, R., de Freitas, P. S. L. Barbieri, J. D. and Daniel, D. F. (2020). Crop
2 coefficient for the early-cycle soybean cultivar SoyTech 815 RR *Revista Brasileira de*
3 *Engenharia Agrícola e Ambiental* 24(2):75-81 [http://dx.doi.org/10.1590/1807-](http://dx.doi.org/10.1590/1807-1929/agriambi.v24n2p75-81)
4 [1929/agriambi.v24n2p75-81](http://dx.doi.org/10.1590/1807-1929/agriambi.v24n2p75-81)
5
- 6 Blake, G. R. and Hartge, K.H. (1986). Bulk density. In: Klute, A., Ed., *Methods of Soil*
7 *Analysis, Part 1—Physical and Mineralogical Methods*, 2nd Ed., *Agronomy Monograph*
8 *9, American Society of Agronomy—Soil Science Society of America, Madison*, 363-382.
9
- 10 Córdova, M., Carrillo-Rojas, G. Crespo, P., Wilcox, and B. Céleri, R. (2015) Evaluation of the
11 Penman-Monteith (FAO 56 PM) Method for Calculating Reference Evapotranspiration
12 Using Limited Data. *Mountain Research and Development*, 35(3):230-239.
13 <https://doi.org/10.1659/MRD-JOURNAL-D-14-0024.1>
14
- 15 Chandra, R. and Kumari, S. (2021). Estimation of crop water requirement for rice- wheat and
16 rice- maize cropping system using CROPWAT model for Pusa, Samastipur district, Bihar.
17 *Journal of AgriSearch*, 8(2). <https://doi.org/10.21921/jas.v8i2.7299>
18
- 19 Dehghanisani, H. and Kosari, H. (2011). Evapotranspiration Partitioning in Surface and
20 Subsurface Drip Irrigation Systems. *InTech*. <https://doi.org/10.5772/17122>
21
- 22 Dingre, S. K., Gorantiwar, S. D., & Kadam, S. A. (2021). *Correlating the field water balance*
23 *derived crop coefficient (Kc) and canopy reflectance-based NDVI for irrigated sugarcane.*
24 *Precision Agriculture*. doi:10.1007/s11119-020-09774-8
25
- 26 Djaman, K., O'Neill, M., Owen, C. K., Smeal, D., Koudahe, K., West, M., Allen, S., Lombard,
27 K., & Irmak, S. (2018). Crop evapotranspiration, irrigation water requirement and water
28 productivity of maize from meteorological data under semiarid climate. *Water*
29 *(Switzerland)*, 10(4). <https://doi.org/10.3390/w10040405>
30
- 31 Djaman, K., Rudnick, D. R. Moukoubi, Y. D., Sow, A. Irmak, S. (2019). Actual
32 evapotranspiration and crop coefficients of irrigated lowland rice (*Oryza sativa L.*) under
33 semiarid climate *Italian Journal of Agronomy* 14(1):19-25.
34 <https://doi.org/10.4081/ija.2019.1059>
35
- 36 Dong, T., Liu, J., Shang, J., Qian, B., Ma, B., Kovacs, J. M., ... Shi, Y. (2019). Assessment of
37 red-edge vegetation indices for crop leaf area index estimation. *Remote Sensing of*
38 *Environment*, 222: 133–143. doi:10.1016/j.rse.2018.12.032
39
- 40 Drechsler, K. Fulton, A. Kisekka, I. (2022). Crop coefficients and water use of young almond
41 orchards. *Irrigation Science* 40:379–395 <https://doi.org/10.1007/s00271-022-00786-y>
42
- 43 Eldeiry, A. A., Garcia, L. A., El-Zaher, A. S. A. and El-Sherbini Kiwan. M., (2005). Furrow
44 Irrigation System Design For Clay Soils In Arid Regions. *Applied Engineering in*
45 *Agriculture*, 21(3): 411–420. doi:10.13031/2013.18460
46
- 47 El-Noemani, A. A., Aboamera M. A. H and Dewedar, O. M. (2015). Determination of crop
48 coefficient for bean (*Phaseolus vulgaris L.*) plants under drip irrigation system,
49 *International Journal of ChemTech Research* 8(12): 203-214
50

- 1 Fan, Y., He, L., Kang, S., Wang, S., and Fang, Y. (2021). A novel approach to dynamically
2 optimize the spatio-temporal distribution of crop water consumption. *Journal of Cleaner*
3 *Production*, 310. doi:10.1016/j.jclepro.2021.127439
4
- 5 Gavlack, R., D. Horneck, and R. Miller. (2005). Plant, soil and water reference methods for
6 Western Region. *Western Region Extension Publication (WREP) 125*, WERA-103
7 Technical committee.
8
- 9 Poyen, F. B., Ghosh, A. K., and Palash, K. (2016). Review on different evapotranspiration
10 empirical equations, *International Journal of Advanced Engineering, Management and*
11 *Science*, 2 (3): 17-24
12
- 13 Gong, X., Qiu, R., Sun, J., Ge, J., Li, Y., & Wang, S. (2020). Evapotranspiration and crop
14 coefficient of tomato grown in a solar greenhouse under full and deficit irrigation.
15 *Agricultural Water Management*, 235. doi:10.1016/j.agwat.2020.10615
16
- 17 Hassan, D. F., Abdalkadhum, A. J., Mohammed, R. and Shaban, A. (2022). Integration Remote
18 Sensing and Meteorological Data to Monitoring Plant Phenology and Estimation Crop
19 Coefficient and Evapotranspiration. *Journal of Ecological Engineering* 23(4):325–335
20 [10.12911/22998993/146267](https://doi.org/10.12911/22998993/146267)
21
- 22 Houba, V. J. G., Temminghoff, E. J. M., Gaikhorst, G. A., & van Vark, W. (2000). Soil analysis
23 procedures using 0.01 M calcium chloride as extraction reagent. *Communications in Soil*
24 *Science and Plant Analysis*, 31(9–10): 1299–1396.
25 <https://doi.org/10.1080/00103620009370514>
26
- 27 Jabro, J. Stevens, W. and Iversen, W. (2020) Comparing Two Methods for Measuring Soil Bulk
28 Density and Moisture Content. *Open Journal of Soil Science*, 10: 233-24
29
- 30 Jamshidi, S., Zand-Parsab, S., Kamgar-Haghighib, A. A., Shahsavarc, A. R., Niyogia, D.
31 (2020). Evapotranspiration, crop coefficients, and physiological responses of citrus trees
32 in semi-arid climatic conditions. *Agricultural Water Management* 227 (2020).
33 <https://doi.org/10.1016/j.agwat.2019.105838>
34
- 35 Khan, M. J., Malik, A., Rahman, M., Afzaal, M. and Mulk, S. (2021). Assessment of Crop
36 Water Requirement for Various Crops in Peshawar, Pakistan Using CROPWAT. *Irrigation*
37 *Drainage System Engineering*, 10(9.):1-10
38
- 39 Kalidass, C., & Mahapatra, A. K. (2014). Evaluation of the proximate and phytochemical
40 compositions of an underexploited legume *Mucua pruriens* var. *utilis* (wall ex
41 wight)l.h.bailey. *International Food Research Journal*, 21(1): 303-308
42
- 43 Kenjabaev, S., Frede, H. G., Begmatov, I., Isaev, S., & Matyakubov, B. (2020). Determination
44 of actual crop evapotranspiration (etc) and dual crop coefficients (kc) for cotton, wheat
45 and maize in fergana valley: integration of the fao-56 approach and budget. *Journal of*
46 *Critical Reviews*, 7(05):340–349. <https://doi.org/10.31838/jcr.07.05.67>
47
- 48 Kullberg, E. G., DeJonge, K. C., & Chávez, J. L. (2017). Evaluation of thermal remote sensing
49 indices to estimate crop evapotranspiration coefficients. *Agricultural Water Management*,
50 179:64–73. <https://doi.org/10.1016/j.agwat.2016.07.007>

- 1
2 López-Urrea, R., Sánchez, J. M., de la Cruz, F., González-Piqueras, J., & Chávez, J. L. (2020).
3 Evapotranspiration and crop coefficients from lysimeter measurements for sprinkler-
4 irrigated canola. *Agricultural Water Management*, 239.
5 <https://doi.org/10.1016/j.agwat.2020.106260>
6
- 7 Myneni, R., Hoffman, S., Knyazikhin, Y., Privette, J., Glassy, J., Tian, Y., ... Running, S.
8 (2002). Global products of vegetation leaf area and fraction absorbed PAR from year one
9 of MODIS data. *Remote Sensing of Environment*, 83(1-2): 214–231.
10 [https://doi.org/10.1016/s0034-4257\(02\)00074-3](https://doi.org/10.1016/s0034-4257(02)00074-3)
11
- 12 National Agricultural Extension and Research Liaison Services (NAERLS) (2010).
13 *Agricultural Performance Survey of 2010 Wet Season in Nigeria*. Zaria, Nigeria, 182.
14
- 15 National Programme for Agriculture and Food Security (NPAFS) (2010). Report of the 2009
16 *Agricultural Production Survey*. Abuja, Nigeria, 86.
17
- 18 Nouri, H., Glenn, E. P., Beecham, S., Boroujeni, S. C., Sutton, P., Alaghmand, S., Noori, B., &
19 Nagler, P. (2016). Comparing three approaches of evapotranspiration estimation in mixed
20 urban vegetation: Field-based, remote sensing-based and observational-based methods.
21 *Remote Sensing*, 8(6). <https://doi.org/10.3390/rs8060492>
22
- 23 Nuwamanya, E., Acheng, S., Vuzi, P., Muyinza, H., Matovu, M., Atwijukire, E., ... Adebayo,
24 B. A. (2019). Effectiveness of pruning and waxing in reducing postharvest physiological
25 deterioration in Uganda local cassava varieties. *African Crop Science Journal*, 27(2): 237-
26 238.
27
- 28 O'Dell, J. W. (1993). Determination of total Kjeldahl nitrogen by semi-automated
29 colorimetry. *Environ. Monit. Syst. Lab*, 1, 15.
30
- 31 Pereira, L. S., Paredes, P., Hunsaker, D. J., López-Urrea, R., & Mohammadi Shad, Z. (2021).
32 Standard single and basal crop coefficients for field crops. Updates and advances to the
33 FAO56 crop water requirements method. *Agricultural Water Management* 243.
34 <https://doi.org/10.1016/j.agwat.2020.106466>
35
- 36 Rallo, G., Paço, T. A., Paredes, P., Puig-Sirera, À., Massai, R., Provenzano, G., & Pereira, L.
37 S. (2021). Updated single and dual crop coefficients for tree and vine fruit crops.
38 *Agricultural Water Management*, 250. doi:10.1016/j.agwat.2020.106645
39
- 40 Reynolds, W. D., Elick, D.E., Youngs, E.G., Amoozegar, A. and Bootink, N.W. 2002. Saturated
41 and Field-saturated Water Flow Parameters, In: Dane, J.H., Topp, G.C. (eds.) , Part 4.
42 SSSA, Madison, WI, USA. 797–878.
43
- 44 Richter, G. L., Zanon, A. J. Streck, N. A. Guedes, J. V. C., Kräulich, B., Rocha, T. S. M., Wink,
45 J. E. M and Cera, J. C. (2014). Estimating leaf area of modern soybean cultivars by a
46 non-destructive method. *Bragantia* 73: 416–425. <https://doi.org/10.1590/1678-4499.0179>
47
48
- 49 Sepaskhah, A. R., & Razzaghi, F. (2009). Evaluation of the adjusted Thornthwaite and
50 Hargreaves-Samani methods for estimation of daily evapotranspiration in a semi-arid

1 region of Iran. *Archives of Agronomy and Soil Science*, 55(1), 51–66.
2 <https://doi.org/10.1080/03650340802383148>

3
4 Silva, W. A. D., Santana, J. S., Silva, C. M. D., & Nunes, A. A. (2017). Crop coefficient
5 regionalization for irrigated agriculture planning in Maranhão state-brazil. *Engenharia*
6 *Agrícola*, 37:953-960.[https://doi.org/10.1590/1809-4430-Eng.Agric.v37n5p953-](https://doi.org/10.1590/1809-4430-Eng.Agric.v37n5p953-960/2017)
7 960/2017

8
9 Shenkut, A., Tesfaye, K., & Abegaz, F. (2013). Determination of Water Requirement and Crop
10 Coefficient for Sorghum (*Sorghum bicolor L.*) at Melkassa, Ethiopia. *Science, Technology*
11 *and Arts Research Journal*, 2(3). <https://doi.org/10.4314/star.v2i3.98717>

12
13 Somers, B., Gysels, V., Verstraeten, W. W., Delalieux, S., & Coppin, P. (2010). Modelling
14 moisture-induced soil reflectance changes in cultivated sandy soils: a case study in citrus
15 orchards. *European Journal of Soil Science*, 61(6): 1091–1105.
16 <https://doi.org/10.1111/j.1365-2389.2010.01305.xn>

17
18 Sultana, N., Salam, A., Ahammad, T., & Zahir, S. M. A. (2022). Remote Sensing Based
19 Assessment of Crop Water Requirement of Boro Rice Using SEBAL Model. *DEW-DROP*
20 8:58–62.

21
22 Vallejos, J. (2008). Hydrostatic compression model for sandy soils. *Canadian Geotechnical*
23 *Journal*, 45(8): 1169–1179. <https://doi.org/10.1139/t08-048>

24
25 Wang, X., Huo, Z., Guan, H., Guo, P., and Qu, Z. (2018). Drip irrigation enhances shallow
26 groundwater contribution to crop water consumption in an arid area. *Hydrological*
27 *Processes*, 32(6): 747–758. <https://doi.org/10.1002/hyp.11451>

28
29
30
31
32