

[Open Peer Review on Qeios](#)

[Commentary] Recognising the Rheumatological Needs of Neurodivergent Females

Ren Martin¹, Rachael Taylor², Clive Kelly³

1 Healios

2 Teesside University

3 Newcastle University

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

We experience life and interact with others in a multitude of ways. The term 'neurodivergence' refers to variations from what is considered typical or normal. Neurodivergence influences an individual's behaviour in social situations and is associated with atypical emotional responses. This can precipitate inequity and rejection. Neurodivergent women and girls experience many physical and psychological health issues, and musculoskeletal disorders account for a significant proportion of these. Research and education into neurodivergent conditions in females should inform the reassessment of clinicians' present approach to those who present with multiple unexplained symptoms. Obtaining official confirmation of a neurodivergent condition improves access to support services and helps them and their family better understand themselves and the challenges they face. This commentary highlights the increased risk of developing rheumatological disease for women and girls with neurodivergent conditions and suggests how clinicians might increase their awareness of this.

Ren Martin

BSc (Hons) Nursing

Adolescent Autism Unit, Teesside

Rachael Taylor

BA (Hons) English, MSC Psych

PhD Student in Psychology at Teesside University

Clive Kelly*

FRCP, MD

Consultant Physician and Rheumatologist

James Cook University Hospital and University of Newcastle upon Tyne

***Corresponding author:** Dr Clive Kelly, Email: cliveryton@gmail.com or clive.kelly2@nhs.net

Keywords: neurodivergence; autism; ADHD; hypermobility; rheumatoid arthritis; osteoporosis; fibromyalgia.

Key Lessons

1. Neurodivergent conditions are under-recognised among women and girls who often present differently to men.
2. Neurodivergent females are at increased risk of developing a wide range of rheumatological disorders.
3. These include fibromyalgia, hypermobility, rheumatoid arthritis, osteoporosis and connective tissue disease.

What is neurodivergence?

While 'neurodiversity' represents the wide variation in the approach people take when communicating and socialising^[1], the term 'neurodivergence' relates more specifically to differences in mental or neurological functioning from the standard norm. This includes autism, ADHD and Tourette's syndrome, and there is evidence that a milder form of neurodivergence accounts for dyslexia and dyspraxia^[2]. Social isolation and self-harm are common among women and girls with these conditions, and their physical symptoms may in part reflect emotional distress. If we are to better respond to their clinical needs, we need to improve our understanding of their way of thinking and behaving. The ability of women and girls with neurodivergence to describe their feelings may differ significantly from that of neurotypicals. They might struggle to verbalise their symptoms in a logical or consistent manner, and may find it hard to deal with questions that try to disentangle what can seem like a jumble of symptoms without a firm timeline. Their expectations from the clinical consultation might differ to that of the clinician, but it is important that doctors recognise this. We should concentrate on the issues that neurodivergent people describe, not the problems that doctors may perceive them to be^[3]. Confirmation of a neurodivergent condition improves access to support services and helps them and their family better understand

themselves and the challenges they face. This commentary highlights the increased prevalence of rheumatological disease in females with neurodivergent conditions, describes these associations and suggests how rheumatologists can help.

The link between neurodivergence and heightened healthcare needs

Although neurodivergent conditions have been perceived as being the province of males, recent evidence has emerged over the last ten years that challenges this assumption [4]. It appears that although females often receive their diagnosis later in life, ADHD and autism may be much more prevalent among women and girls than has been appreciated. This may be due to their tendency to camouflage their difficulties to avoid social rejection or alienation. Because of this, neurodivergent women are more likely to internalise their concerns, resulting in a suppression of their natural tendencies. Such behaviour can trigger intense anxiety, obsession and anger which may manifest itself in a variety of clinical expressions. They can exhibit heightened sensitivity to emotional and sensory stimuli and find that this causes increased muscle tension and tenderness, producing distress and discomfort [5]. Physical symptoms are probably partly genetic and partly environmental in origin. These cause pain and diminished function and need to be recognized and taken seriously. If these features are combined with difficulty in making and maintaining close contact with others, a sense of isolation or abandonment may ensue. Frequently this can cause neurodivergent women and girls to feel misunderstood and to lose faith in their own ability, and to lose confidence in other people's willingness to support them. Autonomic hypersensitivity can be associated with emotional instability and produce a perceived lack of object constancy [6]. Fatigue, muscle tenderness and despair may develop, precipitating a referral to mental health services, the pain team or a rheumatologist.

Autistic people die younger and are more likely to develop a wide range of disorders than their neurotypical peers during their lifetimes [7][8]. A recent review of the literature defined these disorders in detail [9]. Neurodivergent people appear to have worse health care outcomes if they have experienced significant adverse experiences during childhood. Females are at greater risk in this regard, and this might explain their increased use of health care resources [10]. A recent review explored the issues described by autistic women with regards to accessing medical care and identified impaired executive function, poor communication, and hypersensitivity as all being contributory [11]. Failure of health care workers to appreciate these problems led to lower levels of satisfaction and worsening health care outcomes [12]. Surprisingly little literature exists specifically on the needs of women with neurodivergent conditions, despite dysfunction of every organ system being listed among the many disorders they report. However, it is established that autistic females are over-represented among all women who seek support for many medical disorders, and it appears that this is especially true for musculoskeletal conditions.

The musculoskeletal healthcare needs of neurodivergent females

Hypermobility spectrum disorders include Ehlers-Danlos (EDS) which is strongly related to the presence of neurodivergence, especially in girls and women [13]. A reduced sense of balance and poor coordination are often also

observed and may add to the risk of injury, dislocation, or fracture. Formal assessment of the extent of hypermobility using Beighton's test is recommended to quantify joint laxity, which tends to reduce over time. Teaching proprioceptive exercises and the practice of non-weightbearing exercises such as swimming or cycling can be very helpful, especially when allied to the provision of supportive physiotherapy and joint care techniques. A minority of patients are at risk of vascular complications associated with more severe forms of EDS.

Chronic musculoskeletal pain with tenderness to touch and profound fatigue can lead to the suspicion of fibromyalgia in women. This is supported by the presence of autonomic hypersensitivity, which may manifest as postural orthostatic tachycardia syndrome, irritable bowel or recurrent headache. Patients frequently report trouble sleeping and complain of brain fog. Many such women have features of a neurodivergent condition [14] and there maybe also be a family history of autism, ADHD or fibromyalgia" [15][16]. Hypermobile EDS is also often associated with this symptom complex, and it has been postulated that hypermobility mediates the relationship between fibromyalgia and neurodivergence [13][14][16]. Adolescent females with neurodivergent features exhibit more pain than their peers [17], and one report suggested that 77% of young women with autism or ADHD had already manifested features of chronic pain by the age of 27 [18].

Systemic lupus erythematosus (SLE) and rheumatoid arthritis (RA) are among the auto-immune disorders reported with greater frequency in the first degree relatives of neurodivergent women [19]. There is also an increased prevalence of auto-immune conditions among the mothers of neurodivergent women [19][20][21][22]. In addition to maternal auto-immune thyroid disease, both RA [20] and SLE [21][22] are recorded with greater frequency in mothers of neurodivergent girls. While maternal RA is associated with a generalised increase in the risk of autism among progeny, maternal SLE is linked to an increased risk for autism only among Caucasian women [23]. The chances of developing both autism and ADHD are increased among the offspring of mothers with RA [24]. Postulated mechanisms for this include neonatal hypoglycaemia and maternal hyperglycaemia [25]. Chronic inflammatory conditions, such as lupus and rheumatoid arthritis, or the use of corticosteroids, may increase maternal glucose levels, exposing the embryo and the fetus to high-glucose concentrations. Intrauterine hyperglycemia, hyperinsulinemia or both may affect neurogenesis. Raynaud's phenomenon may predict or herald the development of a systemic rheumatological auto-immune disorder in neurodivergent female offspring of mothers with auto-immune disease and can be further exacerbated by stimulants prescribed for concomitant ADHD [26].

Children with autism exhibit lower bone mineral density across all sites when compared to neurotypical controls [27]. Osteoporosis is reported in women with ADHD, and this may be in part a consequence of prescribed stimulants [28]. Bone loss, or reduced bone formation, along with poor balance due to impaired proprioception, causes a much-inflated fracture risk at the hip, spine, and forearm in autistic females at all ages. Their odds ratio for hip fractures increases from 8.1 in girls to 24.8 in autistic women [29]. Other factors that may contribute to this increased fracture risk include vitamin D deficiency and intestinal dysbiosis from restrictive eating disorders in adolescence.

Table 1. Lists the common conditions affecting musculoskeletal health experienced by neurodivergent females.

To show the common musculoskeletal conditions experienced by neurodivergent females:

Hypermobility syndromes

Fibromyalgia

Rheumatoid arthritis

Connective tissue disease

Osteoporosis

Therapeutic challenges

Doctors have been slow to recognise the physical features that neurodivergent conditions can manifest in women. Not all females with autism or ADHD are aware of their condition, and those that are neurodivergent may choose not to disclose this. Historically, a label of psychosomatic illness has often been applied to women and girls who exhibit these features, with the concept that such patients can be hard to help. It is time we re-evaluated this approach and adopted a less judgemental attitude. Consistency and continuity of care are important in building the trust which neurodivergent people often struggle to achieve. If a diagnosis of a neurodivergent condition is suspected, access to appropriate multidisciplinary support may be appreciated whilst avoiding unnecessary cross-referrals.

A careful explanation of the diagnosis and its relationship to other aspects of the individual's symptomatology is especially important for neurodivergent women. They need to feel believed and validated in their symptomatology. Even if their condition proves difficult to treat, it is vital that they feel understood and supported both during the consultation and after. Sharing their story verbatim in a letter to their primary care physician can be helpful, especially if they also receive a copy. This helps to avoid misunderstanding and to ensure that agreed therapeutic plans are followed through. Obtaining consent for the involvement of other members of the multi-disciplinary team is important where this is proposed. Sharing selective and relevant information on the patient's condition with specialist nurses, physiotherapists or occupational therapists requires the patient's knowledge and consent and will likely improve the effectiveness of the intervention.

Many neurodivergent women describe difficulty in tolerating analgesia^[30]. They are more prone to side effects and may experience these at lower doses than neurotypical women^[31]. Starting with a low dose and titrating slowly upwards is a good rule. Certain analgesics also appear to be less effective in autistic females^[32], while other agents used off-licence are reported as being useful in relieving pain and distress^[33]. We may need to reconsider our definition of pain in the setting of neurodivergence, given that functional magnetic resonance imaging demonstrates different cerebral responses in neurodivergent females to their neurotypical peers. Likewise, immunomodulators may be effective at lower than usual doses in neurodivergent women and consensus between the rheumatologist and patient is essential to ensure effective compliance with drug therapy. Access to a helpline is much appreciated by neurodivergent women and girls, especially if there is the option to use email rather than insist on a telephone call^[34]. Depression may be associated with a sense of isolation, as well as with fatigue and despair. Neurodivergence is a risk factor for both anxiety and depression and antidepressants can improve quality of life^[35].

Within a rheumatological setting, it is important for clinicians to acquire an appreciation of the range of common disorders experienced by neurodivergent women and girls. The art of 'learning to listen' remains an essential tool in diagnosis. Neurodivergent people can feel uncomfortable if they are not given enough time to share their concerns, and an open unhurried dialog is more likely to facilitate a diagnosis. If patients are encouraged to share their lived experience, it becomes easier for the clinician to 'join the dots', which may allow the diagnosis of a neurodivergent condition to surface from what may have previously appeared to be a random collection of unrelated symptoms. Likewise, it is essential that clinicians are aware of the broad range of rheumatic conditions commonly experienced by neurodivergent women. If we are to become more effective at managing these conditions, breaking down barriers between services for physical and mental health would be a great help. This may allow the relationship between the limbic, endocrine, and immune systems in neurodivergent individuals to be more fully understood.

How patients and the public contributed to this article

Two of the authors of this paper have lived experience of female neurodivergence, and two authors work in the provision of health care delivery to females with neurodivergent conditions. Females with autism and / or ADHD specifically requested that we emphasise and explain their medical needs to Rheumatologists to improve their access to effective healthcare.

None of the authors have any commercial conflicts of interest to declare.

Each of the authors discussed, designed, wrote and reviewed the manuscript.

References

- ¹ *Singer, J. (1998). Odd People In: The Birth of Community Amongst People on the "Autistic Spectrum": a personal exploration of a New Social Movement based on Neurological Diversity. A thesis presented to the faculty of Humanities and Social Sciences in partial fulfilment of the requirements for the degree of Bachelor of Arts Social Science (Honours), Faculty of Humanities and Social Science, University of Technology, Sydney, 1998.*
- ² *Koi, P. (2021) 'Genetics on the neurodiversity spectrum: Genetic, phenotypic and endophenotypic continua in autism and ADHD', Studies in history and philosophy of science. Part A, 89pp. 52-62. 73*
- ³ *Anna Stenning & Hanna Bertilsdotter Rosqvist (2021) Neurodiversity studies: mapping out possibilities of a new critical paradigm, Disability & Society, 36:9, 1532-1537, DOI: 10.1080/09687599.2021.1919503*
- ⁴ *Young, S., Adamo, N., Ásgeirsdóttir, B.B. et al. Females with ADHD: An expert consensus statement taking a lifespan approach providing guidance for the identification and treatment of attention-deficit/ hyperactivity disorder in girls and women. BMC Psychiatry 20, 404 (2020). <https://doi.org/10.1186/s12888-020-02707-9>*
- ⁵ *Rynkiewicz A, Schuller B, Marchi E, Piana S, Camurri A, Lassalle A, Baron-Cohen S. An investigation of the 'female camouflage effect' in autism using a computerized ADOS-2 and a test of sex/gender differences. Mol Autism. 2016 Jan*

21;7:10. doi: 10.1186/s13229-016-0073-0. PMID: 26798446; PMCID: PMC4721191.

6. [^]Edvinsson D, Lindström E, Bingefors K, Lewander T, Ekselius L. Gender differences of axis I and II comorbidity in subjects diagnosed with attention-deficit hyperactivity disorder as adults. *Acta Neuropsychiatr.* 2013; 25:165-74.
7. [^]Hirvikoski T., Mittendorfer-Rutz E., Boman M., Larsson H., Lichtenstein P., Bölte S. (2016). Premature mortality in autism spectrum disorder. *The British Journal of Psychiatry: The Journal of Mental Science*, 208(3 Special Issue: Physical Health Across the Lifespan), 232-238. <https://doi.org/10.1192/bjp.bp.114.160192>
8. [^]Hand B. N., Angell A. M., Harris L., Carpenter L. A. (2020). Prevalence of physical and mental health conditions in Medicare-enrolled, autistic older adults. *Autism*, 24, 755-764. <https://doi.org/10.1177/1362361319890793>
9. [^]Weir, E., Allison, C., Warrier, V., & Baron-Cohen, S. (2021). Increased prevalence of non-communicable physical health conditions among autistic adults. *Autism*, 25(3), 681-694. <https://doi.org/10.1177/1362361320953652>
10. [^]Weiss J. A., Isaacs B., Diepstra H., Wilton A. S., Brown H. K., McGarry C., Lunsky Y. (2018). Health concerns and health service utilization in a population cohort of young adults with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 48(1), 36-44. <https://doi.org/10.1007/s10803-017-3292-0>
11. [^]Mason D., Ingham B., Urbanowicz A., Michael C., Birtles H., Woodbury-Smith M., et al. (2019). A systematic review of what barriers and facilitators prevent and enable physical healthcare services access for autistic adults. *Journal of Autism and Developmental Disorders*, 49(8), 3387-3400. <https://doi.org/10.1007/s10803-019-04049-2>
12. [^]Nicolaidis C., Raymaker D., McDonald K., Dern S., Boisclair W. C., Ashkenazy E., Baggs A. (2013). Comparison of healthcare experiences in autistic and non-autistic adults: A cross-sectional online survey facilitated by an academic-community partnership. *Journal of General Internal Medicine*, 28(6), 761-769. <https://doi.org/10.1007/s11606-012-2262-7>
13. ^{a, b}Eccles, J., Lodice, V., Dowell, N., and Owens, A. (2014). Joint hypermobility and autonomic hyperactivity: relevance to neurodevelopmental disorders. *J. Neurol. Neurosurg. Psychiatry* 85:8883. doi: 10.1136/jnnp-2014-308883.9
14. ^{a, b}Casanova EL, Baeza-Velasco C, Buchanan CB, Casanova MF. The relationship between autism and Ehlers-Danlos syndromes/hypermobility spectrum disorders. *J Pers Med.* 2020; 10:260.
15. [^]Reyero F, Ponce G, Rodriguez-Jimenez R, Fernandez-Dapica P, Taboada D, Martin V, et al. High frequency of childhood ADHD history in women with fibromyalgia. *Eur Psychiatry.* 2011;26: 482-3
16. ^{a, b}Kelly C, Martin R and Saravanan V. The links between fibromyalgia, hypermobility and neurodivergence. *Touch Reviews* March 15th 2022 <https://www.touchimmunology.com/fibromyalgia/journal-articles/the-links-between-fibromyalgia-hypermobility-and-neurodivergence/>
17. [^]Whitney DG, Shapiro DN. National Prevalence of Pain Among Children and Adolescents With Autism Spectrum Disorders. *JAMA Pediatr.* 2019 Dec 1;173(12):1203-1205. doi: 10.1001/jamapediatrics.2019.3826.
18. [^]Asztély K, Kopp S, Gillberg C, Waern M, Bergman S. Chronic Pain and Health-Related Quality Of Life In Women With Autism And/Or ADHD: A Prospective Longitudinal Study. *J Pain Res.* 2019;12:2925-2932 <https://doi.org/10.2147/JPR.S212422>
19. ^{a, b}Comi AM, Zimmerman AW, Frye VH, Law PA, Peeden JN. Familial clustering of autoimmune disorders and evaluation of medical risk factors in autism. *J Child Neurol.* 1999;14:388-94. doi: 10.1177/088307389901400608. - DOI - PubMed

20. ^{a, b}Sun CK, Cheng YS, Chen IW, Chiu HJ, Chung W, Tzang RF, Fan HY, Lee CW, Hung KC. Impact of parental rheumatoid arthritis on risk of autism spectrum disorders in offspring: A systematic review and meta-analysis. *Front Med (Lausanne)*. 2022 Nov 10;9:1052806. doi: 10.3389/fmed.2022.1052806. PMID: 36438039; PMCID: PMC9687371.
21. ^{a, b}Dalsgaard S. More Evidence Linking Autoimmune Diseases to Attention-Deficit/Hyperactivity Disorder. *JAMA Pediatr*. 2021;175(3):e205502. doi:10.1001/jamapediatrics.2020.5502
22. ^{a, b}Li DJ, Tsai CS, Hsiao RC, Chen YL, Yen CF. Associations between Allergic and Autoimmune Diseases with Autism Spectrum Disorder and Attention-Deficit/Hyperactivity Disorder within Families: A Population-Based Cohort Study. *Int J Environ Res Public Health*. 2022 Apr 8;19(8):4503. doi: 10.3390/ijerph19084503. PMID: 35457368; PMCID: PMC9025211.
23. [^]Zhu, Z., Tang, S., Deng, X. et al. Maternal Systemic Lupus Erythematosus, Rheumatoid Arthritis, and Risk for Autism Spectrum Disorders in Offspring: A Meta-analysis. *J Autism Dev Disord* 50, 2852-2859 (2020). <https://doi.org/10.1007/s10803-020-04400-y>
24. [^]Chiu, HJ., Sun, CK., Tsai, SJ. et al. A nationwide study of the risks of major mental disorders among the offspring of parents with rheumatoid arthritis. *Sci Rep* 12, 4962 (2022). <https://doi.org/10.1038/s41598-022-08834-5>
25. [^]Hoirisch-Clapauch S, Nardi AE. Autism spectrum disorders: let's talk about glucose? *Transl Psychiatry*. 2019 Jan 31;9(1):51. doi: 10.1038/s41398-019-0370-4. PMID: 30705254
26. [^]Umair HM, Sandler RD, Alunno A, Matucci-Cerinic M, Hughes M. Association between central nervous system stimulants used to treat attention deficit hyperactivity disorder (ADHD) and Raynaud's phenomenon: A scoping review. *Semin Arthritis Rheum*. 2021 Dec;51(6):1200-1204. doi: 10.1016/j.semarthrit.2021.09.002. Epub 2021 Sep 16. PMID: 34655948
27. [^]Rostami Haji Abadi, M., Neumeyer, A., Misra, M. et al. Bone health in children and youth with ASD: a systematic review and meta-analysis. *Osteoporos Int* 32, 1679-1691 (2021). <https://doi.org/10.1007/s00198-021-05931-5>
28. [^]Howard J, Walick K, Rivera J. Evidence of an Association between ADHD Medication and Diminished Bone Health in Children and Adolescents. Abstract 641 presented at 2016 Annual Meeting of the American Academy of Orthopaedic Surgeons, Orlando, Florida.
29. [^]Neumeyer AM, O'Rourke JA, Massa A, Lee H, Lawson EA, McDougale CJ, Misra M. Brief report: bone fractures in children and adults with autism spectrum disorders. *J Autism Dev Disord*. 2015 Mar;45(3):881-7. doi: 10.1007/s10803-014-2228-1. PMID: 25193141; PMCID: PMC4590779.
30. [^]Liu, J., Chen, L.L., Shen, S. et al. Challenges in the Diagnosis and Management of Pain in Individuals with Autism Spectrum Disorder. *Rev J Autism Dev Disord* 7, 352-363 (2020). <https://doi.org/10.1007/s40489-020-00199-7>
31. [^]Lévesque, M., Gaumont, I., & Marchand, S. (2011). Pain and autism. *Douleur Et Analgésie*, 24(3), 165-170. <https://doi.org/10.1007/s11724-011-0264-6>.
32. [^]Moore, D. J. (2014). Acute pain experience in individuals with autism spectrum disorders: A review. *Autism*, 19(4), 387-399. <https://doi.org/10.1177/1362361314527839>.
33. [^]Butwicka A, Långström N, Larsson H et al. Increased risk for substance use-related problems in autism spectrum disorders: a population-based cohort study. *J Autism Dev Disord*. 2017; 47: 80-89
34. [^]Livingston, L.A., Ashwin, C. & Shah, P. Electronic communication in autism spectrum conditions. *Molecular Autism*

11, 44 (2020). <https://doi.org/10.1186/s13229-020-00329-2>

35. [^]Liang SC, Sun CK, Fan HY, et al. Therapeutic effects of antidepressants for global improvement and subdomain symptoms of autism spectrum disorder: a systematic review and meta-analysis. *J Psychiatry Neurosci*. 2022 Aug 10;47(4):E299-E310. doi: 10.1503/jpn.210191. PMID: 35948343