

# Child and adolescent self-harm in a pandemic world: Evidence from a decade of data

David Cawthorpe<sup>1</sup>

<sup>1</sup> University of Calgary

**Funding:** No specific funding was received for this work.

**Potential competing interests:** No potential competing interests to declare.

## Abstract

### Background

Little is known about the COVID-19 pandemic impact on child and adolescent mental health, specifically self-harm. This paper serves to form a basis for understanding and planning an appropriate response to the present and longstanding child and adolescent mental health needs with global recommendations for integrated community support and disaster preparedness.

### Methods

Anonymous, aggregated data from an established regional child and adolescent addictions and mental health service was employed to examine differences in the rates of self-harm as the primary reason for referral among the health-seeking population represented by quarter by year from 2010-2022 to examine whether self-harm rates have increased since the onset of the COVID-19 pandemic.

### Results

Females and self-defined sex had higher rates of self-harm referrals compared to males. Both total and first-time self-harm referrals since the COVID outbreak in 2020 did not exceed the highest quarterly rates before COVID since 2010.

### Discussion

Since the COVID-19 pandemic, self-harm rates in one Canadian region remain stable and lower than the highest rates previously observed over the last decade. Given misplaced alarmist news and reports, a coherent, evidence-based, dynamic national response to mental health, social support, and disaster planning is required to fully understand how best to respond to the pandemic in general with a sustainable social support and disaster preparedness policy strategy and specifically the ongoing and pandemic-related mental health needs of the child and adolescent help-seeking population.

**Prof. Dr. David Cawthorpe**

Professor (Adjunct),

Cumming School of Medicine  
Departments of Psychiatry & Community Health Sciences  
Institute for Child & Maternal Health  
Alberta Children's Hospital Research Institute  
The University of Calgary  
email: [cawthord@ucalgary.ca](mailto:cawthord@ucalgary.ca)  
<https://research4kids.ucalgary.ca/profiles/david-cawthorpe>  
<https://www.wpanet.org/comorbidity>

Research and Evaluation,  
Child and Adolescent Addictions, Mental Health and Psychiatry Program,  
Alberta Health Services  
Richmond Road Diagnostic and Treatment Center  
2nd Floor, 1820 Richmond Road S.W.  
Calgary, Alberta T2T 5C7

### Acronyms

CAAMHPP: Child and Adolescent Addictions and Mental Health and Psychiatry Program

CDC: Centers for Disease Control

COVID-19: Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)

Std. Dev.: Standard Deviation

**Keywords:** Epidemiology, Self-Harm, Suicidality, Population, Child, Adolescent, Psychiatry, Mental Disorder, COVID.

## Introduction

In May 2021, the U.S. Centers for Disease Control (CDC) announced a post-COVID-19 surge of “suspected suicide attempt” presentations, especially among adolescent females, across a sample of emergency departments in the United States.<sup>[1]</sup> Even in Canada, report of projected suicide rates<sup>[2]</sup> before the fact of a coherent approach to defining estimated suicidality-related parameters may precipitate a ‘moral panic’ and increase the probability of reflexive, misplaced policy development, strategic planning, and investment.

The CDC announcement may have overestimated youth suicidal presentations to emergency departments. Additionally, the approach to pandemic policy with respect to child and adolescent mental health requires thoughtful guidance and rigorous examination of valid and reliable information from across Canada. There remains the ongoing need to focus on the longstanding service gaps around child and adolescent mental health, as with general mental health, health, and social

support in general<sup>[3][4]</sup>, that will likely transcend the global pandemic; issues not far removed from the requirement for a sustainable, nationally supported, dynamic, community-level, disaster preparedness policy strategy.

This paper presents an analysis since 2010 of self-harm referrals to regional Child and Adolescent Addictions and Mental Health and Psychiatry Program (CAAMHPP) services in Calgary, Alberta, wherein a standard 'self-harm' definition as the primary reason of referral among others is in place. In this region, self-harm includes both suicidal and non-suicidal forms of self-harm.<sup>[5]</sup>

## Methods

The study was a ten-year retrospective case series analysis of self-harm referrals to a regional, publicly funded, child and adolescent addiction and mental health and psychiatry program services that have been fully described.<sup>[6]</sup>

### *Study Setting*

The study was conducted in CAAMHPP services in Alberta Health Services, Calgary Zone.

### *Data source*

The regional access and intake system (RAIS)<sup>[6]</sup> is the CAAMHPP registration system containing the referral dates, demographics (sex), reasons for referral (self-harm) employed in the analysis. Aggregated, de-identified information represented in Tables 2 was extracted from the RAIS for the period including January 1st 20210 to March 18, 2021.

### *Analysis*

The aggregated data were employed to calculate the proportion of the total referrals by quarter that were self-harm referrals within the help-seeking population for each of two groups. One group was the total referrals for self-harm and the other group was the first-time or incident referrals for self-harm for each quarter (3-month intervals) for all years since 2010 until the end of the second quarter in 2022. Chi square analysis revealed the overall self-harm differences comparing males, females, and self-defined sex.

### *Ethical approval*

Data for this paper was collected under ethics ID-REB15-1057.

## Results


Table 1 shows the age and sex distribution of the total referrals and referrals for self-harm from 2010-2022. The self-harming groups for each sex tended to be older and more frequently female or self-defined. The groups are

proportionately different from one another (Chi Square  $p < .0001$ ).

**Table 1.** Age and sex distribution by group of the total referrals and referrals for self-harm from 2010-2022.

Referrals	Statistic	Male	Female	Self-Defined	Total
Total	Mean	11.12	12.95	14.33	12.03
	Std. Dev.	4.75	4.71	3.59	4.81
	n	50082	47811	761	98654
Self-Harm	Mean	14.48	15.1	14.66	14.89
	Std. Dev.	3.03	2.26	2.28	2.54
	n	20380	43772	2519	66671
Total	Mean	12.1	13.98	14.58	13.19
	Std. Dev.	4.58	3.89	2.64	4.29
	n	70462	91583	3280	165325

Figure 1 shows the self-harm proportion of the total unique referrals quarter by year of the proportion of the total and first-time referrals for self-harm.



**Figure 1.** Total unique and first-time quarterly referrals for self-harm by year.

From Figure 1, it may be observed that the proportion of self-harm referrals to CAAMHPP<sup>[6]</sup> increased in the first quarter of 2020 in comparison to 2019. However, since the COVID-19 pandemic outbreak up to the end of the second quarter of 2022, neither the proportions for total or first-time referrals have increased disproportionately compared to the maximum (higher) values over the previous ten years (e.g., 2013).

## Discussion

This study represents only one Canadian catchment area. There may be pockets throughout Canada where the self-harm (e.g., suicidality) might be different. What is called for is a coherent, multi-site, nation-wide approach to define self-harm and collect relevant data. This activity is a necessary precursor for developing a valid and reliable approach to examine whether and where there may have been an actual increase in the rate of self-harm (e.g., suicidality) following the pandemic outbreak of the COVID-19.

To understand and respond to the potential effect of the COVID-19 pandemic, a dynamic, locally integrated, national mental health, social support and disaster preparedness policy at minimum requires a three-fold approach:

1. National health-care integrated surveillance system for rapid measurement of mental and other disorders and diseases.<sup>[7]</sup>
2. Development of basic community-level social support and disaster preparedness.<sup>[8]</sup>
3. Address the long-standing unmet need and gaps of child and adolescent mental health services that in future may encompass more evolved and elastic responses to both local and global crises.<sup>[6]</sup>

## Conclusion

Innovations in child and adolescent mental health services in the present and future pandemic world may not be too far removed from innovations in community-level social support integrated with disaster preparedness planning. Also, the movement to providing online services may be here to stay for those that prefer this as a mode of service, along with the development of more futuristic virtual reality services. Both online and virtual services may help to take pressure off required tertiary face-to-face service encounters. To have coherent national, provincial, and local community-level targeted policy guiding programs that integrate local child-care, elder care, together with basic health, mental health, and social resources could necessarily also be a loci for disaster preparedness<sup>[8]</sup> in addition to sustainably shaping the demand of over-stressed and inadequate social, health, and specifically child and adolescent mental health services.<sup>[6]</sup>

## Limitations

The self-harm definition employed as the primary reason for referral in this study included both suicidal and non-suicidal self-harm. Other services not included in the dataset are available in the Calgary Zone for children and youth, so the reported self-harm rates likely underestimate the true rate of self-harm, even though the dataset does include visits to emergency departments within the catchment.

## Additional References

- Raballo, A., Poletti, M., Valmaggia, L. & McGorry, P. D. Editorial Perspective: Rethinking child and adolescent mental health care after COVID-19. *Journal of child psychology and psychiatry, and allied disciplines* (2020) doi:10.1111/jcpp.13371.
- Riva, G. Virtual reality in psychotherapy: review. *Cyberpsychology Behav. impact Internet, Multimed. virtual Real. Behav. Soc.* 8, 220–240 (2005).

## References

1. <sup>^</sup>Yard, E. et al. *Emergency Department Visits for Suspected Suicide Attempts Among Persons Aged 12–25 Years*

*Before and During the COVID-19 Pandemic — United States, January 2019–May 2021. MMWR Morb Mortal Wkly Rep* 70, 888–894 (2021).

2. <sup>a</sup> McIntyre, R. S. & Lee, Y. *Projected increases in suicide in Canada as a consequence of COVID-19. Psychiatry Res.* 290, 113104 (2020).
3. <sup>a</sup> Cawthorpe, D. R. L. *Children’s Mental Health 1954-2016 – who cares? (2016).*
4. <sup>a</sup> Russell, E. & Patrick, K. *Mental health needs our attention. CMAJ : Canadian Medical Association journal = journal de l’Association medicale canadienne* vol. 190 E34 (2018).
5. <sup>a</sup> Del Brío Ibáñez, P., Vázquez Fernández, M. E. & Imaz Roncero, C. *[Adolescent with non-suicidal self-harm in a psychosocial adversity environment]. Arch. Argent. Pediatr.* 117, e485–e488 (2019).
6. <sup>a, b, c, d, e</sup> Melathopolous, K. & Cawthorpe, D. *Impact of Central Intake Development and System Change on Per Capita Child and Adolescent Mental Health Discharges from 2002 to 2017: Implications for Optimizing System Design by Shaping Demand. Perm. J.* 23, (2019).
7. <sup>a</sup> Cawthorpe, D. *A comparative epidemiology model for understanding mental morbidity and planning health system response to the COVID-19 pandemic. Hear. Mind* 5, 103–111 (2021).
8. <sup>a, b</sup> Anonymous. *CanDoTopia: A Tax Reduction Model For Community Support And Disaster Preparedness. (Amazon, 2021).*