

[Open Peer Review on Qeios](#)

Child and adolescent self-harm in a pandemic world: Evidence from a decade of data

David Cawthorpe¹

¹ University of Calgary

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Background

Little is known about the COVID-19 pandemic impact on child and adolescent mental health, specifically self-harm. This paper serves to form a basis for understanding and planning an appropriate response to the present and longstanding child and adolescent mental health needs with global recommendations for integrated community support and disaster preparedness.

Methods

Anonymous, aggregated data from an established regional child and adolescent addictions and mental health service was employed to examine differences in the rates of self-harm as the primary reason for referral among the health-seeking population represented by quarter by year since 2010 to examine whether self-harm rates have increased since the onset of the COVID-19 pandemic.

Results

Female rates of self-harm referral were greater than male rates. Neither male nor female quarterly rates of total or first-time self-harm referrals exceeded the highest quarterly rates since 2010.

Discussion

Since the COVID-19 pandemic, self-harm rates in one Canadian region remain stable and lower than the highest rates observed over the last decade. Given misplaced alarmist news and reports, a coherent, evidence-based, dynamic national response to mental health, social support, and disaster planning is required to fully understand how best to respond to the pandemic in general with a sustainable social support and disaster preparedness policy strategy and specifically the ongoing and pandemic-related mental health needs of the child and adolescent help-seeking population.

Prof. Dr. David Cawthorpe

Professor (Adjunct),

Cumming School of Medicine

Departments of Psychiatry & Community Health Sciences

Institute for Child & Maternal Health
Alberta Children's Hospital Research Institute
The University of Calgary
email: cawthord@ucalgary.ca
<https://research4kids.ucalgary.ca/profiles/david-cawthorpe>
<https://www.wpanet.org/comorbidity>

Research and Evaluation,
Child and Adolescent Addictions, Mental Health and Psychiatry Program,
Alberta Health Services
Richmond Road Diagnostic and Treatment Center
2nd Floor, 1820 Richmond Road S.W.
Calgary, Alberta T2T 5C7

Acronyms

COVID-19: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)

CDC: Centers for Disease Control

CAAMHPP: Child and Adolescent Addictions and Mental Health and Psychiatry Program

Keywords: Epidemiology, Self-Harm, Suicidality, Population, Child, Adolescent, Psychiatry, Mental Disorder, COVID.

Introduction

In May 2021, the U.S. Centers for Disease Control (CDC) announced a post-COVID-19 surge of “suspected suicide attempt” presentations, especially among adolescent females, across a sample of emergency departments in the United States.^[1] Even in Canada, report of projected suicide rates^[2] before the fact of a coherent approach to defining estimated suicidality-related parameters may precipitate a ‘moral panic’ and increase the probability of reflexive, misplaced policy development, strategic planning, and investment.

The CDC announcement may have overestimated youth suicidal presentations to emergency departments. Additionally, the approach to pandemic policy with respect to child and adolescent mental health requires thoughtful guidance and rigorous examination of valid and reliable information from across Canada. There remains the ongoing need to focus on the longstanding service gaps around child and adolescent mental health, as with general mental health, health, and social support in general^{[3][4]}, that will likely transcend the global pandemic; issues not far removed from the requirement for a sustainable, nationally supported, dynamic, community-level, disaster preparedness policy strategy.

This paper presents an analysis since 2010 of self-harm referrals to regional Child and Adolescent Addictions and Mental Health and Psychiatry Program (CAAMHPP) services in Calgary, Alberta, wherein a standard ‘self-harm’ definition as the primary reason of referral among others is in place. In this region, self-harm includes both suicidal and non-suicidal forms of self-harm.^[5]

Methods

The study was a ten-year retrospective case series analysis of self-harm referrals to a regional, publicly funded, child and adolescent addiction and mental health and psychiatry program services that have been fully described.^[6]

Study Setting

The study was conducted in CAAMHPP services in Alberta Health Services, Calgary Zone.

Data source

The regional access and intake system (RAIS)^[6] is the CAAMHPP registration system containing the referral dates, demographics (sex), reasons for referral (self-harm) employed in the analysis. Aggregated, de-identified information represented in Tables 1 and 2 was extracted from the RAIS for the period including January 1st 20210 to March 18, 2021.

Analysis

The aggregated data were employed to calculate the proportion of the total referrals that were self-harm referrals within the help-seeking population for each of two groups. One group was the total referrals for self-harm and the other group was the first-time or incident referrals for self-harm for each quarter (3-month intervals) for all years since 2010.


Ethical approval

Data for this paper was collected under ethics ID-REB15-1057.

Results

The upper graph in Figure 1 shows the self-harm proportion of the total unique referrals quarter by year and the proportion of first-time referrals for self-harm.

Total Unique and First-time Quarterly Referrals for Self-Harm by Year



From Figure 1, it may be observed that the proportion of self-harm referrals to CAAMHP^[6] increased in the first quarter of 2020 in comparison to 2019. However, since the COVID-19 pandemic outbreak up to the end of the first quarter of 2022, neither the proportions for total or first-time referrals have increased disproportionately compared to the maximum (higher) values over the previous ten years (e.g., 2013).

Discussion

This study represents only one Canadian catchment area. There may be pockets throughout Canada where the self-harm (e.g., suicidality) might be different. What is called for is a coherent, multi-site, nation-wide approach to define self-harm and collect relevant data. This activity is a necessary precursor for developing a valid and reliable approach to examine whether and where there may have been an actual increase in the rate of self-harm (e.g., suicidality) following the pandemic outbreak of the COVID-19.

To understand and respond to the potential effect of the COVID-19 pandemic, a dynamic, locally integrated, national mental health, social support and disaster preparedness policy at minimum requires a three-fold approach:

1. National health-care integrated surveillance system for rapid measurement of mental and other disorders and

diseases.^[7]

2. Development of basic community-level social support and disaster preparedness.^[8]
3. Address the long-standing unmet need and gaps of child and adolescent mental health services that in future may encompass more evolved and elastic responses to both local and global crises.^{[6][9][10]}

Conclusion

Innovations in child and adolescent mental health services in the present and future pandemic world may not be too far removed from innovations in community-level social support integrated with disaster preparedness planning. Also, the movement to providing online services may be here to stay for those that prefer this as a mode of service, along with the development of more futuristic virtual reality services. Both online and virtual services may help to take pressure off required tertiary face-to-face service encounters. To have coherent national, provincial, and local community-level targeted policy guiding programs that integrate local child-care, elder care, together with basic health, mental health, and social resources could necessarily also be a loci for disaster preparedness^[8] in addition to sustainably shaping the demand of over-stressed and inadequate social, health, and specifically child and adolescent mental health services.^[6]

Limitations

The self-harm definition employed as the primary reason for referral in this study included both suicidal and non-suicidal self-harm. Other services not included in the dataset are available in the Calgary Zone for children and youth, so the reported self-harm rates likely underestimate the true rate of self-harm, even though the dataset does include visits to emergency departments within the catchment.

References

1. [^]Yard, E. et al. *Emergency Department Visits for Suspected Suicide Attempts Among Persons Aged 12–25 Years Before and During the COVID-19 Pandemic — United States, January 2019–May 2021*. *MMWR Morb Mortal Wkly Rep* 70, 888–894 (2021).
2. [^]McIntyre, R. S. & Lee, Y. *Projected increases in suicide in Canada as a consequence of COVID-19*. *Psychiatry Res.* 290, 113104 (2020).
3. [^]Cawthorpe, D. R. L. *Children's Mental Health 1954-2016 – who cares? (2016)*.
4. [^]Russell, E. & Patrick, K. *Mental health needs our attention*. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* vol. 190 E34 (2018).
5. [^]Del Brío Ibáñez, P., Vázquez Fernández, M. E. & Imaz Roncero, C. *[Adolescent with non-suicidal self-harm in a*

- psychosocial adversity environment]. *Arch. Argent. Pediatr.* 117, e485–e488 (2019).
6. ^{a, b, c, d, e}Melathopolous, K. & Cawthorpe, D. *Impact of Central Intake Development and System Change on Per Capita Child and Adolescent Mental Health Discharges from 2002 to 2017: Implications for Optimizing System Design by Shaping Demand.* *Perm. J.* 23, (2019).
 7. [^]Cawthorpe, D. *A comparative epidemiology model for understanding mental morbidity and planning health system response to the COVID-19 pandemic.* *Hear. Mind* 5, 103–111 (2021).
 8. ^{a, b}Anonymous. *CanDoTopia: A Tax Reduction Model For Community Support And Disaster Preparedness.* (Amazon, 2021).
 9. [^]Raballo, A., Poletti, M., Valmaggia, L. & McGorry, P. D. *Editorial Perspective: Rethinking child and adolescent mental health care after COVID-19.* *Journal of child psychology and psychiatry, and allied disciplines* (2020) doi:10.1111/jcpp.13371.
 10. [^]Riva G. *Virtual reality in psychotherapy: review.* *Cyberpsychol Behav.* 2005 Jun;8(3):220-30; discussion 231-40. doi: 10.1089/cpb.2005.8.220. PMID: 15971972.