

[Open Peer Review on Qeios](#)

Physiological Adaptation to Altitude: A Comparison of Fast and Slow Ascents to 5,300 m Above Sea Level

Clive Kelly¹, Shireen Saxena, Will Tran, George Wormauld, Rhu Muza, Annabel Lock, Carla Viveiros, Sami Touru, Eric Chen, Dominic Amato, Callina Lee, Shankar Bhattarai, Raj Simkhada, Julian Simeone, Charlotte Parsisson, Kieran Kelly

¹ Newcastle University

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Introduction: Rapid ascent to altitudes of over 5,000m above sea level are associated with dramatic changes in adaptive physiology. We describe the effects of a more gradual ascent on symptoms, oximetry and heart rate, and compare this approach to the effects of a more rapid ascent to the same altitude by a comparable cohort.

Methods: A group of 13 (6 females) representing 10 countries from 5 continents, ascended gradually from Lukla (2,300m) to Everest Base Camp (5,300m) in Nepal over an 8-day period, then descended over a further 4 days. We recorded all symptoms, and medication used, along with pulse oximetry and heart rate (HR) every 500m of ascent. We compared results with those obtained at equivalent altitudes using similar methodology from a fast ascent of Mount Kilimanjaro to an equivalent altitude by a comparable cohort over 4 days.

Results: Our group had a median age of 33 years (range 25-66), and all successfully completed the trek. No severe headache, vomiting, orthopnoea nor productive cough occurred, although minor nausea and mild headache was common. Baseline oximetry fell from a median of 96% (93-97%) to a median of 78% (53-86%) at 8 days but recovered to 94% (89-99%) inside 4 days. Corresponding HR rose from a baseline median of 72bpm (57-85) to a median of 103bpm (78-115) at 8 days, then recovered to 80bpm (54-94) after 4 days. Neither age nor gender correlated with outcomes. Individually, HR correlated inversely with oximetry, but we found no group correlation between these two variables. By contrast, a more rapid 4-day ascent from the same starting height, with similar baseline values for HR and oximetry, to the same final altitude was associated with more severe headache, breathlessness, and vomiting. The fast ascent was associated with a more marked reduction in oximetry to a median of 71% (52-76) and an increase in HR to a median of 110bpm (88-140). The fast ascent group also required significantly more medication and rated their experience as less enjoyable.

Discussion: Oxygen desaturation and tachycardia are inevitable consequences of ascending above 5,000m but the degree to which this occurs can be reduced by slowing ascent times and taking rest days every 1,000m of ascent. This practice is associated with fewer symptoms and greater safety, with less need for either prophylactic or therapeutic medication. Careful consideration should be given to rates of ascent when climbing to altitudes at or above 5,000m.

Introduction

The human body is exposed to many risks when ascending to high elevations. Adverse and changeable weather conditions, cold, high winds, and the risk of ice, rock fall or avalanche may be encountered even at relatively low altitudes. Likewise, the dangers posed by poor visibility, difficult terrain and dehydration are often under appreciated by climbers [1]. More predictable is the change in partial pressure of oxygen (pO₂) in inspired air, which can have profound effects on human physiology.

As altitude increases, atmospheric pressure falls and there is an associated reduction in pO₂. At sea level, atmospheric pressure is 760mmHg, giving an inspired oxygen tension of 148mmHg. Atmospheric pressure is reduced by half at 5500m above sea level to 380mm Hg, with a corresponding fall in pO₂ to 69mmHg. At an altitude of 8900m, the height of mount Everest, these values fall to around 30% of their sea level baseline. As the pressure of inspired oxygen decreases, this leads to reductions in alveolar and arterial oxygen pressures. Indeed, at 3000m arterial oxygen saturation is around 90% [2] and falls to around 80% in travellers in the great ranges (6000-8000m). At this height, the unacclimatised climber will be distressed, displaying breathlessness and significantly reduced exercise tolerance. The stress on the body brought upon by hypoxia is also influenced by the rate of ascent and the severity and duration of exposure [3].

Humans can acclimatise to the hypoxia encountered at altitude. In the acute phase following ascent, the reduction in pO₂ is detected by peripheral chemoreceptors in the carotid and aortic bodies, triggering a hyper-ventilatory response and increasing both tidal volume and respiratory rate [4]. The oxygen-haemoglobin dissociation curve shifts to the right, improving oxygen delivery to tissues, mediated by increased levels of 2,3 DPG and a reduction in tissue pH caused by anaerobic respiration. Hypoxic pulmonary vasoconstriction leads to improved matching of ventilation and perfusion within the lung. The subsequent elevation in pulmonary artery pressure may ultimately lead to right ventricular hypertrophy [5].

There are corresponding cardiovascular changes, with tachycardia, increased cardiac output and rising blood pressure mediated by increased sympathetic drive. Increased diuresis leads to a higher haematocrit as plasma volume reduces and blood becomes more concentrated. Despite these early adaptations, there is a reduction in cognitive function with hypoxia, which may contribute towards accidents and poor decision making at extremes of altitude.

Further acclimatisation occurs over several days. As minute ventilation remains high, the body slowly adapts to the resulting respiratory alkalosis by increasing renal excretion of bicarbonate. This compensatory mechanism takes about 100 hours and is accelerated by acetazolamide [6]. Lactate production falls as capillaries proliferate in skeletal muscle to maintain muscle function [7]. Increased production of erythropoietin by the kidney in response to hypoxia stimulates haematopoiesis. Haematological adaptation to altitude is complete when polycythaemia occurs. This takes 45 days at an altitude of 4000m, but the upper altitude limit of this adaptation has not yet been clearly defined [8]. However, haemoglobin concentrations can rise to 200g/l, and greater blood viscosity increases the risk of venous thromboembolism, retinal damage and stroke [9].

Failure of the human body to adjust adequately to the physiological challenges imposed by hypoxia can cause acute mountain sickness (AMS). Physical fitness does not predict the probability of developing AMS, while genetic factors also

play a part in susceptibility. Blood oxygen levels drop more at night, so ascending slowly and taking occasional 'rest days' to climb high and then return to lower elevations to sleep greatly reduces stress and facilitates acclimatisation. This approach protects against the development of AMS, but the effectiveness of this approach diminishes above 8,000m [2]. Acclimatisation increases the climber's sense of well-being and enjoyment, while improving their sleep and capacity for physical endurance.

Mountain medicine defines three altitude zones which correspond with reducing pO₂ and are associated with increasing risks of developing AMS [10]. These zones are high altitude (1,500-3,500 m), very high altitude (3,500-5,500 m) and extreme altitude (>5,500 m). Medical problems at these altitudes also include the risks of high-altitude pulmonary oedema (HAPE) and high-altitude cerebral oedema (HACE) [11]. Treatment with nifedipine and / or dexamethasone [11] may be required, while the risk of neurological damage is increased at extreme altitude [12]. People who develop AMS demonstrate alterations in anti-diuretic hormone (ADH) and those at risk of developing HAPE may notice reduced urine production prior to developing respiratory symptoms [13]. A reduction in the efficiency of digestion occurs at high altitude [14] and other adverse effects include dehydration, hypothermia, and sunburn. The present study was designed to compare the effects of a gradual ascent to very high altitude with those of a rapid ascent on the body's physiological responses.

Methods

Over twelve days in March 2024, a group of 13 people (6 females) from 10 countries across 5 continents, met in Kathmandu in Nepal to climb to Everest Base Camp. Our group had a median age of 33 years (range 25-66). Led by two male Nepali guides, we flew to Lukla, and where we were joined by a team of five porters who helped carry our luggage. We stayed in Tea Houses and had four season sleeping bags to cope with night-time temperatures which fell to -15C. We ascended gradually from 2,300m to Everest Base Camp at 5,300m, and then further up to Kala Patthar (5,700m) over an 8-day period. The average cumulative height gained each day was 500m, as we had two rest days when we climbed then descended, spending two consecutive nights at the same altitude. Given that our route was undulating, our total ascent amounted to 6,300m, with an average daily height gain of 788m. We then descended back to Lukla (2,300m) over a further 4 days, before flying back to Kathmandu. We recorded all the symptoms we experienced, and any medication we used. We measured each person's pulse oximetry and heart rate (HR) every evening at 6pm. The highest altitude at which recordings were taken was 5,300m. On each occasion we recorded the highest of three readings for oximetry and the lowest of three readings for HR using a finger probe (Oxypulse, Ecomerzpro, Madrid, Spain).

We compared the results from this trek with those obtained in February 2018 from a group of 7 people (4 female) who ascended Mount Kilimanjaro in Tanzania. This group had a median age of 29 years (range 25-60 years). We were led by a guide and assisted by 18 porters who carried our tents and all our food. We camped and required four season sleeping bags to cope with night-time low temperatures of -7C. We ascended rapidly, using the Machame route from a starting height of 1,800m to the summit at 5,900m, over a 4-day period. Our total ascent was therefore 4,160m, and the average cumulative height gained each day was 1,040m, with no rest days and no descent on the way to the summit. We then

descended back to 1,740m over a further day. We recorded all the symptoms we experienced, and any medication we used. We measured each person's pulse oximetry and heart rate (HR) every 500m of ascent, recording the highest of three readings for oximetry and the lowest of three readings for HR using a finger probe (Oxypulse, Ecomerzpro, Madrid, Spain). Again, the highest altitude at which recordings were taken was 5,300m. On this trek, we also measured systolic blood pressure (SBP) each evening at 6pm using an automatic sphygmomanometer (Vital Track, Blue Ocean Company Ltd., Lancs., UK) and recorded the lowest of three readings.

Comparisons between data sets from each group were made by Student's paired t test, while those between groups were made using Student's unpaired t test. Significance was expressed at the $p=0.05$ level.

Results

All members of both groups completed the treks to Everest Base Camp and to Kilimanjaro summit. Table 1 compares the symptoms experienced by both groups. None of the slow ascent group in Nepal developed moderate or severe AMS symptoms, but two met criteria for mild AMS (14%). The main symptoms reported by members of the slow ascent group were mild headache, fast heart rate, polyuria, sunburn, sinusitis and a dry cough. The latter two symptoms were not reported by those who wore masks to minimise dust inhalation. Four also reported mild peripheral oedema, affecting the hands, feet and face, above altitudes of 4,000m. Polyuria was also reported by four and was a significant inconvenience, especially at night.

Table 1. To compare the symptoms experienced by those ascending to 5,300M over 8 days (slow) versus those ascending over 4 days (fast), along with the altitude at which symptoms were first reported.

ALTITUDE (m)	SYMPTOMS (slow ascent)	SYMPTOMS (fast ascent)
1,8	-	-
2,3	-	-
2,8	-	Sunburn (2)
3,3	Chest infection (1)	Tachycardia (3); mild breathlessness (2)
3,8	Polyuria (4); sunburn (3)	Mild headache (7); peripheral oedema (7); polyuria (5)
4,3	Peripheral oedema (4); nausea (2)	Anorexia (7); moderate headache (7); diarrhoea (1)
4,8	Sinusitis (8); tachycardia (2); anorexia (2)	Nausea (7); severe headache (2); moderate breathlessness (3)
5,3	Mild headache (3); dry cough (3)	Vomiting (2); severe breathlessness (2)

Symptoms reported by the fast ascent group in Tanzania were more significant. Two people had severe headache, and one had recurrent vomiting with abdominal pain and became dehydrated. All seven reported anorexia and nausea, with moderate headache and at least mild breathlessness, and 5 (70%) met criteria for at least mild AMS. All had peripheral oedema.

Table 2 records the medication required by each group. Five of the slow ascent group used low dose acetazolamide, and a further three used ibuprofen. One person needed a course of antibiotics for a chest infection, and five required decongestants for sinusitis. By comparison, all of the fast ascent group took acetazolamide and five needed ibuprofen. Two were given low dose (2mg) dexamethasone and one needed nifedipine (20mg). No-one required oxygen but the rapid descent was mandated because of the severity of symptoms at the summit. The slow ascent group generally rated their experience as more enjoyable than those in the fast ascent group.

Table 2. To compare the medication required by those ascending to 5,300M over 8 days (slow) versus those ascending over 4 days (fast).

ALTITUDE (m)	MEDICATION (slow)	MEDICATION (fast)
1,8	Acetazolamide (5)	Acetazolamide (7)
2,3	-	-
2,8	-	Paracetamol (5)
3,3	Co-amoxiclav (1)	-
3,8	-	Ibuprofen (5)
4,3	Paracetamol (3)	Loperamide (1)
4,8	Decongestants (5)	Dexamethasone (2)
5,3	Ibuprofen (3); salbutamol (1)	Cyclizine (2); nifedipine (1)

Table 3 shows the results of the physiological measurements we undertook. In the slow ascent group, baseline oximetry fell from a median of 96% (93-97%) at 2,400m to a median of 78% (53-86%) at 8 days [t=7.29, SE=3.00; p=0.0001] but recovered to 94% (89-99%) inside 4 days on returning to 2,400m. Corresponding HR rose from a baseline of 72bpm (57-85bpm) to a median of 103bpm (78-115bpm) at 8 days [t=20.8, SE=1.28; p=0.001], then recovered to 80bpm (54-94bpm) after 4 days on returning to 2,400m. Neither age nor sex correlated with outcomes. Individually, HR correlated inversely with oximetry, but there was no group correlation between these two variables. By contrast, in the rapid ascent group, baseline oximetry fell from a median of 96% (94-98%) at 1,800m, to a median of 71% (52-76%) at 5,300m at 4 days [t=10.5, SE=2.96; p=0.0001], but recovered to 95% (92-98%) inside one day on descending to 1,800m. Corresponding HR rose from a median baseline of 70bpm (58-80bpm) to a median of 110bpm (88-140bpm) at 5,300m after 4 days [t=40.8, SE=3.95; p=0.0001], then recovered to 80bpm (54-94bpm) after a day on returning to 1,760m. Again, neither age nor sex correlated with outcomes. SBP rose from a median of 122mm Hg (98-138mm) at baseline to a median of 164 mm Hg (139-198mm) at 5,300m [t=18.8, SE=1.82; p=0.001], but returned to normal slowly over five days. No residual neurological features were noted after descending and careful screening showed no evidence of retinal haemorrhages in any of the group the day after descent.

Table 3. To compare heart rate and pulse oximetry in those ascending to 5,300M over 8 days (slow) versus those ascending over 4 days (fast).

ALTITUDE (m)	HEART RATE [HR] (bpm)		PULSE OXIMETRY [PO] (%)		P values	
	Slow	Fast	Slow	Fast	HR	PO
1,8	-	66 (48-72)	-	98 (95-99)	-	-
2,3	72 (57-85)	70 (58-80)	96 (93-97)	96 (94-98)	NS	NS
2,8	78 (58-95)	76 (60-88)	94 (90-98)	94 (91-97)	NS	NS
3,3	83 (67-103)	84 (78-105)	91 (83-94)	91 (88-95)	NS	NS
3,8	83 (63-102)	88 (70-104)	87 (81-94)	85 (74-92)	NS	NS
4,3	94 (70-112)	96 (76-120)	86 (74-94)	82 (71-89)	NS	NS
4,8	94 (69-116)	98 (81-118)	82 (55-93)	77 (58-86)	NS	p=0.048
5,3	103 (78-115)	110 (88-140)	78 (53-86)	71 (52-76)	NS	p=0.038

Although the rises in HR from baseline to 5,300m altitude were highly significant in both groups, the difference between the two groups in the degree of change in HR did not quite meet statistical significance. The falls in oxygen saturation were also highly significant in each group, while the degree of reduction in pulse oximetry was significantly greater in the fast ascent group than the slow ascent group [t=2.11, SE=4.78; p=0.048],

Discussion

Prolonged survival at altitude is certainly possible but the highest altitude at which people have been recorded as living long term is at 5,100m [15], although over 80 million people live at altitudes above 2,500 metres [16]. Different populations have evolved different adaptations to living at high altitude with genetic changes resulting over time [17]. Indigenous inhabitants not only survive but can thrive at high altitude because of evolutionary changes in their respiratory and cardiovascular systems [18][19]. Adaptation to altitude leads to genetic advantages such as larger lungs and greater exercise capacity [20]. Residents of the Andes exhibit polycythaemia [21], while Himalayan inhabitants compensate via increased ventilatory rates and cerebral blood flow [22] which together reduce the risk of AMS [23]. The chances of developing obesity [24] and cardiovascular disease [25] are also reduced among those who live at high altitude.

Hypothermia (a body core temperature of below 35.0 C) is a constant threat at any altitude, but the risk rises with increasing distance from the equator. Even in summer, the mean temperature at the summit of Everest is -19.0 C [26]. Hypothermia may produce confusion [27], which can evolve to hallucination and a false perception of warmth, leading to some climbers taking off their jackets [28] with fatal consequences. The risk of dehydration is accentuated by breathing cold air as moisture from the upper airways is required to warm inhaled air to body temperature [13]. Conversely sunburn is also very common at altitude as a result of increased ultraviolet light in the thin atmosphere [29]. Severe sunburn can accentuate nausea and fatigue and may accelerate fluid loss in the case of severe skin damage [30]. Both hypothermia and hypoxia are more likely to develop in those who visit high altitude areas, rather than in those who reside there

permanently.

Hypoxia is the main factor in the development of altitude illness which can be separated into three related categories: AMS, HAPE, and HACE [31]. AMS is the most common and diagnosis is based on recent ascent to high elevation with the development of headache in all sufferers, typically accompanied by anorexia, fatigue, insomnia or nausea. These symptoms usually develop within 12 hours of ascent to an elevation above 3,000m and often develop overnight. The severity of AMS can be quantified using the Lake Louise score [32]. The prevalence of AMS increases with altitude, and typically affects just 7% at 2,200 m but as many as 52% at 4,560 m [33]. Younger people appear more susceptible to AMS [34], as do females [35].

AMS usually resolves within a day if trekkers do not climb any higher [36]. AMS is usually self-limiting but can be treated with paracetamol or ibuprofen. Acetazolamide facilitates acclimatisation and can be used to treat AMS, but it is more typically used as a prophylactic agent. It causes a bicarbonate diuresis with a metabolic acidosis, increasing ventilatory drive and oxygenation [37]. It accelerates acclimatisation and reduces periodic breathing, which is common at night over 4,000 m. It may induce polyuria and paraesthesia of extremities. A dose of 125 mg twice daily, starting one day prior to ascent and continuing until descent commences, reduces the probability of developing AMS and can be used as treatment if AMS occurs in climbers not already taking it [38]. In more severe cases of AMS, dexamethasone is effective in providing rapid symptomatic relief [39]. In the most severe cases, oxygen supplementation provides immediate benefit but must be combined with a prompt reduction in height of at least 300m [40].

The incidence of severe AMS among all trekkers on Kilimanjaro has been shown to be 8.6% with over 1% of all climbers hospitalised [41]. An earlier study of 130 Finnish trekkers reported a prevalence of at least mild AMS of 75%, and a summit success rate of under 50% [42]. More serious manifestations of altitude illness may occur above 4,300m, with a rough incidence of HAPE occurring in 1% of climbers [43]. It typically manifests as a productive cough with frothy phlegm and associated dyspnoea at rest. In severe cases, sputum is often blood-stained. Oxygen saturation falls dramatically, and oximetry shows levels at least 10% below those recorded by healthy people at the same altitude, with values invariably below 70% and often much lower [44]. The commonest cause of death among trekkers on Kilimanjaro is HAPE [45]. Urgent descent is mandatory, with the use of supplemental oxygen if available. Nifedipine has long been known to be effective in preventing or reducing pulmonary oedema in the short term but does not replace the need for oxygen and descent from altitude [46]. Less common is HACE, which may be triggered by the worsening hypoxia associated with HAPE [47]. HACE is a medical emergency and may lead to confusion, drowsiness, and coma [48]. Urgent descent is mandated, along with the use of dexamethasone and oxygen, where available. Low dose dexamethasone also helps prevent HACE in susceptible individuals [49] and can save lives [50].

Our study showed that a gradual ascent to 5,300m over 8 days, incorporating 2 rest days where we slept at the same altitude, was associated with significant benefits when compared to a rapid ascent over 4 days without taking time to acclimatise. Two of the slow ascenders (14%) developed features of mild AMS, while 70% of those in the fast ascent group developed symptoms of at least mild AMS, with one person (14%) also exhibiting features of early HAPE. Achieving fluid balance posed a challenge for many climbers. Drinking adequate volumes of fluid to prevent dehydration was offset

by the unwanted developments of either peripheral oedema from fluid retention, or polyuria from reduced ADH secretion in many trekkers. Acetazolamide is a diuretic and therefore accentuates polyuria. The need for acetazolamide was less among slow ascenders (35%) than fast ascenders (100%), as was the requirement for ibuprofen and paracetamol. The use of nifedipine and dexamethasone was confined to the fast ascent group, and it is possible that a degree of dehydration might offer some protection against both HAPE and HACE. Oxygen saturations were significantly higher among those who ascended slowly, and this group also exhibited slightly lower pulse rates. Overall, those who ascended slowly adapted more effectively to increasing altitude and reported a more comfortable and enjoyable experience.

Key messages

1. Acute mountain sickness is commonly associated with rapid ascent to altitudes of 5,300m
2. Graduated ascent with rest days to allow acclimatisation reduces the body's physiological burden
3. Gradual rather than rapid ascent is associated with fewer symptoms and less need for medication.

Photographs



Photograph 1. The team at Everest Base Camp in 2024



Photograph 2. Some of the climbing team on Kilimanjaro in 2018

References

- [^] Johnson C, Anderson S, Dallimore J, Winser S, Warrell D. *Oxford Handbook of expedition and wilderness medicine*. Oxford University Press. p618 ISBN 978-0-19-929661-3
- ^{a, b} Grocott, M; Martin, DS.; Levett, D; McMorrow, R; Windsor, J; Montgomery, H. (2009). Arterial Blood Gases and Oxygen Content in Climbers on Mount Everest. *N Engl J Med*. 360 (2): 140–9. doi:10.1056/NEJMoa0801581. PMID 19129527
- [^] Cymerman, A; Rock, PB. *Medical Problems in High Mountain Environments. A Handbook for Medical Officers (Report)*. Vol. USARIEM-TN94-2. US Army Research Inst. of Environmental Medicine Thermal and Mountain Medicine Division Technical Report.
- [^] Young, AJ; Reeves, JT. (2002). *Human Adaptation to High Terrestrial Altitude (PDF)*. *Medical Aspects of Harsh Environments*. Vol. 2. Borden Institute, Washington, DC. CiteSeerX 10.1.1.175.3270.
- [^] Bärtsch, P; Gibbs, JSR (2007). Effect of Altitude on the Heart and the Lungs. *Circulation*. 116 (19): 2191–2202. doi:10.1161/CIRCULATIONAHA.106.650796. PMID 17984389.
- [^] Harris, NS; Nelson, SW (16 April 2008). Altitude Illness – Cerebral Syndromes. *EMedicine Specialties*
- [^] Martin, D; Windsor, J (1 December 2008). From mountain to bedside: understanding the clinical relevance of human acclimatisation to high-altitude hypoxia. *Postgraduate Medical Journal*. 84 (998): 622–627.

doi:10.1136/pgmj.2008.068296. PMID 19201935.

8. [^]Zubieta-Castillo, G.; Zubieta-Calleja, G.R.; Zubieta-Calleja, L.; Zubieta-Castillo, Nancy (2008). *Facts that Prove that Adaptation to life at Extreme Altitude (8842m) is possible. Adaptation Biology and Medicine. 5 (Suppl 5): 348–355.*
9. [^]Peacock A. *Oxygen at high altitude. BMJ. 1998 Oct 17; 317(7165): 1063–1066. doi: 10.1136/bmj.317.7165.1063*
10. [^]Luks AM, Auerbach PS, Freer L, et al. *Wilderness Medical Society Clinical Practice Guidelines for the Prevention and Treatment of Acute Altitude Illness: 2019 Update. Wilderness & Environmental Medicine. 2019;30(4_suppl): S3-S18. doi:10.1016/j.wem.2019.04.006*
11. ^{a, b}Hackett P, Roach R. *High-Altitude Illness. July 12, 2001 N Engl J Med 2001;345:107-114 DOI: 10.1056/NEJM200107123450206*
12. [^]Fayed, N; Modrego, P.J.; Morales, H (2006). *Evidence of brain damage after high-altitude climbing by means of magnetic resonance imaging. The American Journal of Medicine. 119 (2): 168.e1–6. doi:10.1016/j.amjmed.2005.07.062. PMID 16443427. Archived from the original (PDF) on 22 November 2010.*
13. ^{a, b}Anand, IS.; Chandrashekhar, Y. (1996). *Fluid Metabolism at High Altitudes.. In Marriott, B.M.; Carlson, S.J. (eds.). Nutritional Needs In Cold And In High-Altitude Environments: Applications for Military Personnel in Field Operations. Washington (DC): National Academies Press (US): Institute of Medicine (US) Committee on Military Nutrition Research.*
14. [^]Westerterp, Klaas (1 June 2001). *Energy and Water Balance at High Altitude. News in Physiological Sciences. 16 (3): 134–7. doi:10.1152/physiologyonline.2001.16.3.134. PMID 11443234. S2CID 26524828.*
15. [^]West, JB (2002). *Highest permanent human habitation. High Altitude Medical Biology. 3 (4): 401–7. doi:10.1089/15270290260512882. PMID 12631426.*
16. [^]Tremblay, JC; Ainslie, PN (2021). *Global and country-level estimates of human population at high altitude. Proceedings of the National Academy of Sciences of the United States of America. 118 (18): e2102463118. Bibcode:2021PNAS..11802463T. doi:10.1073/pnas.2102463118. PMC 8106311. PMID 33903258.*
17. [^]Azad P, Stobdan T, Zhou D, Hartley I, Akbari A, Bafna V, Haddad GG (December 2017). *High-altitude adaptation in humans: from genomics to integrative physiology. Journal of Molecular Medicine. 95 (12): 1269–1282. doi:10.1007/s00109-017-1584-7. PMC 8936998. PMID 28951950. S2CID 24949046.*
18. [^]Frisancho AR (1993). *Human Adaptation and Accommodation. University of Michigan Press. pp. 175–301. ISBN 978-0-472-09511-7.*
19. [^]Mayell H. (24 February 2004). *Three High-Altitude Peoples, Three Adaptations to Thin Air. National Geographic News. National Geographic Society.*
20. [^]Moore, LG (June 2001). *Human Genetic Adaptation to High Altitude. High Altitude Medicine & Biology. 2 (2): 257–279. doi:10.1089/152702901750265341. PMID 11443005.*
21. [^]Beall, CM. (1 February 2006). *Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude hypoxia. Integrative and Comparative Biology. 46 (1): 18–24. doi:10.1093/icb/icj004. ISSN 1540-7063. PMID 21672719.*
22. [^]Beall, CM.; Goldstein, M. C. (August 1987). *Hemoglobin concentration of pastoral nomads permanently resident at 4,850-5,450 meters in Tibet. American Journal of Physical Anthropology. 73 (4): 433–438. doi:10.1002/ajpa.1330730404. ISSN 0002-9483. PMID 3661681*

23. [^]Witt, Kelsey E.; Huerta-Sánchez, Emilia (22 July 2019). *Convergent evolution in human and domesticated adaptation to high-altitude environments*. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 374 (1777): 20180235. doi:10.1098/rstb.2018.0235. PMC 6560271. PMID 31154977.
24. [^]Voss, JD; Masuoka, P; Webber, BJ; Scher, AI; Atkinson, RL (2013). *Association of Elevation, Urbanization and Ambient Temperature with Obesity Prevalence in the United States*. *International Journal of Obesity*. 37 (10): 1407–12. doi:10.1038/ijo.2013.5. PMID 23357956.
25. [^]Faeh, D; Gutzwiller, F; Bopp, M (2009). *Lower Mortality From Coronary Heart Disease and Stroke at Higher Altitudes in Switzerland*. *Circulation*. 120 (6): 495–501. doi:10.1161/CIRCULATIONAHA.108.819250. PMID 19635973.
26. [^]Handford, C; Thomas, O; Imray, CHE (May 2017). "Frostbite". *Emergency Medicine Clinics of North America*. 35 (2): 281–299. doi:10.1016/j.emc.2016.12.006. PMID 28411928.
27. [^]Fears, J. Wayne (2011-02-14). *The Pocket Outdoor Survival Guide: The Ultimate Guide for Short-Term Survival*. Simon and Schuster. ISBN 978-1-62636-680-0.
28. [^]Brown DJ, Brugger H, Boyd J, Paal P (November 2012). "Accidental hypothermia". *The New England Journal of Medicine*. 367 (20): 19308. doi:10.1056/NEJMra1114208. PMID 23150960. S2CID 205116341.
29. [^]Luks AM, Hackett PH. *Medical conditions and high-altitude travel*. *N Engl J Med*. 2022;386(4):364–73.
30. [^]Blumthaler, M; Ambach, W; Ellinger, R (1997). "Increase in solar UV radiation with altitude". *Journal of Photochemistry and Photobiology B: Biology*. 39 (2): 130–134. doi:10.1016/S1011-1344(96)00018-8
31. [^]Bartsch P, Swenson ER. *Acute high-altitude illnesses*. *N Engl J Med*. 2013;369(17):1666–7.
32. [^]Roach RC, Hackett PH, Oelz O et al. *The 2018 Lake Louise acute mountain sickness score*. *High Alt Med Biol* 2018; 19:4–6.
33. [^]Maggiorini M, Böhler B, Walter M, Oelz O. *Prevalence of acute mountain sickness in the Swiss Alps*. *BMJ*. 1990;301(6756):853–855. <https://doi.org/10.1136/bmj.301.6756.853>.
34. [^]Gianfredi V, Albano L, Basnyat B, Ferrara P. *Does age have an impact on acute mountain sickness? A systematic review*. *J. Trav. Med*. 2020;27(6). <https://doi.org/10.1093/jtm/taz104>.
35. [^]Hou YP, Wu JL, Tan C, Chen Y, Guo R, Luo YJ. *Sex-based differences in the prevalence of acute mountain sickness: a meta-analysis*. *Mil Med Res*. 2019;6(1):38. <https://doi.org/10.1186/s40779-019-0228-3>.
36. [^]Hackett PH, Luks AM, Lawley JS, Roach RC. *High-altitude medicine and pathophysiology*. In: Auerbach PS, editor. *Wilderness medicine, 7th edition*. Philadelphia: Elsevier; 2017: 8–28.
37. [^]Luks AM, Swenson ER. *Medication and dosage considerations in the prophylaxis and treatment of high-altitude illness*. *Chest*. 2008;133(3):744–55.
38. [^]Roach RC, Lawley JS, Hackett PH. *High-altitude physiology*. In: Auerbach PS, editor. *Wilderness medicine, 7th edition*. Philadelphia: Elsevier; 2017. pp. 2–8.
39. [^]West, JB. (2012). "High-Altitude Medicine". *Journal of Respiratory and Critical Care Medicine*. 186 (12): 1229–1237. doi:10.1164/rccm.201207-1323CI. PMID 23103737.
40. [^]Huey RB.; Eguskitza X (2 July 2001). *Limits to human performance: elevated risks on high mountains*. *Journal of Experimental Biology*. 204 (18): 3115–9 doi:10.1242/jeb.204.18.3115. PMID 11581324.
41. [^]Crougths M, Nyakunga G, Sakita F, et al. *Incidence and predictors of severe altitude illness symptoms in Mt.*

Kilimanjaro hikers: a prospective cohort study. Journal of Travel Medicine, 2022, 1–8.

<https://doi.org/10.1093/jtm/taac044>

42. [^]Karinen H, Peltonen J, Tikkanen H. Prevalence of acute mountain sickness among finnish trekkers on Mount Kilimanjaro, Tanzania:an observational study. *High Alt Med Biol* 2008;9:301–6.
43. [^]Meier D, Collet TH, Locatelli I, Cornuz J, Kayser B, Simel DL, Sartori C. Does this patient have acute mountain sickness? The rational clinical examination systematic review. *JAMA*. 2017;318(18):1810–19.
44. [^]Burtscher M, Hefti U, Hefti J. High-altitude illnesses: Old stories and new insights into the pathophysiology, treatment and prevention. *Sports Medicine and Health Science* 3 (2021) 59–69
45. [^]Dekker M, Mremi A, Kilonzo K, et al. Altitude-Related Disorders on Mount Kilimanjaro, Tanzania: Two-Year Survey in a Local Referral Center. *Wilderness and Environmental Medicine* 2021; 32(1): 36–40
46. [^]Bartsch P, Maggiorini M, Ritter M et al. Prevention of High-Altitude Pulmonary Edema by Nifedipine. *N Engl J Med* 1991; 325:1284-1289. Doi: 10.1056/NEJM199110313251805
47. [^]Hackett PH, Roach RC. High altitude cerebral edema. *High Alt Med Biol*. 2004;5(2):136–46.
48. [^]Dekker M, Wilson M, Howlett W. Mountain Neurology: review. *Pract Neurol* 2019; 0:1–8. doi:10.1136/practneurol-2017-001783
49. [^]Imray C, Booth A, Wright A and Bradwell A. Acute altitude illness. *BMJ* 2011; 343 doi: <https://doi.org/10.1136/bmj.d4943>
50. [^]Krakauer, J (1999). *Into Thin Air: A Personal Account of the Mt. Everest Disaster*. New York: Anchor Books/Doubleday. ISBN 978-0-385-49478-6.