

[Open Peer Review on Qeios](#)

Perspectives on the Immune System in Sepsis

Felician Stancioiu¹, Bogdan Ivanescu, Radu Dumitrescu²

¹ Bio-Forum Foundation

² University of Bucharest

Funding: The author(s) received no specific funding for this work.

Potential competing interests: The author(s) declared that no potential competing interests exist.

Abstract

Beyond the modifications shown by the biochemistry labs, profound and ample modifications are seen in septic patients at a molecular level stemming from DNA translation and gene expression, manifested as unique profiles of mRNA (messenger), as well as non-coding, functional RNAs: miRNA (micro) and lncRNAs (long non-coding). Counteracting these modifications requires treatment with pleiotropic molecules and/or combination of molecules and opens the possibility of future treatments with arrays of siRNAs and/or specific panels of small molecules tailored for each patient subpopulation.

Felician Stancioiu, MD¹; Bogdan Ivanescu, MD², Radu Dumitrescu, MD, PhD^{3*}

¹Fundatia Bio-Forum, Bucharest, Romania

²Dr MIT, Bucharest, Romania

³University of Bucharest, Spitalul Medicover, Bucharest, Romania

**Corresponding author*

Dr. Radu Dumitrescu, Director Medical, Spitalul Medicover Bucuresti, Str Pechea nr 8

Email address: radu.dumitrescu@medicover.ro

Tel: 0722776783

Keywords: sepsis, immune system, NF- κ B, lncRNA, miRNA.

The recent, ongoing SARS-Cov2 pandemic has brought to the forefront the care of the critically ill infectious patients, initially mainly as a respiratory pathology - ARDS – but which has proven to be a multisystemic deterioration with extensive vasculitis affecting mostly every organ – lungs, the nervous system, liver, biliary tract, pancreas and the gastrointestinal tract, kidneys, heart, and the multisystem organ failure - MSOF – seen in some of the severely affected patients.

Many of the severe manifestations are due to an over-active, imbalanced immune system which produces a “cytokine storm” - an intense discharge of inflammatory molecules produced by activated neutrophils which is followed by over-

activation of the coagulation cascade with thrombembolism, ischemic tissue changes affecting organ function and ultimately cellular apoptosis.

In some patients these modifications are more ample and affecting more tissues especially as the infection is allowed to progress, making this aspect crucial: early, prompt treatment can impede infection progression to the point it overwhelms the immune system, since the ensuing modifications are multiple, simultaneously affecting various organs and leading to a very fragile functional state which is more difficult to normalize via external interventions (oxygen, iv medication, etc.).

Clinical diagnosis and sepsis biomarkers

Sepsis is defined as the simultaneous presence of 3 elements^{[1][2]}: i. systemic infection (identifiable blood pathogen); ii. dysregulated host response, and iii. life-threatening organ dysfunction; their presence can be identified with specific clinical signs and symptoms, cellular markers (enzymes released after destruction of cells), cytokines (the immune host response to infection), microbiologic markers (presence of specific bacteria or viruses) and cardiovascular modifications (SEPSIS-3).

Organ dysfunction alone is associated with patient mortality greater than 10% and can be identified with a qSOFA (quick Sequential Organ Failure Assessment) score of 2 or more points; qSOFA allows a rapid warning for sepsis by observing the presence of at least 2 of the following: 100 mm Hg or less for systolic blood pressure; altered mentation and a respiratory rate of 22/minute or more.

A Systemic Inflammatory Response Syndrome (SIRS) requires presence of two of more of the following five elements: i. hyper/hypothermia (>38° C or <36° C); ii. pulse > 90 bpm; iii. respiratory rate > 20/min; iv. white blood cells >12,000 or <4000/mm³ or >10% immature bands; v. PaCO₂ <32 mmHg^[3].

Presence of septic shock necessitates vasopressors to keep mean arterial pressure above 65 mm Hg, or lactate blood level greater than 18 mg/dL without hypovolemia, and is associated with a mortality rate greater than 40%.

Prompt diagnosis and early intervention in sepsis patients is essential, and towards this goal evaluation tools are used which quantify risks by using specific scores, especially in the emergency departments: qSOFA; SIRS; APACHE II - the Acute Physiology and Chronic Health Evaluation II; NEWS - the National Early Warning Score, and more recently REMS – the Rapid Emergency Medicine Score^[4].

These clinical tools allow accurate evaluation of risk scores for sepsis and prompt therapeutic intervention in patients admitted through emergency departments for in-hospital and 7-day mortality; however sepsis also occurs in hospitalized patients (10-40% of all sepsis cases in USA); clinical-based scores cannot differentiate between sepsis caused by bacterial vs viral infections to guide efficacious treatment. Blood cultures yield results after too many days when hours matter, and so new diagnostic tools are needed; promising new directions with excellent results are being developed by using mRNAs^[5], which will be discussed below.

Sepsis involves dynamic alterations of the immune system, mostly pro-inflammatory in initial phases and anti-inflammatory subsequently, but sometimes markers of both states are detected simultaneously^{[6][7]}; sepsis also entails modifications of coagulation, metabolic including hepatic, endocrine, cardiovascular including endothelial, nervous system both central and autonomic, renal^{[8][9]}. Most organs are affected at a cellular level via oxidative stress and ischemic modifications of mitochondria, peroxisomes, Golgi system, lysosomes and cell membranes^[10], nuclear transcription of

genes and also by the presence in the cytoplasm of interfering viral material. This complexity overlaps underlying individual characteristics such as age, comorbidities including infection source, trauma including surgery, and current medication.

The number and the dynamic modifications of the factors underlying sepsis pathobiology make the use of animal or computer models of sepsis superfluous and the task of finding biomarkers for sepsis staging, prognosis and treatment very challenging^[11].

Biomarkers most frequently used for sepsis risk stratification and prognosis are lactate, shown to be associated with mortality rates^[12]; fibrinogen^[13]; calprotectin^[14]; Macrophage migration inhibitory factor (MIF)^[15]; copeptin^[16]; Lipopolysaccharide-binding protein^[17]; most used are procalcitonin, which is elevated in sepsis patients but also in bacterial infections, and the nonspecific inflammation marker C-reactive protein (CRP)^[18].

A recent analysis of sepsis biomarkers^[19] revealed 258 such molecules, 28 of which were studied in clinical trials enrolling over 300 participants; 9 had better diagnostic value than either/both procalcitonin or C-reactive protein (CRP), 5 of which were identified in adult non-surgical patients: CD64^[20], heparin binding protein^[21], decoy receptor 3^[22]; Group II phospholipase A2 (PLA2-II)^[23]; sCD163^[24].

Evaluating the transcriptome, metabolome and proteome in sepsis patients with multimodal assays yields better results for risk stratification and prediction, especially when employed on specific populations: pediatric^[25], adults^[26]; this approach may also better differentiate sepsis from noninfectious pathologies like trauma, disseminated intravascular coagulation or pancreatitis.

Pathobiology

The most important and challenging aspect in sepsis is the inflammatory response, present not only as contrasting and biphasic pro-inflammatory and anti-inflammatory states, but also with interconnected and simultaneously active molecules and paths. A good exemplification is offered by the natural killer lymphocytes (NK), which can exert both actions via cytokines with pro-inflammatory actions (interferon-gamma IFN- γ , interleukin-1- IL-1, tumor necrosis factor – alpha -TNF- α) and anti-inflammatory cytokines such as IL-10, IL-4, transforming growth factor-beta - TGF- β ,^[27]; these pro/anti-inflammatory actions seem to be related to the severity and mortality of sepsis, and also the ensuing lymphopenia and immunoparalysis state^[28]. Regulatory mechanisms of inflammation involve multiple molecules and pathways which partially overlap, beginning with gene transcription as the essential component, followed by processing of mRNA, translation, phosphorylation, degradation; of these steps gene transcription and its activation/inhibition are the most important^[29]. Examples of genes undergoing transcription in early stages of inflammation are *Cxcl2*, *Ptgs2/Cox2*, *Tnfa*, and *Il1b*; genes expressed in later stages are *Ccl5*, *Il6*, *Il12b*, *Ifnb1*, *Nos2/INOS*, *Saa3* and *Marco*; inflammation pathways include the nuclear factor- κ B (NF- κ B), mitogen-activated protein kinase - MAPK, and JAK/STAT^[29].

NF- κ B is one of the most important molecules related to inflammation; when inactive it forms dimers located in the cytoplasm where it is bound by I κ B proteins; inflammatory signals such as TNF- α , IL-1 β , bacterial LPS, reactive oxygen species and Toll-like receptors (TLRs) cause I κ B degradation and translocation of the NF- κ B dimers to the nucleus where they promote specific gene transcription encoding for molecules which regulate cell survival, proliferation, differentiation, and also adhesion molecules, cytokines and various proteins^{[30][31]}.

Due to the complexity and overlapping modulation of the inflammatory pathways, a good strategy for getting a clear snapshot of the inflammatory milieu is to look for actively transcribed genes by analyzing the transcriptome for specific **mRNAs**.

Towards this goal was developed a blood-based diagnosis test which quantifies 29 host mRNAs and can differentiate infectious vs noninfectious pathologies, predict disease severity and 30-day mortality and differentiate bacterial vs viral infections^[32]. To detect presence of infection 11 mRNAs are used: the up-regulated genes *GNA15*, *BATF*, *C3AR1*; *CEACAM1*, *ZDHHC19*, and *C9orf95*, and the down-regulated genes *RPGRIP1*, *HLA-DPB1*, *KIAA1370*, *TGFB*; 7 mRNAs are used to distinguish bacterial from viral infections: *IFI27*, *JUP*, *LAX1*, *TNIP1*, *GPAA1*, *CTSB* and *HK3*; and 11 mRNAs can help predict the 30-day mortality risk: *LY86*, *TST*, *KCNJ2*, *HIF1A*, *SEPP1*, *C11orf74*, *CIT*, *DEFA4*, *CD163*, *RGS1*, and *PER1*.

This 29-mRNA test was shown to be significantly better at predicting clinical outcomes in sepsis patients than CRP levels, leukocyte and lymphocyte counts and Charlson comorbidity index and better than IL-6 and APACHE II scores^[5], and also has high accuracy in diagnosing bacterial and viral infections^[33].

Perhaps the most consequential recent development in human biology was the discovery of the fact that the majority of the human genome – more than 80% of the 3 billion nucleotides - have a non-coding, functional role (compared to less than 2% of nucleotides which comprises the known protein-coding genes), and also a majority of single nucleotide polymorphism (SNPs) which were linked to various pathologies via genome-wide association studies (GWAS) were not found in protein-coding exons; 88% of them are located in the intronic or intergenic sites (non-coding loci)^[34]. It was also shown that there are about two thousand different short RNA sequences with 19-22 nucleotides **micro RNAs (miRNAs)** and more than ten thousand long non-coding RNAs (lncRNAs) – which have more than 200 nucleotides – some of those linked to cellular metabolic pathways related to inflammation and ischemia and which play important roles in the immune response and sepsis^[35], as detailed below.

During sepsis specific miRNAs were shown to inhibit the TLR signaling pathway, modulate production of inflammatory cytokines, regulation of endothelial function and the vascular barrier, and in the later sepsis stages also are involved in apoptosis regulation, immunosuppression, coagulation cascade and organ dysfunction. miRNAs can help with better diagnosing and staging of sepsis, and more prompt and effective treatment^[36].

MiRNAs shown to be dysregulated in the peripheral blood of sepsis patients are: miRNA-182, miRNA-486, and miRNA-15a^[37]; miRNA-146a, miRNA-223, miRNA-16, and miRNA-150^[38]; miRNA-125a and miRNA-125b^[38]. Of these, miRNA-223 plasma levels was shown to correlate with inflammation markers and sepsis severity in multiple studies^{[38][39][40][41]}; miRNA-125b has better prognosis prediction and usefulness for disease management than miRNA-125a^[42].

miRNA-146a, miRNA-15a, miRNA-125b, and miRNA-146a prevent NF- κ B activation by inhibiting TRAF6 and IRAK^[38]. A drawback for miRNA-based markers was that the miRNA levels were different in sepsis populations with different characteristics and this must be taken into account in different studies^[38].

Long non-coding RNAs (lncRNAs) are present in all human cells and are specific to cell types and organs, and also to pathologies, with lower expression levels compared to mRNA, and also less sequence conservation^[43]. They are linked to key regulatory mechanisms of inflammation, especially controlling transcription of genes involved in the

inflammatory responses, enhancing or suppressing inflammation in both gene- and time- specific ways. They interact with RNA-binding proteins and participate in chromatin remodeling, function as scaffolds, guides, decoys or signals during gene transcription^[29].

Besides proteins, lncRNAs can also interact with DNA and other RNA through nucleotide base pairing or RNA folding which generate structural domains. Indeed, lncRNAs regulate gene expression at all levels: transcription, post-transcription, translation, post-translation and epigenetic^[44].

During the inflammatory responses lncRNAs play essential roles by regulating multiple pathways, modifying gene expression in inflammatory cells involved in proliferation and differentiation (eg M1/M2 macrophage polarization) including production and release of inflammation-related cytokines.

Furthermore, lncRNAs are dynamically regulated by specific stimuli, so the pattern of lncRNAs can unmask the cell's exposure to distinct inflammatory signals^[45].

Also lncRNAs are targets in various inflammatory pathways and have modified expression profiles in different cells during inflammation; such differences are seen in viral vs bacterial infection, differential TLR activation (TLR4 vs TLR2) and also same antigen – LPS – activating different pathways in different tissues – cardiomyocytes, endothelial cells, renal tubular epithelial cells, monocytes^[46].

TNF α regulates more than 50 pseudogene lncRNAs and hundreds of lncRNAs, responding very selectively to specific cytokines and antigens in a NF- κ B-dependent manner. TLRs recognize specific patterns from pathogens and have essential roles in innate and adaptive immunity; many lncRNAs are also regulated in response to specific stimuli. Gp96 Convergent responds only to TNF α . H2-T32/24AS only to TNF α and TLR3 agonists; Cox2 Divergent to pro-inflammation cytokines and TLR1-4 agonists; while HoxA11AS responds to TLR3 agonists and is down regulated by TNF α stimulation^[45].

lncRNAs can also act as anti-inflammatory factors, can limit/inhibit NF- κ B signaling and transcription of pro-inflammatory cytokines and can down-regulate inflammatory pathways. lncRNA-MEG3 inhibits production and actions of inflammatory molecules IL-1 β and TNF- α and p65/ NF- κ B and also macrophage apoptosis induced by LPS by modifying the levels of Bax/Bcl-2 proteins^[47].

Lethe is a pseudogene lncRNA with 697 bp long, induced specifically by TNF α and IL-1 β , which inhibits NF- κ B via negative feedback; it does so by binding to the NF- κ B p65 dimer and preventing its binding at target genes thus acting as a decoy. Lethe is upregulated by TNF α and IL-1 β , but not TLR agonists, indicating its involvement in inflammation, although not in the native immune response; its level decreases with age, a state associated with increased NF- κ B activity^[43].

Similar to Lethe, NKILA down-regulates NF- κ B-driven inflammation, but through a different mechanism: it blocks I κ B degradation, preventing NF- κ B translocation to the nucleus, (a post-translational modification). Yet another way of inhibiting NF- κ B activity, both basal and TNF- α -stimulated, is showed by lincRNA-p21 which sequesters p65 mRNA and slows translation of p65^[48].

In septic patients, silencing of lncRNA-5657, lncRNA-MALAT1 and lncRNA-THRIL can protect the lung from inflammation damage; limiting the mitochondrial damage associated with increased levels of lncRNA-NEAT1, lncRNA-HOTAIR and lncRNA -CRNDE can protect from cardiomyopathy, increases in plasma lncRNA-ATP13A4-8z correlated

with progression of injury to renal epithelium^[43]; lncRNA-NEAT1 correlated with increased liver injury^[49] but also decreased myocardial injury^[50]; while lncRNA-XIST prevents acute lung injury in sepsis via miR-16-5p^[51]. Figure 1 below shows various lncRNAs, miRNA and molecules acting on immune pathways and linked to aggravation or amelioration of sepsis.

LncRNAs, miRNAs and intracellular molecules associated with aggravating sepsis

CRNDE – miR-181-5p – TLR4


HOTAIR- miR-211

MALAT1 – hsa-miR-346 – SMAD3
– miR-23-3p

NEAT1– miR-370-3p - TSP-1
miR-211 - PI3K/AKT
miR-495-3 - STAT3
miR-125a-5p-TRAF6/TAK1
miR-125 – MCEMP1

PVT1 - p38 MAPK

Wfdc21 - STAT3/TLR4



LncRNA and intracellular molecules with positive effects on sepsis

MEG3 - NF-κB

Figure 1. Nucleic acids and biomolecules linked to sepsis progression or amelioration

Immune system modifications post sepsis

It was observed that the alterations of lymphocytes mirror the severity of sepsis; moderate sepsis with mortality below 10% is generally not followed by a decrease in the number or function of tissue-resident memory T cells (TRMs), the CD103+ cells able to detect antigens and activate IFN-γ production^[52].

Important modifications were observed in immune systems of patients who recovered from sepsis manifested as transient decreases in the number and function of lymphocytes (chronic immunoparalysis), especially of CD8+ T cells, also an altered transcription profile of memory CD8 T cells which make such patients more vulnerable to re-infections such as *Lysteria monocytogenes*^[28] and to intracellular pathogens^[27].

Other immune modifications post-sepsis are decreased number of CD4 and peripheral - TRM -lymphocytes and dendritic cells, which are essential for activation of T cells and clearing localized infections. Patients recovered from sepsis also showed a reduced antigen repertoire and cell pool composition and subpopulations, modification of CD4 T cells phenotype and function^[53]. The mechanism of the loss of B and CD4 T cells in sepsis seem to be caspase-9-mediated, mitochondria-

dependent apoptosis^[54].

Recovery of immune protection after lymphopenia parallels an increase in number in the circulatory memory CD8 cells (T_{circ}) but not TRM^[55] via a process named homeostatic proliferation (HP); it was shown that "HP-memory" cells have similar efficacy against infectious particles as antigen-experienced (conventional) CD8 cells^[56]. It was also shown that generation of competent HP-memory cells requires three elements: i. CD8+ / CD4(+) T cell interaction; ii. CD40L-CD40 interactions with host cells, and iii. presence and release of antigens from endogenous bacteria; absence of these factors yields non-functional HP-memory cells^[56]. The mechanism of the defective priming of HP-memory cells in the absence of CD4+ T cell interaction involves alteration of CD8+ cells with TNF-related apoptosis-inducing ligand (TRAIL) expression; these altered CD8+ cells will undergo apoptosis (activation-induced cell death) after stimulation by antigens. Interestingly, these defective CD8+ cells regain full function after treatment with IL-2, but not IL-7 nor IL-15^[57]. Levels of TRAIL were assessed in patients with sepsis and were shown to increase between days 3-7^[58] but also are inversely associated with sepsis severity and organ dysfunction^{[58][59]}.

However, recently it was shown during research in bone marrow transplantation that such alterations in leukocyte function are reversed in the presence of ascorbate and arginine^{[60][61]}.

Treatment-wise

Clinical trials for medications used in septic patients (ex. inhibitors of TNF-alfa, PAF, IL-1) had poor results when assessed globally^[57], and it was suspected that this indeed is due to patient heterogeneity, so populations were divided into specific phenotypes based on clinical characteristics and biomarkers, and in this way it was observed that specific phenotypes had benefits in morbidity and mortality which were statistically significant. Treatment with recombinant human thrombomodulin was useful only in sepsis patients who have high fibrinogen degradation products - FDP and D-dimer levels, and high mortality rates^[62]; treatment of septic patients with recombinant human activated protein C (rhAPC) also had different results in different patient phenotypes (with different levels of plasminogen activator inhibitor (PAI)-1 and d-dimer)^[63]. Another possible explanation for the poor results seen in therapies with inflammation-related molecules is the high complexity of pathways and the number of molecules involved in inflammation, which make therapies based on single molecules unlikely "one-size-fits-all" panacea; recently this was also seen in COVID-19 treatment, where combination of monoclonal antibodies showed better results than single molecules.

New molecules with pleiotropic actions in inflammation show good promise, and one such molecule is ethyl pyruvate, which was shown to ameliorate the sepsis-induced immunosuppression and protect against secondary infection^[64]; its main immunomodulatory actions are blocking the activation of NF-κB and ERK (Extracellular Signal Regulated Kinase) pathways^[65].

Another direction to improve immune function in sepsis is to add to existing treatments better support for the structural integrity and function of the intestinal epithelium with substances known to improve the levels of the tight junction proteins - claudin, zonula occludens (ZO)-1 and occludin. The digestive system is also endowed with the human body's largest immune organ, which is needed to separate the intestinal bacterial flora from the rest of the body and limit the entrance of ingested antigens. The dipeptide alanil-glutamine (AlaGln) administered iv was shown to improve the

levels of zonula occludens-1 (ZO-1) and claudin-5^[66]. Specifically in sepsis, levels of intestinal epithelial tight junction proteins claudin-3, zonula occludens (ZO)-1 and occludin were improved with administration of emodin^[67] and also other rhubarb monomers: rhein, daucosterol linoleate, 3,8-dihydroxy-1-methyl-anthraquinone-2-carboxylic acid, and 1-O-caffeoyl-2-(4-hydroxy-O-cinnamoyl)-D-glucose^[68].

Another strategy for improving organ function in sepsis, knowing that the glutathione levels are decreased^[69] is iv administration of glutathione for amelioration of the intracellular redox state and function of mitochondria.

Finally, because sepsis involves simultaneously multiple systems and organs we should prioritize treatment with molecules known to have multiple, simultaneous positive actions on the immune, cardiovascular, endocrine and metabolic levels; two such molecules are ascorbic acid and the amino-acid L-arginine.

Ascorbate and arginine are essential molecules for native and adaptive immunity; arginine is the precursor for nitric oxide – NO – an essential microbicide used by macrophages and neutrophils, also essential for activation and maturation of T cells; ascorbate buffers the oxidative stress resulting from the oxidative burst in leukocytes, helps maintain and restore mitochondrial function and thus impede apoptosis, antagonizes the inhibitory effects of 2,3 indoleamine deoxygenase (2,3 IDO) on T cells and together with arginine is essential for T cell maturation and function^[70].

Besides its essential role in maintaining immune function, ascorbate is essential for catecholamine synthesis in the adrenals and thus essential for maintaining the sympathetic tonus in the vascular system without which hemodynamic collapse ensues; is also essential for the structural integrity of organs and vasculature because its essential role in collagen formation which during inflammation is actively degraded by matrix metalloproteases,

Multiple clinical trials have showed the beneficial actions in septic patients of ascorbate (decreased mortality, decreased need for vasopressor and mechanical ventilation when administered at 3-10g/day)^[71] also 28-day mortality, number of ICU-free and hospital-free days at 50 mg/kg q 6 h^{[72][73]} and respectively arginine^{[74][75]}.

Ascorbate can be regarded as a “poster” molecule illustrating the fact that pathobiologic modifications in various organs and system in septic patients are taking place concomitantly even though they are studied separately and treatment results evaluated in clinical trials tend to offer only partial, precise but segmented answers in such complex pathologies where more focus should be placed on the pattern of modifications rather than the individual alterations. When we look beyond the usual molecules used as dynamic markers in sepsis (procalcitonin, lactate, CRP, etc) we observe profound changes in gene expression which drive those modifications while combined therapeutic interventions are usually beyond the scope of most sponsored clinical trials which usually study single new drugs. With these in mind we can say that an important direction for therapeutic progress in complex pathologies with high mortality rates may be represented by individually tailored treatments (ex. with arrays of siRNAs and/or panels of small-molecules) which address both simultaneously and comprehensively the modifications at the nuclear, cytoplasmic and organelle levels in cell function.

Equitable contributions in conceptualization, writing, correction, editing were observed.

No personal data is disclosed in this article.

References

- [^] M. M. Levy, M. P. Fink, J. C. Marshall, E. Abraham, D. Angus, et al. (2001). SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med.* Apr;31(4):1250-6. doi:10.1097/01.CCM.0000050454.01978.3B PubMed PMID: 12682500
- [^] M. Singer, C. S. Deutschman, C. W. Seymour, M. Shankar-Hari, D. Annane, et al. (2016). The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA.* doi:10.1001/jama.2016.0287 PubMed PMID: 26903338; PMCID: PMC4968574; PubMed Central PMCID: 4968574
- [^] (1992). American College of Chest Physicians/Society of Critical Care Medicine Consensus Conference: definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med.* PubMed PMID: 1597042
- [^] O. Ruangsomboon, P. Boonmee, C. Limsuwat, T. Chakorn, A. Monsomboon. (2021). The utility of the rapid emergency medicine score (REMS) compared with SIRS, qSOFA and NEWS for Predicting in-hospital Mortality among Patients with suspicion of Sepsis in an emergency department. *BMC Emerg Med.* 7;21(1):2. doi:10.1186/s12873-020-00396-x PubMed PMID: 33413139; PMCID: PMC7792356; PubMed Central PMCID: 7792356
- ^{a, b} S. C. Brakenridge, P. Starostik, G. Ghita, U. Midic, D. Darden, et al. (2021). A Transcriptomic Severity Metric That Predicts Clinical Outcomes in Critically Ill Surgical Sepsis Patients. *Crit Care Explor.* 14;3(10):e0554. doi:10.1097/CCE.0000000000000554 PubMed PMID: 34671746; PMCID: PMC8522866; PubMed Central PMCID: 8522866
- [^] R. S. Hotchkiss, G. Monneret, D. Payen. (2013). Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol.* 13(12):862–874.
- [^] W. J. Wiersinga, S. J. Leopold, D. R. Cranendonk, T. Poll. (2014). Host innate immune responses to sepsis. *Virulence.* 5(1):36–44.
- [^] M. Singer, V. Santis, D. Vitale, W. Jeffcoate. (2004). Multiorgan failure is an adaptive, endocrine-mediated, metabolic response to overwhelming systemic inflammation. *Lancet.* 364(9433):545–548.
- [^] K. N. Iskander, M. F. Osuchowski, D. J. Stearns-Kurosawa. (2013). Sepsis: multiple abnormalities, heterogeneous responses, and evolving understanding. *Physiol Rev.* 93(3):1247–1288.
- [^] R. S. Hotchkiss, P. E. Swanson, B. D. Freeman. (1999). Apoptotic cell death in patients with sepsis, shock, and multiple organ dysfunction. *Crit Care Med.* 27(7):1230–1251.
- [^] A. Kwan, M. Hubank, A. Rashid, N. Klein, M. J. Peters. (2013). Transcriptional instability during evolving sepsis may limit biomarker based risk stratification. *PLoS One.* 8(3).
- [^] M. E. Mikkelsen, A. N. Miltiades, D. F. Gaieski, M. Goyal, B. D. Fuchs, et al. (2009). Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock. *Crit Care Med.* doi:10.1097/CCM.0b013e31819fcf68 PubMed PMID: 19325467
- [^] A. Juutilainen, S. Hamalainen, K. Pulkki, T. Kuittinen, T. Nousiainen, et al. (2011). Biomarkers for bacteremia and severe sepsis in hematological patients with neutropenic fever: multivariate logistic regression analysis and factor analysis. *Leuk Lymphoma.* 52:2349–2355.

14. [^]N. Jonsson, T. Nilsen, P. Gille-Johnson, M. Bell, C. R. Martling, et al. (2017). *Calprotectin as an early biomarker of bacterial infections in critically ill patients: an exploratory cohort assessment. Crit Care Resusc.*
15. [^]K. Kofoed, O. Andersen, G. Kronborg, M. Tvede, J. Petersen, et al. (2007). *Use of plasma C-reactive protein, procalcitonin, neutrophils, macrophage migration inhibitory factor, soluble urokinase-type plasminogen activator receptor, and soluble triggering receptor expressed on myeloid cells-1 in combination to diagnose infections: a prospective study. Crit Care. 11:R38.*
16. [^]S. Battista, U. Audisio, C. Galluzzo, M. Maggiorotto, M. Masoero, et al. (2016). *Assessment of diagnostic and prognostic role of copeptin in the clinical setting of sepsis. Biomed Res Int. 2016(3624730).*
17. [^]Garcia Guadiana RL, Albaladejo Oton MD, Rebello Acebes S, Esteban TP, Hernando HA, et al. (2018). *Diagnostic accuracy of lipopolysaccharide-binding protein for sepsis in patients with suspected infection in the emergency department. Ann Clin Biochem. 55:143–148.*
18. [^]B. Hamade, D. T. Huang. (2020). *Procalcitonin: Where Are We Now? Crit Care Clin. Epub. Jan;36(1):23-40. doi:10.1016/j.ccc.2019.08.003PubMed PMID: 31733680; PMCID: PMC6866676; PubMed Central PMCID: 6866676*
19. [^]C. Pierrakos, D. Velissaris, M. Bisdorff, J. C. Marshall, J. L. Vincent. *Biomarkers of sepsis: time for a reappraisal. Crit Care. 2020. doi:10.1186/s13054-020-02993-5PubMed PMID: 32503670; PMCID: PMC7273821; PubMed Central PMCID: 7273821*
20. [^]J. Jamsa, T. Ala-Kokko, V. Huotari, P. Ohtonen, E. R. Savolainen, et al. (2018). *C-reactive protein, and procalcitonin in the identification of sepsis in the. J Crit Care. 43:139–142.*
21. [^]Y. Zhou, Z. Liu, J. Huang, G. Li, F. Li, et al. (2019). *Usefulness of the heparin-binding protein level to diagnose sepsis and septic shock according to Sepsis-3 compared with procalcitonin and C reactive protein: a prospective cohort study in China. BMJ Open. 9:e026527.*
22. [^]L. Gao, B. Yang, H. Zhang, Q. Ou, Y. Lin, et al. (2018). *DcR3, a new biomarker for sepsis, correlates with infection severity and procalcitonin. Oncotarget. 9:10934–10944.*
23. [^]T. L. Tan, Y. Y. Goh. (2017). *The role of group IIA secretory phospholipase A2 (sPLA2-IIA) as a biomarker for the diagnosis of sepsis and bacterial infection in adults-a systematic review. PLoS One. 12:e0180554.*
24. [^]L. Feng, X. Zhou, L. X. Su, D. Feng, Y. H. Jia, et al. (2012). *Clinical significance of soluble hemoglobin scavenger receptor CD163 (sCD163) in sepsis, a prospective study. PLoS One. 7:e38400.*
25. [^]H. R. Wong, N. Z. Cvijanovich, N. Anas. (2015). *Developing a clinically feasible personalized medicine approach to pediatric septic shock. Am J Respir Crit Care Med. 191(3):309–315.*
26. [^]R. J. Langley, E. L. Tsalik, J. C. Velkinburgh. (2013). *An integrated clinico-metabolomic model improves prediction of death in sepsis. Sci Transl Med. 5(195).*
27. ^{a, b}S. A. Condotta, D. Rai, B. R. James, T. S. Griffith, V. P. Badovinac. (2013). *Sustained and incomplete recovery of naive CD8+ T cell precursors after sepsis contributes to impaired CD8+ T cell responses to infection. J Immunol. doi:10.4049/jimmunol.1202379PubMed PMID: 23355736; PMCID: PMC3578009; PubMed Central PMCID: 3578009*
28. ^{a, b}I. J. Jensen, X. Li, P. W. McGonagill, Q. Shan, M. G. Fosdick, et al. (2021). *Sepsis leads to lasting changes in phenotype and function of memory CD8 T cells. Elife. doi:10.7554/eLife.70989PubMed PMID: 34652273; PMCID: PMC8589447; PubMed Central PMCID: 8589447*

29. ^{a, b, c}N. W. Mathy, X. M. Chen. (2017). Long non-coding RNAs (lncRNAs) and their transcriptional control of inflammatory responses. *J Biol Chem*. doi:10.1074/jbc.R116.760884PubMed PMID: 28615453; PMCID: PMC5535013; PubMed Central PMCID: 5535013
30. [^]Lenardo M. J., Baltimore D. (1989). NF- κ B: a pleiotropic mediator of inducible and tissue-specific gene control. *Cell*. 58:227–229.
31. [^]Chen F. E., Huang D. B., Chen Y. Q., Ghosh G. (1998). Crystal structure of p50/p65 heterodimer of transcription factor NF- κ B bound to DNA. *Nature*. 391:410–413.
32. [^]J. Ducharme, W. H. Self, T. M. Osborn, N. A. Ledebor, J. Romanowsky, et al. (2020). A Multi-mRNA Host-Response Molecular Blood Test for the Diagnosis and Prognosis of Acute Infections and Sepsis: Proceedings from a Clinical Advisory Panel. *J Med*. 7;10(4):266. doi:10.3390/jpm10040266PubMed PMID: 33297498; PMCID: PMC7762405; PubMed Central PMCID: 7762405
33. [^]W. Bauer, K. Kappert, N. Galtung, D. Lehmann, J. Wacker, et al. (2021). A Novel 29-Messenger RNA Host-Response Assay From Whole Blood Accurately Identifies Bacterial and Viral Infections in Patients Presenting to the Emergency Department With Suspected Infections: A Prospective Observational Study. *Crit Care Med*. 1;49(10):1664-1673. doi:10.1097/CCM.00000000000005119PubMed PMID: 34166284; PMCID: PMC8439671; PubMed Central PMCID: 8439671
34. [^]E. N. C. O. D. E. Project Consortium. (2012). An integrated encyclopedia of DNA elements in the human genome. *Nature*. 6;489(7414):57-74. doi:10.1038/nature11247PubMed PMID: 22955616; PMCID: PMC3439153; PubMed Central PMCID: 3439153
35. [^]T. N. Zhang, D. Li, J. Xia, Q. J. Wu, R. Wen, et al. (2017). Non-coding RNA: a potential biomarker and therapeutic target for sepsis. *Oncotarget*. 10;8(53):91765-91778. doi:10.18632/oncotarget.21766PubMed PMID: 29207683; PMCID: PMC5710963; PubMed Central PMCID: 5710963
36. [^]S. M. K. Kingsley, B. V. Bhat. (2017). Role of microRNAs in sepsis. *Inflamm Res*. 66(7):553–569. doi:10.1007/s00011-017-1031-9
37. [^]R. Ojha, R. Nandani, R. K. Pandey, A. Mishra, V. K. Prajapati. (2019). Emerging role of circulating microRNA in the diagnosis of human infectious diseases. *J Cell Physiol*. doi:10.1002/jcp.27127PubMed PMID: 30146762
38. ^{a, b, c, d, e}X. Shen, J. Zhang, Y. Huang, J. Tong, L. Zhang, et al. (2020). Accuracy of circulating microRNAs in diagnosis of sepsis: a systematic review and meta-analysis. *J Intensive Care*. doi:10.1186/s40560-020-00497-6PubMed PMID: 33292630; PMCID: PMC7607638; PubMed Central PMCID: 7607638
39. [^]D. Zhao, S. Li, J. Cui, L. Wang, X. Ma, et al. (2020). Plasma miR-125a and miR-125b in sepsis: Correlation with disease risk, inflammation, severity, and prognosis. *J Clin Lab Anal*. Feb;34(2):e23036. doi:10.1002/jcla.23036PubMed PMID: 32077163; PMCID: PMC7031612; PubMed Central PMCID: 7031612
40. [^]X. Wu, J. Yang, L. Yu, D. Long. Plasma miRNA-223 correlates with risk, inflammatory markers as well as prognosis in sepsis patients. *Medicine (Baltimore 2018)*. doi:10.1097/MD.00000000000011352PubMed PMID: 29979415; PMCID: PMC6076081; PubMed Central PMCID: 6076081
41. [^]W. Zhang, J. Jia, Z. Liu, D. Si, L. Ma, et al. (2019). Circulating microRNAs as biomarkers for Sepsis secondary to pneumonia diagnosed via Sepsis 3.0. *BMC Pulm Med*. 14;19(1):93. doi:10.1186/s12890-019-0836-4PubMed PMID:

31088429; PMCID: PMC6518454; PubMed Central PMCID: 6518454

42. [^]D. Zhao, S. Li, J. Cui, L. Wang, X. Ma, et al. (2020). Plasma miR-125a and miR-125b in sepsis: Correlation with disease risk, inflammation, severity, and prognosis. *J Clin Lab Anal.* Feb;34(2):e23036. doi:10.1002/jcla.23036PubMed PMID: 32077163; PMCID: PMC7031612; PubMed Central PMCID: 7031612
43. ^{a, b, c}C. Wang, G. Liang, J. Shen, H. Kong, D. Wu, et al. (2021). Long Non-Coding RNAs as Biomarkers and Therapeutic Targets in Sepsis. *Front Immunol.* doi:10.3389/fimmu.2021.722004PubMed PMID: 34630395; PMCID: PMC8492911; PubMed Central PMCID: 8492911
44. [^]Cuevas-Diaz Duran R, Wei H, Kim DH, Wu JQ. (2019). Invited Review: Long non-coding RNAs: important regulators in the development, function and disorders of the central nervous system. *Neuropathol Appl Neurobiol.* Oct;45(6):538-556. doi:10.1111/nan.12541PubMed PMID: 30636336; PMCID: PMC6626588; PubMed Central PMCID: 6626588
45. ^{a, b}N. A. Raponi, K. Qu, J. Zhang, M. Mikhail, R. M. Laberge, et al. (2013). A mammalian pseudogene lncRNA at the interface of inflammation and anti-inflammatory therapeutics. *Elife.* doi:10.7554/eLife.00762PubMed PMID: 23898399; PMCID: PMC3721235; PubMed Central PMCID: 3721235
46. [^]K. K. Singh, P. N. Matkar, S. Muhammad, A. Quan, V. Gupta, et al. (2016). Investigation of Novel LPS-Induced Differentially Expressed Long Non-Coding RNAs in Endothelial Cells. *Mol Cell Biochem.* 421:157–68. doi:10.1007/s11010-016-2797-8
47. [^]X. Pan, L. He. LncRNA MEG3 expression in sepsis and its effect on LPS-induced macrophage function. *Cell Mol Biol (Noisy-le-grand).* 2020. p. 33040826.
48. [^]F. Lu, Y. Hong, L. Liu, N. Wei, Y. Lin, et al. (2021). Long noncoding RNAs: A potential target in sepsis-induced cellular disorder. *Exp Cell Res.* 15;406(2):112756. doi:10.1016/j.yexcr.2021.112756PubMed PMID: 34384779
49. [^]C. C. Zhang, F. Niu. (2019). LncRNA NEAT1 promotes inflammatory response in sepsis-induced liver injury via the Let-7a/TLR4 axis. *Int Immunopharmacol.* 22(75). doi:10.1016/j.intimp.2019.105731
50. [^]S. M. Wang, G. Q. Liu, H. B. Xian, J. L. Si, S. X. Qi, et al. (2019). LncRNA NEAT1 alleviates sepsis-induced myocardial injury by regulating the TLR2/NF- κ B signaling pathway. *Eur Rev Med Pharmacol Sci.* 23(11):4898–4907.
51. [^]X. Song, L. Li, Y. Zhao, Y. Song. (2021). Down-regulation of long non-coding RNA XIST aggravates sepsis-induced lung injury by regulating miR-16-5p. *Hum Cell.* Sep;34(5):1335-1345. doi:10.1007/s13577-021-00542-yPubMed PMID: 33978928; PMCID: PMC8114023; PubMed Central PMCID: 8114023
52. [^]S. J. Moioffer, D. B. Danahy, S. Wall, I. J. Jensen, F. V. Sjaastad, et al. (2021). Severity of Sepsis Determines the Degree of Impairment Observed in Circulatory and Tissue-Resident Memory CD8 T Cell Populations. *J Immunol.* 1;207(7):1871-1881. doi:10.4049/jimmunol.2001142PubMed PMID: 34479943; PMCID: PMC8496966; PubMed Central PMCID: 8496966
53. [^]J. Cabrera-Perez, S. A. Condotta, V. P. Badovinac, T. S. Griffith. (2014). Impact of sepsis on CD4 T cell immunity. *J Leukoc Biol.* doi:10.1189/jlb.5MR0114-067RPubMed PMID: 24791959; PMCID: PMC4197564; PubMed Central PMCID: 4197564
54. [^]R. S. Hotchkiss, K. W. Tinsley, P. E. Swanson, Schmiegel RE Jr, J. J. Hui, et al. (2001). Sepsis-induced apoptosis causes progressive profound depletion of B and CD4+ T lymphocytes in humans. *J Immunol.* 1;166(11):6952-63. doi:10.4049/jimmunol.166.11.6952PubMed PMID: 11359857

55. [^]D. B. Danahy, R. K. Strother, V. P. Badovinac, T. S. Griffith. (2016). *Clinical and Experimental Sepsis Impairs CD8 T-Cell-Mediated Immunity. Crit Rev Immunol.* 36(1):57–74. doi:10.1615/CritRevImmunol.2016017098PubMed PMID: 27480902; PMCID: PMC5314458; PubMed Central PMCID: 5314458
56. ^{a, b}S. E. Hamilton, S. C. Jameson. (2008). *The nature of the lymphopenic environment dictates protective function of homeostatic-memory CD8+ T cells. Proc Natl Acad Sci U A.* 25;105(47):18484–9. doi:10.1073/pnas.0806487105PubMed PMID: 19020089; PMCID: PMC2587628; PubMed Central PMCID: 2587628
57. ^{a, b}S. A. Condotta, J. Cabrera-Perez, V. P. Badovinac, T. S. Griffith. (2013). *T-cell-mediated immunity and the role of TRAIL in sepsis-induced immunosuppression. Crit Rev Immunol.* 33(1):23–40. doi:10.1615/critrevimmunol.2013006721PubMed PMID: 23510024; PMCID: PMC3625932; PubMed Central PMCID: 3625932
58. ^{a, b}H. Yoo, J. Y. Lee, J. Park, J. H. Yang, G. Y. Suh, et al. (2020). *Association of Plasma Level of TNF-Related Apoptosis-Inducing Ligand with Severity and Outcome of Sepsis. J Clin Med.* 1;9(6):1661. doi:10.3390/jcm9061661PubMed PMID: 32492832; PMCID: PMC7356503; PubMed Central PMCID: 7356503
59. [^]E. J. Schenck, K. C. Ma, D. R. Price, T. Nicholson, C. Oromendia, et al. (2019). *Circulating cell death biomarker TRAIL is associated with increased organ dysfunction in sepsis. JCI Insight.* doi:10.1172/jci.insight.127143PubMed PMID: 31045578; PMCID: PMC6538332; PubMed Central PMCID: 6538332
60. [^]K. Scheit, G. Bauer. (2014). *Synergistic effects between catalase inhibitors and modulators of nitric oxide metabolism on tumor cell apoptosis. Anticancer Res.* 34(10):5337–5350.
61. [^]J. Manning, B. Mitchell, D. A. Appadurai. (2013). *Vitamin C promotes maturation of T-cells. Antioxid Redox Signal.* 19(17):2054–2067. doi:10.1089/ars.2012.4988
62. [^]D. Kudo, T. Goto, R. Uchimido, M. Hayakawa, K. Yamakawa, et al. (2021). *Coagulation phenotypes in sepsis and effects of recombinant human thrombomodulin: an analysis of three multicentre observational studies. Crit Care.* doi:10.1186/s13054-021-03541-5PubMed PMID: 33741010; PMCID: PMC7978458; PubMed Central PMCID: 7978458
63. [^]H. Zhao, J. N. Kennedy, S. Wang, E. B. Brant, G. R. Bernard, et al. (2021). *Revising Host Phenotypes of Sepsis Using Microbiology. Front Med Lausanne.* doi:10.3389/fmed.2021.775511PubMed PMID: 34805235; PMCID: PMC8602092; PubMed Central PMCID: 8602092
64. [^]W. Chen, J. Lian, J. J. Ye, Q. F. Mo, J. Qin, et al. (2017). *Ethyl pyruvate reverses development of Pseudomonas aeruginosa pneumonia during sepsis-induced immunosuppression. Int Immunopharmacol.* (v;52:61-69). doi:10.1016/j.intimp.2017.08.024PubMed PMID: 28863323
65. [^]Q. Huang, Y. Fu, S. Zhang, Y. Zhang, S. Chen, et al. (2020). *Ethyl pyruvate inhibits glioblastoma cells migration and invasion through modulation of NF-κB and ERK-mediated EMT. PeerJ.* doi:10.7717/peerj.9559PubMed PMID: 32742812; PMCID: PMC7380274; PubMed Central PMCID: 7380274
66. [^]M. Bartosova, R. Herzog, D. Ridinger, E. Levai, H. Jenei, et al. (2020). *Alanyl-Glutamine Restores Tight Junction Organization after Disruption by a Conventional Peritoneal Dialysis Fluid. Biomolecules.* 10(8). doi:10.3390/biom10081178
67. [^]Y. Li, R. Guo, M. Zhang, P. Chen, J. Li, et al. (2020). *Protective effect of emodin on intestinal epithelial tight junction barrier integrity in rats with sepsis induced by cecal ligation and puncture. Exp Ther Med.*

doi:10.3892/etm.2020.8625PubMed PMID: 32346413; PubMed Central PMCID: 7185184

68. [^]L. Wang, Y. L. Cui, Z. Zhang, Z. F. Lin, D. C. Chen. *Rhubarb Monomers Protect Intestinal Mucosal Barrier in Sepsis via Junction Proteins*. *Chin Med J Editor*. 2017. doi:10.4103/0366-6999.205855PubMed PMID: 28485323; PMCID: PMC5443029; PubMed Central PMCID: 5443029
69. [^]Gianni M. D. Biolo, Antonione PhD, Raffaella MD, Cicco. Marcello MD. *Glutathione metabolism in sepsis*, *Critical Care Medicine*: September 2007. p. 591-595. doi:10.1097/01.CCM.0000278913.19123.13
70. [^]F. Stancioiu, G. Z. Papadakis, S. Kteniadakakis, B. N. Izotov, M. D. Coleman, et al. (2020). A."A dissection of SARS CoV2 with clinical implications (Review)". *Int J Mol Med*. 46(2):489–508. doi:10.3892/ijmm.2020.4636
71. [^]Y. Wang, H. Lin, B. W. Lin, J. D. Lin. (2019). *Effects of different ascorbic acid doses on the mortality of critically ill patients: a meta-analysis*. *Ann Intensive Care*. 2019;9(1):58. doi:10.1186/s13613-019-0532-9
72. [^](2020). *Effect of Vitamin C Infusion on Organ Failure and Biomarkers of Inflammation and Vascular Injury in Patients With Sepsis and Severe Acute Respiratory Failure: The CITRIS-ALI Randomized Clinical Trial [published correction appears in JAMA*. *JAMA*. 2019;322(13):1261-1270. doi:10.1001/jama.2019.11825
73. [^]W. Y. Kim, J. W. Jung, J. C. Choi, J. W. Shin, J. Y. Kim. (2019). *Subphenotypes in Patients with Septic Shock Receiving Vitamin C, Hydrocortisone, and Thiamine: A Retrospective Cohort Analysis*. *Nutrients*. 2019;11(12):2976. doi:10.3390/nu11122976
74. [^]M. S. Gough, M. A. Morgan, C. M. Mack, D. C. Darling, L. M. Frasier. (2011). *The ratio of arginine to dimethylarginines is reduced and predicts outcomes in patients with severe sepsis*. *Crit Care Med*. 39:1351–1358.
75. [^]M. Visser, M. A. Vermeulen, M. C. Richir, T. Teerlink, A. P. Houdijk. (2012). *Imbalance of arginine and asymmetric dimethylarginine is associated with markers of circulatory failure, organ failure and mortality in shock patients*. *Br J Nutr*. 107:1458–1465.