

[Open Peer Review on Qeios](#)

# Constructing particle generator schemes including supersymmetry for cycles of string vibrations through genus of hypercomplex manifolds

Deep Bhattacharjee

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

## Abstract

Considering the origin of the particles an approach has been provided to show the origin of Bosons, Fermions along with supersymmetry (SUSY) from string vibrations in  $g \geq 1$  genus of a hypercomplex manifold through  $\eta \times \eta$  matrix for the generator  $\nabla_{(p,q)}$ .

**Deep Bhattacharjee**<sup>1,2,3,\*,\dagger</sup>

<sup>1</sup> *INS Research, Department of Geometry & Topology, India.*

<sup>2</sup> *Research Assistant for computation of Theoretical Actual Intelligence (ActI) Formalisms, CXAI Technologies Ltd., Cyprus.*

<sup>3</sup> *Electro-Gravitational Space Propulsion Laboratory, India.*

\*Email address: [itsdeep@live.com](mailto:itsdeep@live.com)

\daggerThe author holds no conflict of interests related to this paper.

**Keywords:** String Theory – Particles – Supersymmetry.

## Introduction

Hypercomplex manifold is an essential ingredient in string theory. The topological string story formalises the concept of compact Kähler manifolds having Ricci flatness that in essence is the Calabi-Yau manifolds where for the Calabi-Yau geometry; Calabi-Yau threefolds are non-trivial in string theoretic foundations. Being the fundamental structure of vibrations; those strings resonant in a specific pattern with each pattern gives rise to each category of particles. All that is needed is the genus of the hypercomplex manifolds to carry out the vibrations and the permutation cycles associated with them as each schemes of permutation corresponds to a generator for each class of particles, be it bosons, fermions or their supersymmetric partner. Thus, the necessity arrives to calculate a formalism that in a precise way formulate those vibration cycles and depict what categories of particle emerge from them. This paper typically focusses on this vibration category and schemes related to the particle generation scenario.

This paper focuses on the hypercomplex manifold with no such specific dimensions although in the cases of general M-Theory, a compact Kähler having Ricci flatness or Calabi-Yau is a hypercomplex manifold of six compactified dimensions. Therefore, generalizing the notion of dimensions the strings being considered here as fibres for a coherent formation goes 'in' and 'out' of the genera and the specific symphony of the vibration cycle with respected permutations gives rise to the particles be it boson, fermion or their supersymmetric partner.

This permutation cycles depend totally on the chirality as computed towards the methodological aspects of the paper where it is necessary to develop the formations as per four signatures of the chiral operator  $L$ ,

$$\langle +, -, \pm, \bar{\mp} \rangle$$

Where each of such signature when marks the initiation point then each type of particles arises where in case of a scheme considered here as  $\epsilon_{ij}$  when  $i = j$  then for the directions or chirality as taken  $\langle + \rangle$ , the cycle starts with even generates boson or  $\partial$  while for the chirality  $\langle - \rangle$ , the cycle in odd generates fermions or  $\bar{\partial}$ . The odd and even cycle is same in the matrix operation  $\eta \times \eta$  for  $\epsilon_{ij}$  when  $i = j$  in both the cases of  $L^+$  and  $L^-$  which shows the supersymmetric partner for,

$$\left\{ \begin{array}{l} \mathcal{L}^+ \rightsquigarrow \text{for every fermion } \bar{\partial} \text{ there is a boson } \partial \Rightarrow \bar{\partial}\partial \text{ as the initiation point} \\ \text{which alters in the next cycle and then returns back in the previous cycle} \\ \downarrow \\ \mathcal{L}^- \rightsquigarrow \text{for every boson } \partial \text{ there is a fermion } \bar{\partial} \Rightarrow \bar{\partial}\partial \text{ as the initiation point} \\ \text{which alters in the next cycle and then returns back in the previous cycle} \end{array} \right. \quad \text{Potential is } \rho$$

Whereas the alternative form is considered for the same matrix  $\eta \times \eta$  for  $\epsilon_{ij}$  for  $i \neq j$  which shows the chiral operator  $L$

having signatures,

$$(\pm, \mp)$$

Extend the notion of the potential having the vibration cycle such that the derivative of the potential  $\rho$  is extended into such a state that there is no such supersymmetric formulations or particle formations – that is to say,

$$\left\{ \begin{array}{l} \mathcal{L}^{\pm} \rightsquigarrow \text{extends } \mathcal{L}^{\pm} \text{ and ultimately gives rise to only bosons } \partial\bar{\partial} \text{ upto } k \\ \downarrow \\ \mathcal{L}^{\mp} \rightsquigarrow \text{extends } \mathcal{L}^{\mp} \text{ and ultimately gives rise to only fermions } \bar{\partial}\partial \text{ upto } k \end{array} \right. \quad \exists \text{potential is } \rho^{-1}, \rho^{+1}$$

Where two peculiar instances of particle generation are shown for the chiral operator  $\mathcal{L}$  in the same matrix  $\eta \times \eta$  up to  $k$  such that for the chiral signatures,

$$(+ +, - -)$$

$$\left\{ \begin{array}{l} \mathcal{L}^{++} \rightsquigarrow \text{generates only bosons upto } k \\ \downarrow \\ \mathcal{L}^{--} \rightsquigarrow \text{generates only fermions upto } k \end{array} \right. \quad \exists \text{potential is } \rho^{+0}, \rho^{-0}$$

The detailed calculations will be computed in the next section.

## Methodology

A topological hypercomplex manifold  $T^*$  having genus  $g \geq 1$  such that  $M \subset T^*$  where  $M$  represents the hyperbolic ring of a genus  $g$  for the codimension inside the genus being represented as  $d$  which in essence is <sup>[1][2][3][4][5]</sup>,

$$g = \{\phi\}$$

There exists the  $n$  – dimensions of the hypercomplex manifold  $T^*$  having the spacetime coordinate  $(\sigma, \rho)$  it can be shown that there exists a defective unit  $\epsilon$  for each genus having a signature  $(\sigma, \rho)|_{\epsilon}$  where  $\epsilon$  depends on odd or even cycles of each strings  $s$  having a summation of their coherent structure to be represented as <sup>[6][7][8]</sup>,

$$\sum_s$$

Determines the genus signature of  $n$  – dimensions for the defective unit  $\epsilon$  taking the sub  $i, j$  for a scheme  $\mu$  such that <sup>[9][10]</sup>,

$$\ln \epsilon_{ij} = \begin{cases} i = j \text{ for } \mu = \text{odd or even} \\ i \neq j \text{ for } \mu = \text{reciprocal} \end{cases}$$

Where the genus signature can be spotted over a multiple sequence of 'in' and 'out' fibrations in such a way that <sup>[1][12][13]</sup>,

For the coherent fibre  $O$  which in essence determines the summation of the strings  $\sum_s$  there exists a factor of coherent fibre length  $J$  that determines the length of the cycle  $\ell$  that is exactly the length of the fibre determines the length of the 'in' and 'out' cycle between the genus  $g \geq 1$  to input the defective unit  $\epsilon_{ij}$  where this can be easily shown that the cycle  $\ell$  is given for the matrix  $\eta \times \eta$ ,



$$\Rightarrow \text{Tr}_{(\partial + \bar{\partial}, \dots, k)} \frac{1}{2} \frac{1}{(2\pi i)} \oint_{\rho} \oint_{\rho} \oint_{\rho} \dots \oint_{\rho_k} dJ dJ \dots dJ \approx \partial \rho$$

Where  $L^+ \equiv L^-$  for  $\epsilon_{ij}$  for  $i = j$  where  $L^-$  can be shown to have the opposite identity but still equal in terms of defect unit for the matrix depiction,





evolution  $\{T^\infty \nearrow\}$  governing the process of spacetime  $(\sigma, \rho)$  with normal space  $\sigma$  and frozen space  $l_\sigma$ ; more precisely a mixture to be given where there are two possibilities,

1.  $l_\sigma \not\subseteq \sigma$
2.  $l_\sigma \subseteq \sigma$

Therefore, the frozen in either of the Points [1 or 2] whichever takes place can be represented via the below matrix,

$$\sigma, \rho|_{l_\sigma} \cong \left\{ \begin{array}{l} \text{for chirality } \mathcal{L}^{++} \left( \begin{array}{cccc} \ddots & & & \\ & \partial\partial & & \\ & & \partial\partial & \\ & & & \partial\partial \\ & & & \ddots \\ & & & & k \end{array} \right) \xrightarrow{\Rightarrow} \partial\rho^{+0} \\ \\ \text{for chirality } \mathcal{L}^{--} \left( \begin{array}{cccc} \ddots & & & \\ & \bar{\partial}\bar{\partial} & & \\ & & \bar{\partial}\bar{\partial} & \\ & & & \bar{\partial}\bar{\partial} \\ & & & \ddots \\ & & & & k \end{array} \right) \xrightarrow{\Rightarrow} \partial\rho^{-0} \end{array} \right.$$

Thus, the cycle can be denoted as,

$$\begin{array}{ccccccccccc} \rightarrow & \rightarrow & L^+ & \rightarrow & \rightarrow & L^\pm & \rightarrow & \rightarrow & L^{++} & \rightarrow & \rightarrow \\ \uparrow & & & & & & & & & & \downarrow \\ \uparrow & & & & & & & & & & \downarrow \\ \uparrow & & & & & & & & & & \downarrow \\ \leftarrow & \leftarrow & L^- & \leftarrow & \leftarrow & L^{--} & \leftarrow & \leftarrow & L^{\mp} & \leftarrow & \leftarrow \end{array} \cong \Sigma l_{(p,q)}$$

Where  $(p, q)$  denotes the + and – respectively in all of the chirality where in cases the time coordinate  $\phi$  appears because of the infinite temporal evolution  $\{T^\infty \nearrow\}$  to formulate  $l_{(p,q)|\rho}$ .

Here in the genus signature  $(\sigma, \rho)|_\epsilon$  – there is a differential operator  $\epsilon$  for the entire hypercomplex manifold  $T^*$  : this particular operator singles out each genus from the previous genus and act correspondingly over a summation of all genera with respect to the fibration cycle  $\Sigma l_{(p,q)} \equiv \psi$  for each genus such that thee exists the above depicted chain where the independency of the genus is preserved with the connecting operator – the gap between two hyperbolic rings of the genus<sup>[1][8][13][14]</sup>,

$$\sum_{\kappa \equiv m=1}^{\infty} g_m - \sum_{n=1}^{\infty} g_n \exists m \neq n$$

For the differential operator  $\epsilon$  to take the value that has been closed through,

$$\psi \in (\Sigma \kappa) \int_{\epsilon(\Sigma \kappa)} / \sim$$

Thus, for,

$$\int_{\psi \text{ with } \psi \in (\Sigma \kappa)} \int_{\epsilon(\Sigma \kappa)} / \sim$$

It is easy to determine the  $(p, q)$  norms of the vibration cycle  $\Sigma / l_{(p,q)} \equiv \psi$  one can determine the types of particles being generated where this can be deduced as the generator of  $(p, q) \nabla_{(p,q)}$  for  $p = +$  and  $q = -$  where if we take  $\partial \equiv p = +$  and  $\bar{\partial} \equiv q = -$ : The generator can be introduced via the  $\eta \times \eta$  matrix – one gets the boson, fermion and supersymmetry as,

$$\left\{ \begin{array}{l} \nabla_{(p,q)} \implies \left\{ \begin{array}{l} \partial \partial \quad \hookrightarrow \quad \text{generates Boson} \\ \bar{\partial} \bar{\partial} \quad \hookrightarrow \quad \text{generates Fermion} \\ \partial \bar{\partial} \quad \hookrightarrow \quad \text{generates Fermion for every Boson (SUSY)} \\ \bar{\partial} \partial \quad \hookrightarrow \quad \text{generates Boson for every Fermion (SUSY)} \end{array} \right. \end{array} \right\}$$

## Conclusions

The origin of particles has been shown to represent the scenario for the creation of supersymmetry through the chiral operators  $L$  and  $\eta \times \eta$  matrix that in essence creates the generator  $\nabla_{\pm} \equiv \nabla_{(p,q)}$  for the hypercomplex potential  $\partial^{(6)} \rho \rho \rho^{+1} \rho^{-1} \rho^{+0} \rho^{-0}$  to represent  $(p, q)$  norms for the particle creation through the permutation cycle of the vibrations  $\psi$  via the differential operator  $\epsilon$  in four-possible ways:  $\partial \partial, \bar{\partial} \bar{\partial}, \partial \bar{\partial}, \bar{\partial} \partial$ .

## References

- <sup>a, b</sup> Bhattacharjee, D. (2022k). *Generalization of Quartic and Quintic Calabi – Yau Manifolds Fibered by Polarized K3 Surfaces*. Research Square (Research Square). <https://doi.org/10.21203/rs.3.rs-1965255/v1>
- <sup>a</sup> Katz, S. (2007). *Enumerative geometry and string theory*. Choice Reviews Online, 44(10), 44-5694. <https://doi.org/10.5860/choice.44-5694>
- <sup>a</sup> Albeverio, S., Jost, J., Paycha, S., & Scarlatti, S. (1997). *A Mathematical Introduction to String Theory*. A Mathematical Introduction to String Theory, 143.
- <sup>a</sup> Bhattacharjee, D. (2022d). *Imaginary Cycles of Permutations for Genus g=3 in Complex Geometries*. EasyChair Preprint No. 8200. <https://easychair.org/publications/preprint/CWFx>
- <sup>a</sup> Bhattacharjee, D. (2022, June 28). *An outlined tour of geometry and topology as perceived through physics and*

- mathematics emphasizing geometrization, elliptization, uniformization, and projectivization for Thurston's 8-geometries covering Riemann over Teichmüller spaces. *TechRxiv*. <https://doi.org/10.36227/techrxiv.20134382.v1>
6. <sup>^</sup>Penrose, R. (2005). *The road to reality: a complete guide to the laws of the universe*. *Choice Reviews Online*, 43(01), 43-0377. <https://doi.org/10.5860/choice.43-0377>
  7. <sup>^</sup>Greene, B. (2000, February 3). *The Elegant Universe*. In *Superstrings, Hidden Dimensions, and the Quest for the Ultimate Theory*. Vintage.
  8. <sup>a, b</sup>Bhattacharjee, D. (2022, May 10). *Establishing equivalence among hypercomplex structures via Kodaira embedding theorem for non-singular quintic 3-fold having positively closed (1,1)-form Kähler potential  $2-1\partial\bar{\partial}^*p$* . *Research Square*. <https://doi.org/10.21203/rs.3.rs-1635957/v1>
  9. <sup>^</sup>Gross, M. (2002, November 27). *Calabi-Yau Manifolds and Related Geometries*. In *Lectures at a Summer School in Nordfjordeid, Norway, June 2001*. Springer. <https://doi.org/10.1007/b8363310.1007/978-3-642-19004-9>
  10. <sup>^</sup>Bhattacharjee, D., Samal, P., Bose, P. N., Behera, A. K., & Das, S. (2023, April 5). *Suspension  $\eta$  for  $\beta$  bundles in  $\pm 1$  geodesics in  $g \geq 1$  genus creations for loops for a Topological String Theory Formalism*. *TechRxiv*. <https://doi.org/10.36227/techrxiv.22339732.v1>
  11. <sup>^</sup>Collier, P. (2013, January 1). *A Most Incomprehensible Thing*. In *Notes Towards a Very Gentle Introduction to the Mathematics of Relativity*.
  12. <sup>^</sup>Kaku, M. (1994, March 24). *Hyperspace*. In *A Scientific Odyssey Through Parallel Universes, Time Warps, and the Tenth Dimension*. <https://doi.org/10.1604/9780195085143>
  13. <sup>a, b, c</sup>Bhattacharjee, D. (2022, July 1). *Suspension of structures in two-dimensional topologies with or without the presence of  $\geq 1$  genus deformations for canonical  $2\eta$  stabilizer points*. *Research Square*. <https://doi.org/10.21203/rs.3.rs-1798323/v1>
  14. <sup>a, b</sup>Bhattacharjee, D. (2022, July 15). *M-Theory and F-Theory over Theoretical Analysis on Cosmic Strings and Calabi-Yau Manifolds Subject to Conifold Singularity with Randall-Sundrum Model*. *Asian Journal of Research and Reviews in Physics*, 25-40. <https://doi.org/10.9734/ajr2p/2022/v6i230181>