

[Open Peer Review on Qeios](#)

# Effect of Animal-Assisted Activities on Symptoms and Emotions of Children with Neoplastic Disease: A Systematic Review with Meta-Analysis

Luca Giuseppe Re<sup>1</sup>, Silvia Porcarelli<sup>2</sup>, Camilla Ripari<sup>2</sup>, Sara Marotta<sup>2</sup>

<sup>1</sup> University of Milan

<sup>2</sup> Ospedale Niguarda Ca' Granda

**Funding:** No specific funding was received for this work.

**Potential competing interests:** No potential competing interests to declare.

## Abstract

**Background:** Animal-assisted activities (AAAs) have long been present in pediatric oncology as a nonpharmacological intervention aimed at helping children cope with symptoms and negative emotions during hospitalization and antineoplastic treatment. Among the systematic reviews in the literature, there is a lack of one with meta-analysis that includes only RCTs centered on the effect of the intervention on symptoms and emotions in children with neoplastic disease.

**Objective:** To synthesize the effect of AAAs on symptoms and emotions of children with neoplastic disease.

**Methods:** Studies were searched from biomedical databases, trial registries, and web resources. The risk of bias of included studies was assessed with RoB 2, and the overall effect size of the intervention was calculated by creating random-effects meta-analyses graphically represented by forest plots. The summary of findings was illustrated with a table in accordance with the GRADE method.

**Results:** Three parallel-group randomized controlled trials with low risk of bias were included (N = 151, mean age: 8.5-11.2 years). AAAs are promising for anxiety reduction (N = 134; SMD = -0.07 [95% CI: -0.40, 0.27],  $p > 0.05$ ) and quality of life improvement (N = 84; SMD = -0.11 [95% CI: -0.53, 0.31],  $p > 0.05$ ) in children and for anxiety reduction (N = 154; SMD = -0.50 [95% CI: -1.52, 0.52],  $p > 0.05$ ) in parents/caregivers. The certainty/quality of evidence is low to very low.

**Conclusions:** The effect of animal-assisted activities to reduce anxiety and improve the quality of life of children with neoplastic disease and to reduce the anxiety of parents/caregivers is promising. However, more studies need to be conducted on the topic that overcome the important methodological limitations manifested by current research in order to collect more data in favor of implementing the intervention in pediatric oncology.

**Keywords:** animal-assisted activities, children, emotions, meta-analysis, neoplasia, symptoms.

## Introduction

Although cancer remains a leading cause of death in the pediatric population<sup>[1]</sup>, its prognosis has improved over the past decades<sup>[2]</sup>. The counterpart of this positive outcome is the use of complicated, intensive, and prolonged treatment protocols involving multiple hospitalizations and/or outpatient care over a period of time generally exceeding one year<sup>[3][4]</sup>. During this time, the child is exposed to invasive procedures that cause him or her physical and emotional suffering<sup>[5]</sup>. Pain is the most prevalent symptom, followed by fatigue, anxiety, fear, and depression<sup>[6][7][8][9][10]</sup>. Added to this is the sense of distress, which assails the child in seeing his or her body image gradually change<sup>[6]</sup>. The effects are manifested in terms of psychological disorders, social withdrawal and maladaptive behaviors, with a negative effect on the treatment process, the overall clinical condition<sup>[6][11][12][13]</sup>, self-image and the ways of interacting with family members and peers<sup>[6][7]</sup>.

The exploration and use of alternative therapies to complement traditional medical care, alleviate symptoms, manage and control negative emotions, promote positive ones, and improve the child's perceived quality of life during hospitalization and antineoplastic treatment must be a primary goal of nursing care in pediatric oncology<sup>[14][15]</sup>. These therapies include pet therapy, which has long been present in a wide variety of settings, including the oncology setting, benefiting large segments of the population, including the pediatric population<sup>[16]</sup>.

The first documented study of the scientific use of animals for therapeutic purposes dates back to 1859 by Florence Nightingale, who discovered that the presence of small pets was an important support to the healing process of the acutely ill and an excellent companion for the chronically ill<sup>[17]</sup>. In 1964, child psychologist Boris Levinson was the first to coin the term "pet therapy"; he observed that his own dog spontaneously interacted with a child with autism, activating responses and reactions in him that had not been achieved with traditional treatment<sup>[18]</sup>. In 1977, veterinarian Leo K. Bustad and psychiatrist Michael J. McCulloch formed the Delta Foundation in Portland, Oregon, USA, which later became the Delta Society in 1981, with the goal of improving people's health and well-being through interactions with animals. In the late 1980s and early 1990s, among the initiatives promoted by the Delta Society was the Pet Partners® program, the first comprehensive and standardized training course on animal-assisted activities and therapy for volunteers and health care workers. In 2012, the Delta Society formally changed its name to Pet Partners, currently one of the largest organizations in the field of hospital-based pet therapy in the United States. In Italy, pet therapy has been recognized as an official treatment since 2003<sup>[19]</sup>. Thanks to the partnership between the Istituto Superiore di Sanità, Centro di Referenza Nazionale per gli Interventi Assistiti da Animali (National Reference Center for Animal-Assisted Interventions) and the Ministry of Health, the National Guidelines for Animal-Assisted Interventions were issued in 2015 in order to protect people's health and the welfare of the animals used, define operational standards for the correct and uniform application of the interventions, provide guidance on the tasks and responsibilities of the professional figures and operators involved (e.g., nurses, veterinarians, physicians, psychologists, educators), and identify training methods<sup>[20]</sup>.

The term "pet therapy" is widespread but inaccurate; more appropriately, one should speak generally of "animal-assisted intervention" (AAI), but specifically of "animal-assisted activity" (AAA) or "animal-assisted therapy" (AAT) [8][21][22]. Both of these interventions involve interaction between a pet and a human being, take place mostly in outpatient clinics, hospitals, or residential facilities, and are designed to complement and supplement traditional therapeutic modalities [6][23]. They are sometimes used interchangeably in the literature but possess distinctive characteristics [15].

AAT can be an essential part of individualized treatment aimed at people with physical, social, emotional, or cognitive difficulties, or suffering from conditions such as autism or depression; it requires stated goals for each session and is aimed at a specific clinical outcome [6][15][24][25][26]. The animal is carefully selected and undergoes a formal training period; the handler (who often coincides with the trainer) must undergo specific training based on the treatment to be delivered [25][27][28][29]. Progress must be documented, and sessions are multiple and scheduled to last a set period of time depending on the person's needs and resources [6][15][25].

AAA aims to manage symptomatology and negative emotions, offer comfort, and improve people's quality of life with brief (usually 15-30 minutes) informal pet visits accompanied by a handler (the owner or trainer); the manner of interaction with the animal is at the discretion of the person in agreement with the handler, and there are no specific therapeutic goals. No formal training of the animal or special training of the handler is usually required, and the same animal is not always paired with the same person [6][15][25]. Progress is not documented, and there are few sessions (sometimes only one) [6][15][25]. The theoretical frameworks underlying the effects observed after AAA in oncology include the biophilia hypothesis, social support theory, general human-animal bonding theory, cognitive stress activation theory, the object-self hypothesis, and the science of unitary humans (Holder 2020b). Specifically, the biophilia hypothesis holds that humans have a natural attraction to other living things [30][31][32][33] and this triggers the initial impulse to interact with the animal. This hypothesis would be able to justify the benefit observed even during a single session of short duration, a typical feature of AAAs [34].

According to Johnson et al. [35], specific populations that may benefit most from AAAs include individuals with malignancy, especially children. For them, the positive effects occur at several levels: (a) they reduce anxiety and pain, promote positive emotions, and improve mood [6][13][21][36][37][38][39]; (b) they increase interaction skills by acting as a "social lubricant" [8][15][21][38][40]; (c) they help normalize the experience of hospitalization [38][41], allow for the acquisition of self-esteem and confidence, and nurture a sense of responsibility, enabling the child to orient to the future by overcoming the sick condition [6]; (d) they improve cooperation in treatment and encourage active participation [12][42]; (e) they reduce blood pressure and heart rate [43]; (f) they increase plasma concentrations of endorphins, which help relieve pain, reduce stress, and generate a feeling of well-being; and (g) they decrease plasma concentrations of cortisol, which negatively affects the degree of stress.

There are three systematic reviews in the literature that have addressed the effectiveness of AAIs on children. A systematic review with meta-analysis [44] investigated the efficacy of AAIs on some clinical outcomes in children and adults; however, participants also had pathologic conditions that were not neoplastic in nature, and there was no subgroup analysis by age group. A systematic review [45] summarized the available evidence on the effectiveness of AAIs in

pediatric oncology; however, the authors included both observational and experimental studies and did not perform a quantitative synthesis. Finally, a recent systematic review with meta-analysis [46] summarized the effectiveness of AAT on pain management in hospitalized children; however, again, participants had been hospitalized for pathological conditions including non-neoplastic conditions, and the authors included both randomized and nonrandomized controlled experimental studies. Thus, a systematic review with meta-analysis of randomized controlled clinical trials summarizing the effect of AAAs on children with neoplastic pathology is lacking. It is considered important to undertake this study because the results could fill a significant gap in the knowledge of the effectiveness of this intervention in pediatric oncology and prove useful in obtaining new information to help optimize and personalize its application.

## Objective

The study aims to summarize the effect of AAAs on the symptoms and emotions of children with neoplastic disease.

## Methods

To achieve the objective, a systematic review with meta-analysis was conducted in accordance with PRISMA guidelines [47]. The review protocol was registered with the International Prospective Register of Systematic Reviews (PROSPERO) (ID: CRD42024540556).

## Eligibility Criteria

The research question was formulated in accordance with the PICOS framework [48] (Table 1); the inclusion criteria were as follows: (1) participants: subjects with established neoplastic disease, aged 0-18 years, to be treated with oncology care during inpatient, outpatient, or day hospital; (2) intervention: animal-assisted activities; (3) control: standard of care; (4) outcomes: (a) primary - child's pain or anxiety, measured by any instrument at the end of the first visit or at the end of the study period; (b) secondary - child's fear, distress, fatigue, depression, stress, mood, discomfort, quality of life, measured by any instrument at the end of the first visit or at the end of the study period; (c) parent/caregivers' anxiety, measured by any instrument at the end of the first visit or at the end of the study period; (5) study design: parallel-group randomized controlled trials.

Studies were excluded: (1) with mixed populations (adults and children) in which the effect of the intervention on children could not be assessed; (2) in which the intervention was delivered to a group of children and not to the individual subject; (3) that compared the intervention in question with other interventions; (4) in which at least one of the outcomes, primary or secondary, of interest was not present; (5) conducted in the community.

**Table 1.** PICOS framework.

P	Patient	Children with neoplastic disease
I	Intervention	Animal-assisted activities
C	Comparison	Standard of care
O	Outcome	Symptoms, emotions
S	Study design	Parallel-group randomized controlled trials
<i>Research Question</i> - In children with neoplastic disease, are animal-assisted activities more effective on symptoms and emotions than standard of care?		

## Information sources and search strategy

To answer the research question, the biomedical databases Cochrane Library, MEDLINE via PubMed, EMBASE via Elsevier, CINAHL via EBSCOhost, PsycINFO via Ovid, Web of Science via Clarivate, Scopus via ELSEVIER, AMED via EBSCOhost, sciELO, LILACS, CNKI, J-GLOBAL, J-STAGE were queried. In addition, web resources BASE, TRIP Medical Database, IBSS via ProQuest, Social Science Premium Collection via Proquest, Dissertations & Theses Global via Proquest, Google Scholar, and clinical trial registries ICTRP, ClinicalTrials.gov, EU Clinical Trials Register, ISRCTN were consulted. "Animal-assisted therapy," "neoplasm," and related synonyms were used as keywords to search the documents. The keywords were connected to each other with Boolean AND/OR operators; the search string was adapted to the characteristics peculiar to the queried information sources. The search strategy implemented on MEDLINE is shown in Table 2 as an example. References of eligible studies and available reviews were searched to retrieve other relevant studies. The search was restricted to records pertaining to the pediatric population and containing keywords or synonyms in the title and/or abstract. No language or publication date limits were imposed.

**Table 2.** Search strategy.

Animal Assisted Therapy	Neoplasm	Preschool Child
OR	OR	OR
Animal Assisted Therapies	Neoplasms	Preschool Children
OR	OR	OR
Animal-Assisted Therapy	Tumor	Pre-school Child
OR	OR	OR
Animal-Assisted Therapies	Tumors	Pre-school Children
OR	OR	OR
Animal Facilitated Therapy	Tumour	Preschooler
OR	OR	OR
Animal Facilitated Therapies	Tumours	Pre-schoolers
OR	OR	OR
Pet Therapy	Neoplasia	Child
OR	OR	OR
Pet Therapies	Neoplasias	Children
OR	OR	OR

Pet Facilitated Therapy		Cancer		Adolescent
OR		OR		OR
Pet Facilitated Therapies		Cancers		Adolescents
OR		OR		OR
Pet-Assisted Therapy		Malignant Neoplasm		Adolescence
OR		OR		OR
Pet-Assisted Therapies		Malignant Neoplasms		Teen
OR	AND	OR	AND	OR
Pet Therapy Animal		Malignancy		Teens
OR		OR		OR
Pet Therapy Animals		Malignancies		Teenager
OR		OR		OR
Emotional Support Animal		Neoplastic Disease		Teenagers
OR		OR		OR
Emotional Support Animals		Neoplastic Diseases		Youth
OR		OR		OR
Comfort Animal		Neoplastic Entity		
OR		OR		
Comfort Animals		Neoplastic Mass		
OR		OR		
Emotional Support Dog		Tumoral Mass		
OR		OR		
Emotional Support Dogs		Tumoural Mass		
OR		OR		Youths
Therapy Dog		Tumorous Mass		
OR		OR		
Therapy Dogs				
OR				
Pet Therapy Dog		Tumourous Mass		
OR				
Pet Therapy Dogs				

## Study selection and data extraction

After the production of a shared search strategy, the authors independently queried the information sources by eliminating duplicates and selecting records based on relevance of title and/or abstract or, in doubtful cases, after full-text analysis. The record screening process was managed with a Microsoft Excel version 2016 spreadsheet. Any disagreements were resolved by comparison and discussion. From each included study, using a standardized and shared template, the authors independently extracted the following information: first author and year of publication; country; hospital or research facility and setting; sample characteristics; inclusion and exclusion criteria; intervention and control

characteristics; outcome and its measurement tools; and any notes.

### Risk of bias

The authors independently assessed the risk of bias of included studies with RoB 2<sup>[49]</sup>. Any disagreement was resolved by comparison and discussion.

### Data analysis and synthesis

The authors independently extracted the data and resolved any differences of opinion through comparison and discussion. The variables of interest for each outcome were sample size, mean, and standard deviation. In the presence of median, range, or interquartile range, conversion equations were used <sup>[50][51][52]</sup>. The overall effect size of the intervention was calculated with the standardized mean difference (SMD) and Cohen's  $d$  <sup>[53]</sup>. Effect size was considered small, moderate, or large for thresholds of  $d$  of 0.2, 0.5, 0.8, respectively <sup>[53]</sup>. In the presence of at least two studies per outcome, a meta-analysis was performed by applying a random-effects model and generating the corresponding forest plot. A 95% confidence interval was considered as the deviation from the point estimate for each individual study and from the overall estimated value for the aggregated studies. The presence of statistical heterogeneity ( $p < 0.05$ ) was highlighted with Cochran's Q-test <sup>[54]</sup> and quantified with Higgins'  $I^2$  index <sup>[55]</sup>. Values of  $I^2 \leq 30\%$ ,  $30\% < I^2 \leq 60\%$ ,  $60\% < I^2 \leq 90\%$ , or  $I^2 > 90\%$  were assigned a low, moderate, high, or very high degree of statistical heterogeneity, respectively <sup>[56]</sup>. Statistical processing was performed with ProMeta© version 3.0 software.

### Publication bias

Publication bias was assessed by inspecting the funnel plot<sup>[57]</sup> and applying the trim and fill method<sup>[58]</sup> in the presence of at least ten studies. Objective assessment of publication bias was performed with Egger's test<sup>[59]</sup> and Begg and Mazumdar's test<sup>[60]</sup>.

### Sensitivity analysis

In the presence of studies at high risk of bias, sensitivity analysis was performed by regenerating the meta-analysis after their exclusion.

### Additional analysis

Subgroup analyses were planned to assess the effect of animal-assisted activities according to participants' gender and age.

### Summary of findings

The authors independently performed the overall assessment of certainty/quality of evidence using the GRADE

method [61] applied to the meta-analysis results. Disagreements that emerged were resolved by comparison and discussion.

## Results

### Selection of studies

Searching for useful records took place on April 18, 2024. The PRISMA flowchart [47] in Figure 1 illustrates the record selection process. A total of 178 records were identified. Net of duplicates and irrelevant records after reading titles and abstracts, 30 studies, for as many records, were analyzed in full text and evaluated for eligibility. Twenty-seven were excluded because they did not meet the inclusion criteria, while three were included in the systematic review and quantitative synthesis [62][63][64]. No other studies of interest were found after consulting the references of eligible studies and relevant reviews.




Figure 1. PRISMA Flow Diagram.

### Characteristics of the studies

The studies cover a 6-year time frame from 2018 [64] to 2024 [63] (Table 3) and were conducted in the United States. One



of these studies is multicenter [64]. All studies received funding; the last author of one study [62] reported a possible conflict of interest. There were a total of 151 children recruited and 187 parents/caregivers. The age range was 3-17 years, the mean age 8.5-11.2 years (63.2-69.2% Caucasian), and the percentage of males 47.4-57.7%. The prevalent diagnoses were acute lymphoblastic leukemia (51.9%) [64], leukemia or lymphoma (61.5%) [62], and advanced solid tumor (52.6%) [63]. The percentage of participants who owned a pet was 57.9-84.6%. The most frequent exclusion criteria were the presence of cognitive impairment in the child and/or parent/caregiver and fear or allergy to animals. Animal-assisted activities all included the presence of a trained dog and its handler. The breed of dogs was specified in only one study [64], and these were predominantly Labradors and Golden retrievers. The intervention was weekly for all studies, was individual, and could take place in the outpatient clinic, in the inpatient room, or in dedicated spaces within the hospital. The duration of the intervention was 10-20 minutes; activities were at the discretion of the child and/or handler but generally involved the child petting, cuddling, or playing with the dog. Studies have compared animal-assisted activities with standard of care; of the latter, no study has provided a description. The outcomes assessed after the intervention were as follows: (a) child anxiety [62][63][64], measured by PedsQL VAS - anxiety item [65] or STAI-CH [66]; (b) child quality of life [62][64], measured by PedQL VAS [65] or PedsQL [67]; (c) caregiver/parent anxiety [62][63][64], measured by STAI Short Form [68] or STAI [69]. In addition, one study [62] measured the levels of clinically important pathogens on each child's hands, and one study [64] measured the child's heart rate and blood pressure. The outcomes were assessed at baseline, after the first view [62], at the first and fourth views [63] or at the end of the intervention [64], in this case as arithmetic averages of the values measured in the previous sessions. The duration of the studies was 1-4 months. In two studies [62][63] the COVID-19 pandemic conditioned the recruitment of participants.

**Table 3.** Main features of included studies.

Study (year)	Country	Setting	Sample	Inclusion criteria	Exclusion criteria	Intervention	Control	Outcome	Notes
Chubak (2023)	United States	Seattle Children's Hospital (Seattle, WA)	N = 26 (IG = 12, CG = 14), mean age 11.2 years, range 5-12 years, males 57.7%, Caucasian race 69.2%, diagnosed with leukemia or lymphoma 61.5%, pet owner 84.6%  Caregivers or parents: N = 19	English-speaking subjects aged 5 to 17 years regardless of the type of neoplasm and gender, race, or ethnicity	Allergy to or fear of dogs, subjects undergoing bone marrow transplantation or isolation precautions, with skin on hands not intact, without an English-speaking parent or legal guardian	Weekly inpatient room visits  One visit per child lasting 20 minutes  Hand hygiene before and after visit  Activities at the discretion of the child and handler	Standard of care	Quality of life of the child Anxiety of caregivers/parents  Detection of clinically relevant pathogens on children's hands after the first AAA	Data collection performed before randomization, before and after the first AAA, approximately 9 days after the first AAA, at discharge, 2-3 days after discharge (follow-up 1), 9 weeks after discharge (follow-up 2)  Outcome assessed after the first AAA; subsequent assessments not illustrated due to poor

Author (Year)	Country	Study Location	Participants (N, IG, CG, Age, Sex, Race, Diagnosis, Pet Owner)	Language	Eligibility Criteria	Intervention	Control	Outcomes	Notes
Mahoney (2024)	United States	Vanderbilt Children's Hospital (Nashville, TN)	N = 19 (IG = 9, CG = 10), mean age 9 years, range 3-17 years, males 47.4%, Caucasian race 63.2%, diagnosed with solid tumor 52.6%, pet owner 57.9%	English-speaking	Subjects aged 3-17 years, diagnosed with advanced cancer (relapsed or refractory)	Children or parents with cognitive impairment or fear of or allergy to dogs	Standard of care	Child anxiety Anxiety of caregivers/parents	Due to poor adherence to follow-ups. At the first AAA, three children did not complete the questionnaire. Data collection performed before randomization and weekly after each AAA (4 visits). Outcomes assessed after the first and last AAAs
McCullough (2018)	United States	Vanderbilt Children's Hospital (Nashville, TN); Randall Children's Hospital at Legacy Emanuel (Portland, OR); UC Davis Children's Hospital (Sacramento, CA); St. Joseph's Children's Hospital (Tampa, FL); UMass Memorial Medical Center - Children's Medical Center (Worcester, MA)	N = 106 (IG = 60, CG = 46), mean age 8.5 years, range 3-17 years, males 53.8%, Caucasian race 67.9%, diagnosed with acute lymphoblastic leukemia 51.9%, pet owner 67%	English or Spanish	Subjects aged 3 to 17 years, diagnosed with a type of cancer that would have required at least monthly outpatient treatment (e.g., leukemia, lymphoma, solid tumor, brain tumor), English or Spanish as the primary language of the child and his or her parents	Children or parents with significant cognitive impairment, allergy or fear of dogs	Standard of care	Anxiety of the child Anxiety of caregivers/parents Child's quality of life Blood pressure and heart rate of the child	Data collection conducted before randomization and weekly after each visit. Outcomes assessed before randomization and at the end of the intervention. Seven children discontinued the study due to changes in treatment plan and/or noncompletion of questionnaires

CG = Control Group; IG = Intervention Group.

## Risk of bias

In a study [64], the risk of bias is low; in the other two, the risk of bias is of some concern, since there is no information on assignment sequence concealment (Figure 2 and Figure 3).




Figure 2. Risk of bias (summary plot).




Figure 3. Risk of bias (traffic light).

## Primary outcomes

### Child pain

No studies have evaluated this outcome.

### Child's anxiety

The child's anxiety was assessed by all studies. The values considered were those measured after the first visit [62][63] or the mean value calculated at the end of the study [64]. The analysis of the effect of AAAs on child anxiety compared with standard of care included 134 participants. The SMD (95% CI) was -0.07 [-0.40,0.27], I2 = 0.00 in favor of the

intervention in a statistically nonsignificant manner (Table 4).

Study	AAA			Standard of care			Std. Mean Difference IV, Random, 95% CI	Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Chubak 2023	5.1	8.7	11	4.2	5.5	12		17.98%	0.12 [-0.67, 0.91]
Mahoney 2024	29.9	4.5	9	31.3	6.9	10		15.03%	-0.23 [-1.09, 0.64]
McCullough 2018	28.8	6.7	52	29.3	5.4	40		66.99%	-0.08 [-0.49, 0.33]
<b>Total (95% CI)</b>								<b>100.00%</b>	<b>-0.07 [-0.40, 0.27]</b>

Q = 0.35 (p = 0.838); I<sup>2</sup> = 0.00; T<sup>2</sup> = 0.00; T = 0.00

Table 4. Child's anxiety: AAAs vs standard of care.

### Secondary outcomes

No studies assessed child fear, distress, fatigue, depression, stress, discomfort, or mood.

#### Child's quality of life

The child's quality of life was assessed in two studies<sup>[62][64]</sup>. The values considered were those measured after the first visit<sup>[62]</sup> or the average value calculated at the end of the study<sup>[64]</sup>. The analysis of the effect of AAA on the child's quality of life compared to standard of care included 84 participants. The SMD (95% CI) is equal to -0.11 ([-0.53,0.31], I<sup>2</sup> = 98.59%) in favor of the intervention in a statistically insignificant way (Table 5).

Study	AAA			Standard of care			Std. Mean Difference IV, Random, 95% CI	Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Chubak 2023	11	12.5	11	13.6	12.8	12		17.98%	-0.20 [-0.99, 0.59]
McCullough 2018	69	15.6	34	70.4	22	27		66.99%	-0.08 [-0.58, 0.42]
<b>Total (95% CI)</b>									<b>100.00%</b>

Q = 0.06 (p = 0.803); I<sup>2</sup> = 0.00; T<sup>2</sup> = 0.00; T = 0.00

Table 5. Child's quality of life: AAAs vs standard of care.

### Parental/caregivers anxiety

Parental/caregiver anxiety was assessed in all studies. The values considered were those measured after the first visit<sup>[62][63]</sup> or the average value calculated at the end of the study<sup>[64]</sup>. The analysis of the effect of AAA on parents/caregivers' anxiety compared to standard of care included 154 participants. The SMD (95% CI) is equal to -0.50 ([-1.52,0.52], I<sup>2</sup> = 83.48) in favor of the intervention in a statistically insignificant way (Table 6).

Study	AAA			Standard of care			Std. Mean Difference IV, Random, 95% CI	Weight	Std. Mean Difference IV, Random, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Chubak 2023	38.9	11.4	9	46.3	8.9	9		33.88%	-0.69 [-1.60, 0.22]
Mahoney 2024	28.8	9.1	10	47	16.5	11		34.41%	-1.26 [-2.20, -0.38]
McCullough 2018	66.4	11.4	68	62.9	13.7	47		31.71%	0.28 [-0.09, 0.65]
Total (95% CI)			87			67		100.00%	-0.50 [-1.52, 0.52]

$Q = 12.11$  ( $p = 0.002$ );  $I^2 = 83.48$ ;  $T^2 = 0.67$ ;  $T = 0.82$

**Table 6.** Parental/caregivers anxiety: AAAs vs standard of care.

## Additional analysis

### Gender

No study has measured the effect of the intervention according to the gender of participants.

### Age

No study has measured the effect of the intervention according to the age group of participants.

## Sensitivity analysis

Sensitivity analysis was not performed as none of the included studies is at high risk of bias.

## Publication bias

The studies included are less than ten; therefore, the funnel plot was not created, and the trim and fill method was not applied for the graphic evaluation of risk of bias. However, the objective assessment suggests that risk of publication bias, although possible, seems unlikely; in fact, the Egger test and the Begg and Mazumdar test are not statistically significant ( $p = 0.942$  and  $p = 0.602$ , respectively).

## Summary of findings

With the GRADE method, findings relating to the effect of AAAs on children with neoplastic pathology were summarized. Despite the trend in favor of intervention, the certainty/quality of evidence on children's anxiety and quality of life is low, and on parents'/caregivers' anxiety is very low (Table 7).

**Table 7.** Effect of AAAs on symptoms and emotions of children with neoplastic disease: summary of findings.

Summary of findings. Animal-assisted activities (AAAs) for symptoms and emotions in children with neoplastic disease.

AAAs compared to standard of care for symptoms and emotions in cancer children

Patient or population: children with cancer

Setting: hospital, clinic, day hospital

Intervention: AAAs

Comparison: standard of care

Outcome	Anticipated absolute effects* (95% CI)		N° of participants (studies)	Certainty/quality of the evidence (GRADE)	Comments**
	Risk with standard care	Risk with AAAs			
Children anxiety	-	The mean level of anxiety with AAAs was 0.07 standard deviation lower (0.40 lower to 0.27 higher).	134 (3 RCTs)	⊕⊕⊕⊕ Low <sup>a</sup>	There is no evidence of an effect of AAAs.
Children quality of life	-	The mean level of anxiety with AAAs was 0.11 standard deviation lower (0.53 lower to 0.31 higher).	84 (2 RCTs)	⊕⊕⊕⊕ Low <sup>a</sup>	There is no evidence of an effect of AAAs.
Parental/caregivers anxiety	-	The mean level of anxiety with AAAs was 0.50 standard deviation lower (1.52 lower to 0.52 higher).	154 (3 RCTs)	⊕⊕⊕⊕ Very low <sup>a,b</sup>	There is no evidence of an effect of AAAs.

\*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

\*\*0.2 represents a small difference, 0.5 a moderate difference, and 0.8 a large difference.

AAAs: Animal-Assisted Activities; CI: confidence interval; RCT: Randomized Controlled Trial; QoL.

#### GRADE Working Group grades of evidence

High certainty - We are very confident that the true effect lies close to that of the estimate of the effect

Moderate certainty - We are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low certainty - Our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect

Very low certainty - We have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect

<sup>a</sup> Downgraded twice for imprecision: analysis based on < 100 participants per group.

<sup>b</sup> Downgraded once for inconsistency due to substantial heterogeneity (60% < I<sup>2</sup> < 90%).

## Discussion

### Main results

The study aimed to summarize the effect of animal-assisted activities (AAAs) on symptoms and emotions of children with

neoplastic disease. The included studies evaluated the benefit of the intervention on children's anxiety and quality of life and on parental/caregivers' anxiety. The results, statistically non-significant, reveal that a) the intervention is promising for reducing children's anxiety (SMD = -0.07; N = 134) and for improving their quality of life (SMD = -0.11; N = 84). The effect size is small; b) the intervention also shows a positive trend for decreasing parental/caregivers' anxiety (SMD = -0.50; N = 154). The effect size is moderate. The level of certainty/quality of evidence is low or very low, so confidence in the effect estimate is limited or very limited: it is likely that the real effect is substantially different.

## Comparison with other reviews

The results obtained from our study are consistent with those of previous systematic reviews<sup>[44][45][46]</sup>, which demonstrates the benefit of the intervention both on child anxiety<sup>[44][46]</sup> and on parental/caregivers' anxiety<sup>[45][46]</sup>. As regards the effect on children's quality of life, it is not possible to make a comparison because previous reviews did not evaluate this outcome.

## Implications for practice

In children with an average age of 8.5-11.2 years suffering from leukemia, lymphoma, or solid tumors in advanced stages, AAAs in which the animal is a dog are promising for reducing their anxiety and improving their quality of life and for decreasing the anxiety of their parents/caregivers. The results are still provisional, due to few studies available (all carried out in a single country) and the small sample size, which determines low statistical power.

Although statistical heterogeneity is zero for both anxiety and quality of life of the child, there are several sources of clinical heterogeneity: (a) the average age of the participants is between 8.5 and 11.2 years (school-age children), but the age range is very wide (3-17 years), and this may have determined a different effectiveness of the intervention depending on the child's level of neurocognitive development; (b) the children are mostly Caucasian and live in the United States, so it is not a given that the intervention will work on children of other races or who live in other countries; (c) the type of neoplasm and the degree of severity are different, and these aspects could have determined a different sensitivity and predisposition in children. Other sources of heterogeneity are the following: a) the role - active, passive, or neutral - of the handler is not known; this makes it difficult to distinguish the net effect of the intervention because any positive interactions between the handler and parents/caregivers, promoted by the social lubricant function of AAAs, could in themselves have had a role in reducing the child's anxiety; b) the standard of care was not described; (c) for all studies, the visits took place weekly, but one study<sup>[62]</sup> limited the maximum number of visits per child to four, while the other two studies, which lasted 12 weeks<sup>[63]</sup> and four months<sup>[64]</sup>, did not declare a maximum limit of visits; (d) outcome measurement tools have different intrinsic characteristics (e.g., duration of compilation, method of administration, sensitivity, and specificity). Finally, the intervention seems to work when the animal is a dog, but there is no evidence that the same benefit is obtained with other animals. The dog is the most commonly used animal for AAAs due to its ease of training, docility, obedience, and predictability of behavior<sup>[15][70][71]</sup>; furthermore, compared to other animals, it is more in tune with human moods and emotions<sup>[72]</sup>.

One of the main concerns that often affects the application of AAAs in a healthcare setting is the infectious risk and the allergic risk. Although animals carry germs and can unintentionally serve as mechanical vectors of hospital-associated pathogens, contributing to their transmission between patients, there is insufficient evidence demonstrating increased infection rates during AAAs [73]. Furthermore, with the application of adequate hospital infection control protocols, the associated risks are minimized [74]. People most likely to contract diseases from animals include infants, children under five years of age, organ transplant patients, people with HIV/AIDS, or those being treated for cancer [73]. However, for these subjects, the risk of infection can be significantly mitigated with simple preventive measures, including washing hands with soap and water or an alcohol-based antiseptic solution before and after touching the animal, carefully selecting both the patient and the animal, and ensuring that the animal is subjected to rigorous veterinary care [41][71][73][75]. Confirming this, in one of the included studies [62], there was no significant transfer of germs from the animal to the child during visits. To reduce the risk of allergic reactions to the animal's hair, it is recommended to bathe within 24 hours of the session, perform grooming just before the session, and wear clothing that blocks loose hair [76].

### Implications for research

Although the literature on the therapeutic implications of AAAs in oncology is limited, there are strong indications of their benefit. However, research focused on the effectiveness of intervention continues to not be definitive for several reasons: a) in general, there is a lack of rigor in theoretical frameworks underlying therapeutic human-animal interactions. This problem perpetuates a lack of empirical evidence based on clear hypotheses, hinders understanding of the mechanisms that drive the observed benefit, and therefore makes it difficult to optimize the intervention; b) many studies are preliminary in nature (e.g., pilot studies) and/or have methodological weaknesses that produce statistically insignificant effects even in the presence of clear clinical significance. Furthermore, when good-quality controlled experimental designs are implemented, not all previously observed positive results are validated; c) since AAAs are an unstructured intervention, it is complex to evaluate their effectiveness; d) there is a lack of data to support the effectiveness of the long-term intervention (e.g., at least one year of follow-up).

According to what has been written, greater rigor is therefore needed in the conduct of clinical research on AAAs; for example, the studies should be multicenter to reach a sample size that guarantees good statistical power, they should have the support of a more solid theoretical framework capable of understanding the mechanism underlying the observed results, and they should be capable of modifying some aspects of the intervention to personalize it and make it more effective for the child to whom it is proposed. It remains problematic to evaluate the effectiveness of AAAs, but perhaps efforts could be made to borrow some aspects of TAAs in order to make the intervention more structured without distorting its original characteristics.

### Limitations

The small number of participants did not allow subgroup analysis by age and gender; furthermore, the intervention was applied to subjects predominantly of Caucasian race and resident in the same country. These limitations taken together



suggest the need to adopt great caution both on the reliability of results and on their external validity. Finally, there is a lack of randomized controlled clinical trials focused on the effect of AAAs on symptoms or emotions such as pain, fear, distress, stress, discomfort, fatigue, depression, mood.

## Conclusions

Animal-assisted activities have a promising effect on the anxiety and quality of life of children with neoplastic disease and on the anxiety of parents/caregivers. However, the certainty/quality of evidence is low or very low, so at the current state of research, the results should be considered with great caution because a definitive judgment cannot be made on the real effect of the intervention.

## References

- <sup>^</sup>Steliarova-Foucher E, Colombet M, Ries LAG, Moreno F, Dolya A, Bray F, Hesselning P, Shin HY, Stiller CA; IICC-3 contributors. *International incidence of childhood cancer, 2001-10: a population-based registry study. Lancet Oncol* 2017;18(6):719-731. [https://doi.org/10.1016/S1470-2045\(17\)30186-9](https://doi.org/10.1016/S1470-2045(17)30186-9)
- <sup>^</sup>Siegel RL, Miller KD, Wagle NS, Jemal A. *Cancer statistics, 2023. CA Cancer J Clin* 2023;73(1):17-48. <https://doi.org/10.3322/caac.21442>
- <sup>^</sup>Pan HT, Wu LM, Wen SH. *Quality of life and its predictors among children and adolescents with cancer. Cancer Nurs* 2017;40(5):343-351. <https://doi.org/10.1097/NCC.0000000000000433>
- <sup>^</sup>Long KA, Marsland AL. *Family adjustment to childhood cancer: a systematic review. Clin Child Fam Psychol Rev* 2011;14(1):57-88. <https://doi.org/10.1007/s10567-010-0082-z>
- <sup>^</sup>Aldiss S, Hostman M, O'Leary C, Richardson A, Gibson F. *What is important to young children who have cancer while in hospital? Child Soc* 2009;23(2):85-98. <https://doi.org/10.1111/j.1099-0860.2008.00162.x>
- <sup>a, b, c, d, e, f, g, h, i, j, k</sup>Jenkins M, Ruchrdanz A, McCullough A, Casillas K, Fluke JD. *Canines and childhood cancer. Examining the effects of therapy dogs with childhood cancer patients and their families: literature review; 2012. https://www.americanhumane.org/app/uploads/2016/08/january2012clcompressed.pdf* Accessed May 24, 2024
- <sup>a, b</sup>Simon MV. *Participation and control experienced during animal-assisted activities by children hospitalised with cancer (Doctoral dissertation); 2014. https://repository.nwu.ac.za/bitstream/handle/10394/15550/Simon\_MV\_2014.pdf?sequence=1* Accessed June 3, 2024
- <sup>a, b, c</sup>Urbanski BL, Lazenby M. (2012). *Distress among hospitalized pediatric cancer patients modified by pet-therapy intervention to improve quality of life. J Pediatr Oncol Nurs* 2012;29(5):272-282. <https://doi.org/10.1177/1043454212455697>
- <sup>^</sup>Van Cleve L, Munoz CE, Savedra M, Riggs M, Bossert E, Grant M, Adlard K. 2012. *Symptoms in children with advanced cancer: Child and nurse reports. Cancer Nurs* 2012;35:115-125. <https://doi.org/10.1097/NCC.0b013e31821aedba>
- <sup>^</sup>Zisk-Rony RY, Lev J, Haviv H. *Nurses' report of in-hospital pediatric pain assessment: Examining challenges and*

- perspectives. Pain Manag Nurs* 2015;16(2):112-120. <https://doi.org/10.1016/j.pmn.2014.05.003>
11. <sup>a</sup>Duran J, Bravo L, Torres V, Craig A, Heidari J, Adlard, K, Secola R, Granados R, Jacob E. (2020). Quality of life and pain experienced by children and adolescents with cancer at home following discharge from the hospital. *J Pediatr Hematol Oncol* 2020;42(1):46–52. <https://doi.org/10.1097/MPH.0000000000001605>
  12. <sup>a, b</sup>Gagnon, J, Bouchard F, Landry M, Belles-Isles M, Fortier M, Fillion L. (2004). Implementing a hospital-based animal therapy program for children with cancer: A descriptive study. *Can Oncol Nurs J* 2004; 14(4):217-222. <https://doi.org/10.5737/1181912x144217222>
  13. <sup>a, b</sup>Silva NB, Osório FL. Impact of an animal-assisted therapy programme on physiological and psychosocial variables of paediatric oncology patients. *PLoS One*. 2018;13:e0194731. <https://doi.org/10.1371/journal.pone.0194731>
  14. <sup>a</sup>Bussotti EA, Leão ER, Chimentão DMN, Silva CPR. Assistência individualizada: posso trazer meu cachorro?. *Rev Esc Enferm USP* 2005;39:195-201. <https://www.scielo.br/j/reeusp/a/5TssW7ZMQ87wMZ9kz66mJtK/?format=pdf&lang=pt>
  15. <sup>a, b, c, d, e, f, g, h</sup>Goddard AT, Gilmer MJ. (2015). The role and impact of animals with pediatric patients. *Pediatr Nurs* 2015; 41(2): 65. PMID: 26292453
  16. <sup>a</sup>Ein N, Li L, Vickers K. The effect of pet therapy on the physiological and subjective stress response: A meta-analysis. *Stress Health* 2018;34(4): 477-489. <https://doi.org/10.1002/smi.2812>
  17. <sup>a</sup>Nightingale F. *Notes on nursing: What it is, and what it is not*. Lippincott Williams & Wilkins;1992.
  18. <sup>a</sup>Levinson BM. Pet psychotherapy: use of household pets in the treatment of behavior disorder in childhood. *Psychol Rep* 1965;17(3): 695-698. <https://doi.org/10.2466/pr0.1965.17.3.695>
  19. <sup>a</sup>Ciceroni C, Mugnai F. L'impiego degli animali con utilità sociale nella disabilità e nel supporto psicologico. *Manuale sulla disabilità: dai bisogni educativi speciali ai programmi di integrazione scolastica. -(Collana medico-psico-pedagogica)*, 243-263. 2012.
  20. <sup>a</sup>Istituto Superiore di Sanità. *Linee Guida Nazionali per gli Interventi Assistiti con gli Animali (IAA). L'attività istituzionale*. 2020. [https://www.iss.it/pet-therapy/-/asset\\_publisher/qFwujz9PsR3h/content/linee-guida-nazionali-per-gli-interventi-assistiti-con-gli-animale-iaa](https://www.iss.it/pet-therapy/-/asset_publisher/qFwujz9PsR3h/content/linee-guida-nazionali-per-gli-interventi-assistiti-con-gli-animale-iaa)- Accessed May 13, 2024
  21. <sup>a, b, c</sup>Caprilli S, Messeri A. Animal-assisted activity at A. Meyer Children's Hospital: a pilot study. *Evid Based Complement Alternat Med* 2006;3:379-383. <https://doi.org/10.1093/ecam/nel029>
  22. <sup>a</sup>Pet Partners. *Glossary*. 2024. <https://petpartners.org/publications/glossary/> Accessed June 17, 2024
  23. <sup>a</sup>Chubak J, Hawkes R. Animal-assisted activities: Results from a survey of top-ranked pediatric oncology hospitals. *J Pediatr Oncol Nurs* 2016;33(4):289-296. <https://doi.org/10.1177/1043454215614961>
  24. <sup>a</sup>Cerulli C, Minganti C, De Santis C, Tranchita E, Quaranta F, Parisi A. Therapeutic horseback riding in breast cancer survivors: a pilot study. *J Altern Complement Med*. 2014;20:623-629. <https://doi.org/10.1089/acm.2014.0061>
  25. <sup>a, b, c, d, e</sup>Holder TR, Gruen ME, Roberts DL, Somers T, Bozkurt A. (2020). A systematic literature review of animal-assisted interventions in oncology (Part I): Methods and results. *Integr Cancer Ther* 2020; 19:1534735420943278. <https://doi.org/10.1177/1534735420943278>
  26. <sup>a</sup>Natoli E. Activities and therapy mediated by animals (pet-therapy): international picture and state of the art in Italy. *Ann Ist Super Sanita`* 1997;33:267-272. PMID: 9470251

27. <sup>^</sup> Gilmer MJ, Baudino MN, Tielsch Goddard A, Vickers DC, Akard TF. *Animal-assisted therapy in pediatric palliative care. Nurs Clin North Am* 2016;51:381-395. <https://doi.org/10.1016/j.cnur.2016.05.007>
28. <sup>^</sup> Halm MA. *The healing power of the human-power connection. Am J Crit Care* 2008;17:373–376. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=fe37b30bbc24998ed46e0995a960d3a602687006> Accessed April 30, 2024
29. <sup>^</sup> Kamioka H, Okada S, Tsutani K, Park H, Okuizumi H, Handa S, Oshio T, Park S-J, Kitayuguchi J, Abe T, Honda T, Mutoh Y. *Effectiveness of animal-assisted therapy: A systematic review of randomized controlled trials. Complement Ther Med* 2014;22(2):371–390. <https://doi.org/10.1016/j.ctim.2013.12.016>
30. <sup>^</sup> Brodie SJ, Biley FC. (1999). *An exploration of the potential benefits of pet-facilitated therapy. J Clin Nurs* 1999;8(4): 329-337. <https://doi.org/10.1046/j.1365-2702.1999.00255.x>
31. <sup>^</sup> Kellert SR, Wilson EO. *The Biophilia Hypothesis. Island Press; 1993.* [https://philpapers.org/rec/KELTBH?utm\\_source=miragenews&utm\\_medium=miragenews&utm\\_campaign=news](https://philpapers.org/rec/KELTBH?utm_source=miragenews&utm_medium=miragenews&utm_campaign=news)
32. <sup>^</sup> Willens JS. *Animal-assisted therapies are becoming more common. Pain Manag Nurs* 2013;14 (4):183. <https://doi.org/110.1016/j.pmn.2013.10.001>
33. <sup>^</sup> Yin J, Arfaei N, MacNaughton P, Catalano PJ, Allen JG, Spengler JD. *Effects of biophilic interventions in office on stress reaction and cognitive function: a randomized crossover study in virtual reality. Indoor Air* 2019;29:1028-1039. <https://doi.org/10.1111/ina.12593>
34. <sup>^</sup> Holder TR, Gruen ME, Roberts DL, Somers T, Bozkurt A. (2020). *A systematic literature review of animal-assisted interventions in oncology (Part II): Theoretical mechanisms and frameworks. Integrat Cancer Ther* 2020; 19:1534735420943269. <https://doi.org/10.1177/1534735420943269>
35. <sup>^</sup> Johnson RA, Meadows RL, Haubner JS, Sevedge K. *Animal-assisted activity among patients with cancer: Effects on mood, fatigue, self-perceived health, and sense of coherence. Oncol Nurs Forum* 2008;35(2):225-232. <https://doi.org/10.1188/08.ONF.225-232>
36. <sup>^</sup> Baek SM, Lee Y, Sohng KY. *The psychological and behavioural effects of an animal-assisted therapy programme in Korean older adults with dementia. Psychogeriatrics* 2020;20(5):645–653. <https://doi.org/10.1111/psyg.12554>
37. <sup>^</sup> Kaminski M, Pellino T, Wish J. *Play and pets: The physical and emotional impact of child-life and pet therapy on hospitalized children. Child Health Care* 2002;31:321-335 [https://web.archive.org/web/20170812095955id\\_/http://www.kenrodogtraining.com/upload/play.pdf?prefix=www.m](https://web.archive.org/web/20170812095955id_/http://www.kenrodogtraining.com/upload/play.pdf?prefix=www.m) Accessed Jun 21, 2024
38. <sup>a, b, c</sup> Sobo EJ, Eng B, Kassity-Krich N. *Canine visitation (pet) therapy: Pilot data on decreases in child pain perception. J Holist Nurs* 2006;24:51-57. <https://doi.org/10.1177/0898010105280112>
39. <sup>^</sup> Wu AS, Niedra R, Pendergast L, McCrindle BW. *Acceptability and impact of pet visitation on a pediatric cardiology inpatient unit. J Pediatr Nurs* 2002;17:354-362. <https://doi.org/10.1053/jpdn.2002.127173>
40. <sup>^</sup> Stewart LA, Dispensa F, Parker L, Chang CY, Cunnien T. *A pilot study assessing the effectiveness of an animal-assisted outreach program. J Creativ Mental Health* 2014;9(3):332–345. <https://doi.org/10.1080/15401383.2014.892862>
41. <sup>a, b</sup> Brodie S, Biley FC, Shewring M. (2002). *An exploration of the potential risks associated with using pet therapy in*

- healthcare settings. *J Clin Nurs* 2002;11:444-456. <https://doi.org/10.1046/j.1365-2702.2002.00628.x>
42. <sup>^</sup>Bouchard F, Landry M, Belles-Isles M, Gagnon J. A magical dream: A pilot project in animal-assisted therapy in pediatric oncology. *Can Oncol Nurs J* 2004;14(1):14-17. <https://doi.org/10.5737/1181912x1411417>
43. <sup>^</sup>Beetz AM. Theories and possible processes of action in animal assisted interventions. *Applied Develop Sci* 2017;21(2): 139–149. <https://doi.org/10.1080/10888691.2016.1262263>
44. <sup>a, b, c</sup>Waite TC, Hamilton L, O'Brien W. A meta-analysis of animal assisted interventions targeting pain, anxiety and distress in medical settings. *Complement Ther Clin Pract* 2018;33:49-55. <https://doi.org/10.1016/j.ctcp.2018.07.006>
45. <sup>a, b, c</sup>Cotoc C, An R, Klonoff-Cohen H. Pediatric oncology and animal-assisted interventions: A systematic review. *Holist Nurs Pract* 2019;33(2):101-110. <https://doi.org/10.1097/HNP.0000000000000313>
46. <sup>a, b, c, d</sup>Zhang Y, Yan F, Li S, Wang Y, Ma Y. Effectiveness of animal-assisted therapy on pain in children: A systematic review and meta-analysis. *Int J Nurs Sci* 2021; 8(1), 30-37. <https://doi.org/10.1016/j.ijnss.2020.12.009>
47. <sup>a, b</sup>Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hróbjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;88:105906. <https://doi.org/10.1136/bmj.n71>
48. <sup>^</sup>Amir-Behghadami M, Janati A. Population, Intervention, Comparison, Outcomes and Study (PICOS) design as a framework to formulate eligibility criteria in systematic reviews. *Emerg Med J* 2020;37(6):387. <https://doi.org/10.1136/emered-2020-209567>
49. <sup>^</sup>Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng H-Y, Corbett MS, Eldridge SM, Emberson JR, Hernán MA, Hopewell S, Hróbjartsson A, Junqueira DR, Jüni P, Kirkham JJ, Lasserson T, Li T, McAleenan A, Reeves BC, Shepperd S, Shrier I, Stewart LA, Tilling K, White IR, Whiting PF, Higgins JPT. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366. <https://doi.org/10.1136/bmj.l4898>
50. <sup>^</sup>Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol* 2005;5:13. <https://doi.org/10.1186/1471-2288-5-13>
51. <sup>^</sup>Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res* 2018;27(6):1785-1805. <https://doi.org/10.1177/0962280216669183>
52. <sup>^</sup>Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014;14:135. <https://doi.org/10.1186/1471-2288-14-135>
53. <sup>a, b</sup>Cohen J. Statistical power analysis. *Curr Direct Psychol Sci* 1992;1(3):98-101. <https://doi.org/10.1111/1467-8721.ep10768783>
54. <sup>^</sup>Cochran WG. The comparison of percentages in matched samples. *Biometrika* 1950;37(3/4):256-266. PMID: 14801052
55. <sup>^</sup>Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327(7414):557-560. <https://doi.org/10.1136/bmj.327.7414.557>
56. <sup>^</sup>Calderón MA, Boyle RJ, Penagos M, Sheikh A. Immunotherapy: the meta-analyses. What have we learned? *Immunol Allergy Clin North Am* 2011;31(2):159-173. <https://doi.org/10.1016/j.iac.2011.02.002>
57. <sup>^</sup>Sterne JA, Egger M. Funnel plots for detecting bias in metaanalysis: guidelines on choice of axis. *J Clin Epidemiol*

- 2001;54(10):1046-1055. [https://doi.org/10.1016/s0895-4356\(01\)00377-8](https://doi.org/10.1016/s0895-4356(01)00377-8)
58. <sup>^</sup>Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000;56(2):455-463. <https://doi.org/10.1111/j.0006-341x.2000.00455.x>
59. <sup>^</sup>Egger M, Smith GD, Schneider M, Minder C. Bias in metaanalysis detected by a simple, graphical test. *BMJ* 1997;315(7109):629-634. <https://doi.org/10.1136/bmj.315.7109.629>
60. <sup>^</sup>Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;1088-1101. PMID: 7786990
61. <sup>^</sup>Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, Vist GE, Falck-Ytter Y, Meerpohl J, Norris S, Guyatt GH. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64(4):401-406. <https://doi.org/10.1016/j.jclinepi.2010.07.015>
62. <sup>a, b, c, d, e, f, g, h, i, j, k, l, m, n, o</sup>Chubak J, Adler A, Bobb JF, Hawkes RJ, Ziebell RA, Pocobelli G, Ludman EJ, Zerr DM. (2024). A Randomized Controlled Trial of Animal-assisted Activities for Pediatric Oncology Patients: Psychosocial and Microbial Outcomes. *J Pediatr Health Care* 2024;38(3):354–364. <https://doi.org/10.1016/j.pedhc.2023.09.010>
63. <sup>a, b, c, d, e, f, g, h, i, j</sup>Mahoney AB, Akard TF, Cowfer BA, Dietrich MS, Newton JL, Gilmer MJ. Impact of Animal-Assisted Interaction on Anxiety in Children With Advanced Cancer and Their Caregivers. *J Palliat Med* 2024;27(1):75–82. <https://doi.org/10.1089/jpm.2023.0091>
64. <sup>a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p</sup>McCullough A, Jenkins MA, Ruehrdanz A, Gilmer MJ, Olson J, Pawar A, Holley L, Sierra-Rivera S, Linder DE, Pichette D, Grossman NJ, Hellman C, Guérin NA, O’Haire ME. Physiological and behavioral effects of animal-assisted interventions on therapy dogs in pediatric oncology settings. *Applied Anim Behav Sci* 2018;200:86-95. <https://doi.org/10.1016/j.applanim.2017.11.014>
65. <sup>a, b</sup>Sherman SA, Eisen S, Burwinkle TM, Varni JW. The PedsQL Present Functioning Visual Analogue Scales: preliminary reliability and validity. *Health Qual Life Outcomes* 2006;4:75. <https://doi.org/10.1186/1477-7525-4-75>
66. <sup>^</sup>Spielberger CD. *State-Trait Anxiety Inventory for Children*. Palo Alto, CA: Consulting Psychologist Press; 1973.
67. <sup>^</sup>Varni JW, Seid M, Rode CA. The PedsQL: measurement model for the pediatric quality of life inventory. *Med Care* 1999;37(2):126–139. <https://doi.org/10.1097/00005650-199902000-00003>
68. <sup>^</sup>Marteau TM, Bekker H. (1992). The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *Br J Clin Psychol* 1992;31(3):301-306. <https://doi.org/10.1111/j.2044-8260.1992.tb00997.x>
69. <sup>^</sup>Spielberger CD. *State-Trait Anxiety Inventory for Adults*. Mind Garden Inc.: Menlo Park, CA; 1983; pp. 1–75.
70. <sup>^</sup>Glenk LM, Kothgassner OD, Stetina BU, Palme R, Kepplinger B, Baran H. Therapy dogs' salivary cortisol levels vary during animal-assisted interventions. *Anim Welf* 2013;22:369-378. <https://doi.org/10.7120/09627286.22.3.369>
71. <sup>a, b</sup>Lefebvre SL, Waltner-Toews D, Peregrine AS, Reid-Smith R, Hodge L, Weese JS. Characteristics of programs involving canine visitation of hospitalized people in Ontario. *Infect Control Hosp Epidemiol* 2006; 27(7):754-758. <https://doi.org/10.1086/505099>
72. <sup>^</sup>Wang GD, Zhai W, Yang HC, Wang L, Zhong L, Liu YH, Fan RX, Yin TT, Zhu CL, Poyarkov AD, Irwin DM, Hytönen MK, Lohi H, Wu CI, Savolainen P, Zhang YP. Out of southern East Asia: the natural history of domestic dogs across the world. *Cell Res* 2016;26:21-33. <https://doi.org/10.1038/cr.2015.147>

73. <sup>a, b, c</sup>Centers for Disease Control and Prevention, National Center for Infectious Diseases. *Healthy Pets, Healthy People*; 2024. <https://www.cdc.gov/healthy-pets/index.html> Accessed July 3, 2024
74. <sup>^</sup>Dalton KR, Waite KB, Ruble K, Carroll KC, DeLone A, Frankenfield P., Serpell JA, Thorpe Jr RJ, Morris DO, Agnew J, Rubenstein RC, Davis MF. *Risks associated with animal-assisted intervention programs: A literature review. Complement Ther Clin Pract* 2020;39:101145. <https://doi.org/10.1016/j.ctcp.2020.101145>
75. <sup>^</sup>Robinson RA, Pugh RN. (2002). *Dogs, zoonoses and immunosuppression. J R Soc Promot Health* 2002;122(2):95-98. <https://doi.org/10.1177/146642400212200210>
76. <sup>^</sup>Sehulster L, Chinn RY, CDC, HICPAC (2003). *Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). MMWR Recomm Rep* 2003;52(RR-10):1-42. PMID: 12836624