

Liver Function Test Abnormalities and Associated Factors Among Liver Disease Patients at the University of Gondar Comprehensive Specialized Hospital Northwest, Ethiopia: Cross Sectional Study

Abateneh Melkamu¹, Yalew Much¹, Abebe Yensew¹, Melaku Laikemariam¹, Befikad Mandefro², Adane Adugna¹, Gashaw Azanaw Amare¹, Aytenew Atnaf¹

¹ Debre Markos University

² Arba Minch University

Funding: No specific funding was received for this work.

Potential competing interests: No potential competing interests to declare.

Abstract

Background: Liver disease is a disturbance of the normal, and healthy performance of the liver. It is directly linked to liver synthesis, excretion and metabolism as a result it causes liver function disorder. Thus, the purpose of this study was to evaluate the extent of liver function abnormalities and the factors that are related to it in patients with liver disease.

Methods: The study was done on 307 liver-diseased patients using consecutivesampling from August 22, 2022 to October 24, 2022 at the University of Gondar Comprehensive Specialized Hospital using cross-sectional study design. About 3 ml of venous blood specimen was collected and analyzed by *Bechman coulter 700 chemistry analyzer*. Data were analyzed using SPSS version 25 software. Frequencies and proportions were used to characterize the results. Bivariable and multivariable logistic regression were utilized to investigate the factors associated with abnormalities in liver function. P value <0.05 was considered as statistically significant.

Result: Among 307 patients, 117 (38.11%, 95% CI: 32.64%, 43.57%), 212 (69.06%; 95% CI: 63.85%, 74.25%), 168 (54.72%, 95% CI: 49.12%, 60.32%), 171 (55.70%, 95% CI: 50.11%, 61.28%) patients had abnormal ALT, AST, and total bilirubin respectively. Nearly one fourth (25.40%) of the patients had all liver function tests abnormalities. Being male (AOR = 0.17; 95% CI: 0.08, 0.38), presence of viral hepatitis (AOR = 3.85; 95% CI: 1.23, 12.02) are significantly associated with abnormal liver ALT levels. Presence of blood transfusion history (AOR=0.45; 95%CI: 0.21, 0.95), lack of vegetable feeding habit (AOR=2.73; 95%CI: 1.34, 5.59), lack of physical exercise habit (AOR=2.97; 95%CI: 1.44, 6.11) and ALD (AOR=17.09; 95%CI: 1.96, 148.98) significantly associated with AST abnormality. Being anemic (AOR=3.33; 95%CI: 1.58, 7.08), lack of vegetable feeding habit (AOR=1.98; 95%CI: 1.06, 3.69), and lack of physical exercise habit (AOR=4.03; 95%CI: 1.90, 8.57) significantly associated with total bilirubin abnormality.

Conclusion: In this study, patients with liver disease had substantial liver function test abnormalities. Viral hepatitis associated to rising only ALT, while ALD only associated with the increment of AST. Patients having a transfusion

history, lacking physical activity, and lack of vegetable associate with rising bilirubin. Therefore, clinicians should consider these factors in the management and treatment of liver disease patients.

Abateneh Melkamu^{1,*}, Yalew Muche¹, Abebe Yensew¹, Melaku Laikemariam², Befikad Mandefro³, Adane Adugna¹, Gashaw Azanaw Amare¹, and Ayteneu Atnaf¹

¹*Department of Medical Laboratory Science, College of Medicine and Health Sciences, Debre Markos University, Debre Markos, Ethiopia*

²*Department of midwifery, College of Medicine and Health Sciences, Debre Markos University, Debre Markos, Ethiopia*

³*Department of medical laboratory science, College of Medicine and Health Sciences, Arba Minch University, Arba Minch, Ethiopia*

***Corresponding Author:** Abateneh Melkamu, Department of Medical Laboratory Science, College of Medicine and Health Sciences, Debre Markos University, Debre Markos, Ethiopia. P.O. Box 269; email: abatenehmelkamu@gmail.com, abateneh_melkamu@dmu.edu.et;

Keywords: Alanine Transaminase, Aspartate Transaminase, Bilirubin, Hepatic disease, Liver disease, Gondar, Ethiopia.

Background

The liver is the largest essential organ in the body, located in the right upper quadrant below the diaphragm. It performs about 5,000 different physiological activities. The liver performs the process of detoxifying different metabolites, producing digestive enzymes, and synthesizing proteins to keep other bodily systems running smoothly ^[1].

Liver disease is one of the world's most critical public health problems ^[2] that can affect liver cells and impair normal liver function ^[3]. It is characterized by increased liver biochemistry, coagulopathy, and hepatic encephalopathy without underlying chronic liver disease ^[4]. The most common causes of the rising prevalence of liver disease are non-alcoholic fatty liver disease (NAFLD), which is brought on by obesity, diabetes, autoimmunity, hemochromatosis, and chronic viral hepatitis, primarily from the hepatitis B and hepatitis C viruses. Other causes and types of liver disease caused by medication and chemicals ^[2].

The LFT is among the most commonly used laboratory investigations to measure its normality. Abnormal results indicate liver disease, even when a person does not have symptoms. Despite of this, LFT detects the severity of liver diseases, the effect of therapies, and drug-induced liver injury ^[5]. The most common causes of the rising burden of abnormal liver function tests in liver disease are nonalcoholic fatty liver disease, alcohol-related liver disease, and viral hepatitis ^[6].

Liver function tests are groups of blood, urine, and stool tests that provide information about a patient's liver condition.

Blood LFT include liver enzyme such as aspartate aminotransferase (AST), alanine aminotransferase (ALT) additionally total bilirubin are main valuable biomarkers of liver injury in a patient with some degree of abnormal liver function [7].

Liver illness is becoming increasingly widespread in developing nations, with an estimated two million fatalities each year and accounting for 4% of all deaths [8]. There is research on the prevalence of liver disease in Ethiopia, despite its consequences, there is a scarcity of information on the liver function test abnormalities and associated factors among liver disease patients in the study area. In this study, the authors aim to evaluate the extent of liver function abnormalities, associated factors among liver disease patients at the University of Gondar Comprehensive Specialized Hospital Northwest, Ethiopia, in 2022

Methods and Materials

Study design, area, and period

Cross-sectional research was carried out at the University of Gondar Comprehensive Specialized Hospital (UoG-CSH) from August 22,2022, to October 24, 2022. The hospital is located in Gondar. Gondar is located 727 kilometers northwest of Ethiopia's main city, Addis Ababa, and 175 kilometers from Bahir Dar, the seat of the Amhara National Regional State. According to the 2015 report of the central statistical agency of Ethiopia, Gondar has a population of 323,900 [9]. The town has one public comprehensive specialty hospital, which is one of the oldest teaching hospitals in the country and offers health services for more than 7 million people in Gondar town and its catchment regions [10].

Operational definitions

- **Habit of drinking tea or coffee:** Habitual tea/coffee drinkers are defined as tea/coffee drinking of 120 mL or more per day for at least one year [11].
- **Physical exercise:** Participant who performs daily active exercise for about 30 minutes/day [12].
- **Habit of feeding vegetables:** It is a consistent action to integrate a variety of vegetables into one's diet or meal plan [13].
- **Habit of feeding meat:** It is the consumption of animal-derived proteins such as beef, poultry, pork, fish, or other meats as a primary source of nutrition in one's daily diet [14].
- **AST normal:** If the serum levels were from 0 to 35 U/L for males and 0 to 31 U/L for females [15].
- **ALT normal:** If the serum levels were from 0 to 45 U/L for males and 0 to 34 U/L for females [15].
- **Normal serum total bilirubin:** If the reference range from 0 to 1.2 mg/dl [15].

Population, variables, and sampling techniques

Patients with liver illness who attended the UoG-CSH during the data collecting period and met the inclusion criteria were considered a study population. Although the liver function parameters (ALT, AST, and total bilirubin) were taken as

dependent variables, the socio-demographic variables, clinical variables and behavioural related variables were taken as independent variables. A total of 307 consecutively selected liver disease patients were included in this study. Study participants with critically ill patients and pregnant women, had excluded from the study.

Data collection methods and data quality management

A pretested structured questionnaire was used to gather the sociodemographic, lifestyle, and information about diets using a face-to-face interview. A data extraction sheet was used to get the clinical data from the patient's medical files. Three milli liter of venous blood collected by needle and syringe method. liver function tests were carried out by a Beckman Coulter DXC 700, (*United States of America*) chemistry analyzer and. To maintain the quality of the data, quality control testing was performed for each procedure. Furthermore, standard operating procedures were strictly followed. Training was given to all data collectors prior to the actual data collection. The investigator strictly controls the data collector and general activity during the data collection period.

Statistical analysis

Epidata version 3.1 was utilized for data entry, coding, and cleaning. The statistical package for the social sciences (SPSS) version 25 software was used to import and analyze the data. To summarize the data, summary statistics, proportions, and frequencies were employed. Pearson rank chi-square assumption fulfillment was checked for categorical variables. Bivariable and multivariable logistic regression were used to determine factors associated with liver function abnormalities. To verify the model fitness assumption, the Hosmer-Lemeshow goodness of fit test was performed, requiring a p-value of larger than 0.05. Finally, the odds ratio with a 95% confidence interval was used to express the strength of the association. The multivariable analyses with p-value < 0.05 indicated variables had a significant association with the result.

Results

Socio-demographic and clinical characteristics of study participants

This research had 307 individuals. Among them, 220 (71.66%), 213 (69.38%), 181 (59.28%), and 143 (46.58%) were male, from rural areas, married, and unable to read and write. Furthermore, the average age of the research participants was 38.38 + 15.13 years, with a range of 6 to 82 years.

From a total of study participants, 157 (51.14%), 87 (28.34%), 37 (12.05%), and 26 (8.47%) of study participants had chronic liver disease (CLD), viral hepatitis, acute liver disease, and ALD, respectively. Among viral hepatitis study participants, about 71.26% and 28.74 had HBV and HCV, respectively. On the other hand, about 51 (16.61%), 39 (12.70%), and 33 (10.75%) were anemic, had a history of blood transfusions, and had heart disease, respectively additionally about 86 (28.01%) of the study participants had taken medication other than liver disease drugs. (Table 1).

Table 1. Clinical characteristics of liver disease patients at UoG-CSH Northwest, Ethiopia (n =307).

Variable	Category	Frequency	Percent (%)
Use of medication other than liver disease	Yes	86	28.01
	No	221	71.99
History of anemia	Yes	51	16.61
	No	256	83.61
History of tuberculosis	Yes	12	3.90
	No	295	96.10
Presence of DM	Yes	13	4.20
	No	294	95.80
Presence of cardiac disease	Yes	31	10.10
	No	276	89.90
History of blood transfusion	Yes	39	12.70
	No	268	87.30
History of surgery	Yes	10	3.30
	No	297	96.70
Presence of cancer	Yes	8	2.60
	No	299	97.40
Presence of hypertension	Yes	21	6.80
	No	286	93.2
Presence of HIV/AIDS	Yes	6	2.0
	No	301	98.0
Types of liver disease	Acute liver disease	37	12.05
	CLD	157	51.14
	ALD	26	8.47
	Viral hepatitis	87	28.34
Presence of viral hepatitis	Yes	87	28.34
	No	220	71.66

Abbreviations: **ALD:** Alcoholic Liver Disease, **CLD:** Chronic Liver Disease, **HIV;** Human Immunodeficiency Virus, **DM;** Diabetes Miletus

Nutritional and life style characteristics of study participants

Of the study participants, about 269 (87.91%), 287 (93.49%), and 237 (77.1%) had a habit of feeding meat, drinking tea or coffee, and consuming vegetables, respectively. despite this, many of the study participants about 262 (85.34%) had not regularly physical exercise habit, and only 11 (3.58%) of study participant had smoking habit (Table 2).

Table 2. Lifestyle characteristics of liver disease patients at UoG-CSH Northwest, Ethiopia (n=307).

Variable	Category	Frequency	Percent (%)
Tea or coffee drinking habit	Yes	270	88.24
	No	37	11.76
Meat feeding habit	Yes	284	92.51
	No	23	7.49
Vegetable feeding habit	Yes	237	7.2
	No	70	22.8
Alcohol drinking habit	Yes	205	66.8
	No	102	33.2
Cigarette smoking habit	Yes	11	3.58
	No	296	96.42
Physical exercise habit	Yes	45	14.66
	No	262	85.34

Magnitude of liver function abnormalities

From the total study participants, 117 (38.11%, 95% CI: 32.64%, 43.57%), 212 (69.06%; 95% CI: 63.85%, 74.25%), 168 (54.72%, 95% CI: 49.12%, 60.32%), of the study participants had abnormal ALT, AST, and total bilirubin respectively. Furthermore, 78 (25.40%) study participants had all liver function abnormalities (Figure 1).




Figure 1. Magnitude of liver function test abnormalities among liver disease patients at the UoG-CSH, Northwest Ethiopia (n=307)

Table 3. Liver function abnormalities tabulated with ALT, AST, and total bilirubin among liver diseased study participants at the UoG-CSH, Northwest Ethiopia (n =307).

Variable	Category	ALT		AST		Total Bilirubin	
		Abnormal	Normal	Abnormal	Normal	Abnormal	Normal
Sex	Male	70	150	148	72	122	98
	Female	47	40	64	23	46	41
Age	<18 (children)	4	24	18	10	14	14
	18-45 (young adult)	70	118	127	61	106	82
	>45 (old adult)	43	48	67	24	48	43
Residence	Urban	36	58	64	30	51	43
	Rural	81	132	148	65	117	96
Occupation	Farmer	74	103	126	51	98	79
	Housewife	16	17	25	8	16	17
	Merchant	5	18	13	10	12	11
	Government	5	16	13	8	9	12
	Private employee	10	9	13	6	14	5
	Other	7	27	22	12	19	15
Educational level	No education	60	83	96	47	82	61
	Primary school	41	78	87	32	64	55
	High school	11	15	18	8	16	10
	University, college	5	14	11	8	6	13
Marital status	Married	66	115	121	60	95	86
	Unmarried	51	75	91	35	73	53
Use of medication other than liver disease	Yes	33	53	60	26	44	42
	No	84	137	152	69	124	97
History of anemia	Yes	26	25	41	10	38	13
	No	91	165	171	85	130	126
History of tuberculosis	Yes	5	7	7	5	7	5
	No	112	183	205	90	161	134
Presence of DM	Yes	4	9	11	2	6	7
	No	113	181	201	93	162	132
Presence of cardiac disease	Yes	9	22	23	8	16	15
	No	108	168	189	87	152	124
History of blood transfusion	Yes	11	28	21	18	17	22
	No	106	162	191	77	151	117
Have a history of surgery	Yes	3	7	6	4	5	5
	No	114	183	206	91	163	134
Presence of cancer	Yes	2	6	5	3	6	2
	No	115	184	207	92	162	137
Presence of hypertension	Yes	7	14	17	4	10	11
	No	110	176	195	91	158	128

Presence of HIV disease	Yes	4	2	4	2	2	4
	No	113	188	208	93	166	135
Consumption of tea or coffee	Yes	104	166	189	81	149	121
	No	13	24	23	14	19	18
Meat feeding habit	Yes	107	177	195	89	155	129
	No	10	13	17	6	13	10
Vegetable feeding habit	Yes	82	155	155	82	122	115
	No	35	35	57	13	46	24
Alcohol drinking habit	Yes	73	132	146	59	119	86
	No	44	58	66	36	49	53
Cigarette smoking habit	Yes	4	7	7	4	4	7
	No	113	183	205	91	164	132
Physical exercise habit	Yes	15	30	22	23	13	32
	No	102	160	190	72	155	107
Types of liver disease	Acute liver disease	8	29	25	12	18	19
	CLD	48	109	95	62	80	77
	ALD	7	19	25	1	18	8
	Viral hepatitis	54	33	67	20	52	32
Presence of virus hepatitis (HBV and HCV)	Yes	30	57	55	32	44	43
	No	87	133	157	63	124	96

Abbreviations: **ALD:** Alcoholic Liver Disease, **CLD:** Chronic Liver Disease, **HIV=** Human Immunodeficiency Virus, **DM:** Diabetes Miletus, **Other** includes student and no jobs

Factors associated with liver function abnormalities

The bivariable logistic regression analysis, patients' age group, gender, habit of eating vegetables, viral hepatitis, and anemia were linked to abnormal liver ALT levels. After multivariable logistic regression analysis, gender being male (AOR = 0.17; 95% CI: 0.08, 0.38), presence of viral hepatitis (AOR = 3.85; 95% CI: 1.23, 12.02) were significantly associated (Table 4).

Table 4. Bivariable and multivariable logistic regression of ALT among liver disease patients attending at the UoG-CSH, Northwest, Ethiopia (n=307)

Variables	Category	ALT		Bivariable analysis		Multivariable
		Abnormal (%)	Normal (%)	COR (95%CI)	P-value	AOR (95%CI)
Sex	Male	70 (31.82)	150(61.18)	0.39 (0.23,0.66)	<0.001	0.17(0.08,0.38) *
	Female	47(54.02)	40(45.98)	1		1
Age	<18 (children)	4(16.66)	24(83.34)	1		1
	18-45 (young adult)	70(37.24)	118(62.76)	3.55(1.18,10.68)	0.024	2.92(0.49,17.24)
	>45 (old adult)	43(47.23)	48(52.77)	5.35(1.72,16.73)	0.004	4.66(0.73,29.68)

Residence	Rural	36(38.29)	58(61.71)	0.98(0.59,1.62)	0.964	NA
	Urban	81(38.02)	132(61.98)	1		
Occupation of study participants	Farmer	74(41.81)	103(58.19)	2.29(0.80,6.55)	0.119	3.21(0.53,19.2)
	Housewife	16(48.49)	17(51.51)	3.01(0.89,10.14)	0.075	0.84(0.11,6.24)
	Merchant	5(21.74)	18(78.26)	0.88(0.21,3.64)	0.870	1.11(0.14,8.45)
	Private employee	10(52.63)	9(47.37)	3.55(0.92,13.69)	0.065	2.95(0.40,21.52)
	Other	7(20.59)	27(79.41)	0.82(0.22,3.05)	0.779	1.46(0.20,10.34)
	Government employee	5(23.81)	16(76.19)	1		1
Educational level	No education	60(41.95)	83(58.05)	2.02(0.69,5.92)	0.198	1.48(0.21,10.13)
	Primary school	41(34.45)	78(65.55)	1.47(0.49,4.37)	0.752	1.84(0.27,12.43)
	High school	11(42.31)	15(57.69)	2.05(0.56,7.41)	0.272	2.73(0.37,20.08)
	University & college	5(26.32)	14(73.68)	1		1
Marital status	Unmarried	66(36.46)	115(63.54)	0.84(0.52,1.34)	0.477	NA
	Married	51(40.47)	75(59.53)	1		
Use of medication other than liver disease	Yes	33(38.37)	53(61.63)	1.01(0.60,1.69)	0.953	NA
	No	84(38.01)	137(61.99)	1		
History of anemia	Yes	26(50.98)	25(49.02)	1.88(1.02,3.45)	0.040	1.53(0.76,3.10)
	No	91(35.55)	165(64.45)	1		1
Presence of DM	Yes	5(41.66)	7(58.34)	0.71(0.21,2.36)	0.579	NA
	No	112(37.96)	183(62.04)	1		
Presence of cardiac disease	Yes	9(29.04)	22(70.96)	0.63(0.28,1.43)	0.275	NA
	No	108(39.14)	168(60.86)	1		
History of blood transfusion	Yes	11(28.21)	28(71.79)	0.60(0.28,1.25)	0.176	0.70(0.30,1.61)
	No	106(39.55)	162(60.45)	1		1
Presence of hypertension	Yes	7(33.34)	14(66.66)	0.8(0.31,2.04)	0.641	NA
	No	110(38.46)	176(61.54)	1		
Tea or coffee drinking habit	No	104(38.52)	166(61.48)	0.86(0.42,1.77)	0.691	NA
	Yes	13(35.14)	24(64.86)	1		
Meat feeding habit	Yes	107(37.67)	177(63.33)	0.78(0.33,1.85)	0.582	NA
	No	10(43.47)	13(56.53)	1		
Vegetable feeding habit	No	35(50.00)	35(50.00)	1.89(1.10,3.24)	0.021	1.54(0.83,2.86)
	Yes	82(34.60)	155(65.40)	1		1
Alcohol drinking habit	Yes	73(35.61)	132(64.39)	0.72(0.44,1.18)	0.202	0.65(0.37,1.16)
	No	44(43.14)	58(56.86)	1		1
Cigarette smoking habit	Yes	4(36.36)	7(63.64)	0.92(0.26,3.23)	0.903	NA
	No	113(38.18)	183(61.82)	1		
Physical exercise habit	No	15(33.33)	30(66.67)	0.78(0.40,1.52)	0.476	NA
	Yes	102(38.93)	160(61.07)	1		
Types of liver disease	CLD	48(30.57)	109(69.43)	1.59(0.68,3.74)	0.283	0.99(0.33,2.98)
	ALD	7(26.93)	19(73.07)	1.33(0.41,4.29)	0.627	1.38(0.32,5.83)
	Viral hepatitis	54(62.06)	33(37.94)	5.93(2.42,14.51)	<0.001	3.85(1.23,12.02) *

	Acute liver disease	8(21.62)	29(78.38)	1		1
Presence of virus hepatitis (HBV and HCV)	Yes	30(34.48)	57(65.52)	0.80 (0.47,1.35)	0.411	NA
	No	87(39.54)	133(60.46)	1		

Abbreviation: * = Significant Variable, **ALD:** Alcoholic Liver Disease, **CLD:** Chronic Liver Disease **DM=**Diabetes mellitus **AOR=** Adjusted odds ratio, **COR:** Crude Odd Ratio, **CI:** Confidence Interval, **1=** reference group **other=** includes student and have no jobs, **NA=**not applicable

Using bivariable regression analysis, AST abnormalities was linked to independent factors such as the history of blood transfusion, vegetable eating habits, physical activity, and ALD. After multivariable logistic regression analysis, presence of blood transfusion history (AOR=0.45; 95%CI: 0.21, 0.95), lack of vegetable feeding habit (AOR=2.73; 95%CI: 1.34, 5.59), and not doing physical exercise (AOR=2.97; 95%CI: 1.44, 6.11) and ALD (AOR=17.09; 95%CI: 1.96, 148.98) remained significantly associated (Table 5).

Table 5. Bivariable and multivariable logistic regression of abnormal AST among liver disease patients attending at the UoG-CSH, Northwest, Ethiopia, 2022 (n=307)

Variable	Category	AST		Bivariable analysis		Multivariable analysis
		Abnormal (%)	Normal (%)	P-value	COR (95%CI)	AOR (95%CI)
Sex	Male	148(67.27)	72(32.73)	0.284	0.73 (0.42,1.28)	NA
	Female	64(73.56)	23(26.44)		1	
Age	<18 (children)	18(64.29)	10(35.71)		1	NA
	18-45 (young adult)	127(67.55)	61(32.45)	0.731	1.15(0.50,2.65)	
	>45 (old adult)	67(73.63)	24(26.37)	0.341	1.55 (0.62,3.82)	
Residence	Rural	148(69.48)	65(30.52)	0.807	1.06(0.63,1.79)	NA
	Urban	64(68.09)	30(31.91)		1	
Occupation	Farmer	126(71.18)	51(28.81)	0.382	1.52(0.59,3.88)	NA
	Housewife	25(75.75)	8(24.25)	0.280	1.92(0.58,6.30)	
	Merchant	13(56.53)	10(43.47)	0.717	0.8(0.23,2.67)	
	Private employee	13(68.42)	6(31.58)	0.666	1.33(0.36,4.93)	
	Other	22(64.70)	12(35.30)	0.834	1.12(0.36,3.48)	
	Government employee	13(61.90)	8(38.10)		1	
Educational level	No education	96(67.13)	47(32.87)	0.426	1.48(0.56,3.93)	1.40(0.42,4.62)
	Primary school	87(73.10)	32(26.90)	0.180	1.97(0.72,5.35)	2.11(0.63,7.06)
	High school	18(69.23)	8(30.77)	0.434	1.63(0.47,5.62)	1.50(0.35,6.29)
	University & College	11(57.90)	8(42.10)		1	1
Marital status	Unmarried	121(66.85)	60(33.15)	0.317	1.12(0.78,2.12)	NA
	Married	91(72.22)	35(27.78)		1	
Use of medication other than liver disease	Yes	60(69.77)	26(30.23)	0.866	1.04(0.60,1.79)	

	No	152(68.78)	69(31.22)		1	
History of anemia	Yes	41(80.40)	10(19.60)	0.059	2.03(0.97,4.26)	2.11(0.96,4.61)
	No	171(66.80)	85(32.20)		1	1
Presence of DM	Yes	11(84.61)	2(15.39)	0.230	2.54(0.55,11.71)	NA
	No	201(68.36)	93(31.64)		1	
Presence of cardiac disease	Yes	23(74.20)	8(25.80)	0.515	1.32(0.56,3.07)	NA
	No	189(68.47)	87(31.52)		1	
History of blood transfusion	Yes	21(53.85)	18(46.15)	0.030	0.47(0.23,0.93)	0.45(0.21 0.95) *
	No	191(71.27)	77(28.73)		1	1
Presence of hypertension	Yes	17(80.95)	4(19.05)	0.230	1.98(0.64,6.06)	NA
	No	195(68.18)	91(31.82)		1	
Tea or coffee drinking habit	No	189(70.00)	81(30.00)	0.335	0.70(0.34,1.43)	NA
	Yes	23(62.16)	14(37.84)		1	
Meat feeding habit	Yes	195(68.66)	89(31.34)	0.601	0.77(0.29,2.02)	NA
	No	17(73.91)	6(26.09)		1	
Vegetable feeding habit	No	155(65.40)	82(34.60)	0.012	2.31(1.19,4.48)	2.73(1.34,5.59) *
	Yes	57(81.43)	13(18.57)		1	1
Alcohol drinking habit	Yes	146(71.22)	59 (28.78)	0.246	1.34(0.81,2.23)	NA
	No	66(64.70)	36(35.30)		1	
Cigarette smoking habit	Yes	7(63.64)	4(36.36)	0.693	0.77(0.22,2.71)	NA
	No	205(69.26)	91(30.74)		1	
Physical exercise habit	No	22(48.89)	23(51.11)	0.002	2.75(1.44,5.25)	2.97(1.44,6.11) *
	Yes	190(72.52)	72(27.48)		1	1
Type of liver disease	CLD	95(60.50)	62(39.50)	0.428	0.73(0.44,1.57)	0.83(0.36,1.89)
	ALD	25(96.15)	1(3.85)	0.021	11.99(1.44,99.37)	17.09(1.96,148.98) *
	Viral hepatitis	67(77.01)	20(22.99)	0.274	1.60(0.68,3.76)	1.75(0.70,4.38)
	Acute liver disease	25(67.57)	12(32.43)		1	1
Presence of virus hepatitis (HBV and HCV)	Yes	55(63.22)	32(36.78)	0.165	0.68(0.40,1.16)	0.74(0.41,1.33)
	No	157(71.36)	63(28.64)		1	1

Abbreviation: *= Significant Variable, **ALD:** Alcoholic Liver Disease, **CLD:** Chronic Liver Disease **DM=**Diabetes mellitus, **AOR=** Adjusted odds ratio, **COR=**Crude Odd Ratio, **CI=**Confidence Interval, **1=** the reference group, **other=** includes student and have no jobs, **NA=**not applicable

In bivariable logistic regression analysis revealed associations between total bilirubin elevation and independent factors such as lack of education, anemia, vegetable eating habits, and physical activity. After multivariable logistic regression analysis, being anemic (AOR=3.33; 95%CI: 1.58, 7.08), lack of vegetable feeding habit (AOR=1.98; 95%CI: 1.06, 3.69), and not doing physical exercise (AOR=4.03; 95%CI: 1.90, 8.57) remained significantly associated (Table 6).

Table 6. Bivariable and multivariable logistic regression of abnormal Total bilirubin among liver disease patients attending at the UoG-CSH, Northwest, Ethiopia (n=307)

Variable	Category	Total Bilirubin		Bivariable analysis		Multivariable analysis
		Abnormal (%)	Normal (%)	P-value	COR (95%CI)	AOR (95%CI)
Sex	Male	122(55.45)	98(44.55)	0.682	1.10 (0.67,1.82)	NA
	Female	46(52.87)	41(47.13)		1	
Age	<18 (children)	14(50.00)	14(50.00)		1	NA
	18-45 (young adult)	106(56.38)	82(47.62)	0.527	1.29(0.58,2.86)	
	>45 (old adult)	48(52.75)	43(47.25)	0.799	1.11 (0.47,2.60)	
Residence	Rural	117(54.93)	96(45.07)	0.913	1.02(0.63,1.67)	NA
	Urban	51(54.26)	43(45.74)		1	
Occupation	Farmer	98(55.37)	79(44.63)	0.280	1.65(0.66,4.12)	0.44(0.07,2.74)
	Housewife	16(48.48)	17(51.52)	0.686	1.25(0.41,3.77)	0.28(0.04,2.01)
	Merchant	12(52.17)	11(47.83)	0.537	1.45(0.44,4.78)	0.45(0.64,3.22)
	Private employee	14(73.68)	5(26.32)	0.054	3.73(0.97,14.22)	1.33(0.16,10.94)
	Other	19(55.88)	15(44.12)	0.349	1.68(0.56,5.06)	0.71(0.10,4.75)
	Government employee	9(42.86)	12(57.14)		1	1
Educational level	No education	82(57.34)	61(42.66)	0.040	2.91(1.04,8.09)	4.91(0.72,33.55)
	Primary school	64(53.78)	55(46.22)	0.079	2.52(0.89,7.07)	4.53(0.68,30.11)
	High school	16(61.54)	10(38.46)	0.051	3.46(0.99,12.08)	5.42(0.74,39.68)
	University & College	6(31.58)	13(68.42)		1	1
Marital status	Unmarried	95(52.49)	86(47.51)	0.346	1.24(0.78,1.97)	NA
	Married	73(57.94)	53(42.06)		1	
Use of medication other than liver disease	Yes	44(51.16)	42(48.84)	0.435	0.81(0.49,1.35)	NA
	No	124(56.10)	97(43.90)		1	
History of anemia	Yes	38(74.50)	13(25.50)	0.003	2.83(1.44,5.56)	3.33(1.58,7.08) *
	No	130(50.78)	126(49.22)		1	1
Presence of DM	Yes	7(58.33)	5(41.67)	0.528	0.69(0.22,2.12)	NA
	No	161(54.58)	134(45.42)		1	
Presence of cardiac disease	Yes	16(51.61)	15(48.39)	0.714	0.87(0.41,1.82)	NA
	No	152(55.07)	124(44.93)		1	
History of blood transfusion	Yes	17(43.59)	22(56.41)	0.138	0.59(0.30,1.17)	0.62(0.29,1.29)
	No	151(56.34)	117(43.66)		1	1
Presence of hypertension	Yes	10(47.62)	11(52.38)	0.499	0.73(0.30,1.78)	NA
	No	158(55.24)	128(44.76)		1	
Tea or coffee drinking habit	No	149(55.19)	121(44.81)	0.661	0.85(0.43,1.70)	NA
	Yes	19(51.35)	18(48.65)		1	
Meat feeding habit	Yes	155(54.58)	129(45.42)	0.857	0.92(0.39,2.17)	NA
	No	13(56.52)	10(43.48)		1	
Vegetable feeding habit	No	122(51.48)	115(48.52)	0.037	1.80(1.03,3.14)	1.98(1.06,3.69) *

	Yes	46(65.71)	24(34.29)		1	1
Alcohol drinking habit	Yes	119(58.05)	86(41.95)	0.098	1.49(0.92,2.41)	1.21(0.71,2.07)
	No	49(48.04)	53(51.96)		1	1
Cigarette smoking habit	Yes	4(36.36)	7(63.64)	0.223	0.45(0.13,1.60)	NA
	No	164(55.40)	132(44.60)		1	
Physical exercise habit	No	13(28.89)	32(71.11)	0.000	3.56(1.78,7.10)	4.03(1.90,8.57) *
	Yes	155(59.16)	107(40.84)		1	1
Type of liver disease	CLD	80(50.96)	77(49.04)	0.801	1.09(0.53,2.24)	1.42(0.58,3.48)
	ALD	18(69.23)	8(30.77)	0.107	2.37(0.82,6.80)	3.21(0.90,11.36)
	Viral hepatitis	52(61.90)	32(38.10)	0.255	1.56(0.72,3.40)	1.88(0.72,4.93)
	Acute liver disease	18(48.65)	19(51.35)		1	1
Presence of virus hepatitis (HBV and HCV)	Yes	44(50.57)	43(49.43)	0.359	0.79(0.48,1.30)	NA
	No	124(56.36)	96(43.64)		1	

Abbreviation: *= Significant Variable, **ALD:** Alcoholic Liver Disease, **CLD:** Chronic Liver Disease **DM=**Diabetes mellitus, **AOR=** Adjusted odds ratio, **COR=**Crude Odd Ratio, **CI=**Confidence Interval, **1=** the reference group, **other=** includes student and have no jobs, **NA=**not applicable

Discussion

Liver disease is a global public health issue that causes mortality and morbidity. It is one of the most common causes of coagulopathy and impaired liver function in both developed and developing countries. This study investigated on the extent and associated factors influencing liver function abnormalities in liver disease patients visiting UoG-CSH.

The findings of this study showed that the overall prevalence of abnormal liver function test results was 117 (38.11%, 95% CI: 32.64%, 43.57%), 212 (69.06%; 95% CI: 63.85%, 74.25%), 168 (54.72%, 95% CI: 49.12%, 60.32%), of the study participants had abnormal ALT, AST, and total bilirubin respectively. Furthermore, 78 (25.40%) study participants had all liver function abnormalities. When liver cells are damaged, they release ALT, AST, and increase bilirubin into the bloodstream. ALT is preferably specific indicator than AST for active hepatic cell damage due to its location. The rising of blood AST indicates the occurrence of other events besides liver damage [16].

In this study Patient's sex and hepatic virus were showed statistically significant association with rising of ALT. The likelihood of patient's ALT abnormality occurrence being male was reduced by 83%, which contradicts with the study done in China [17], Taiwan [18]. This could be associated with BMI mostly males had good BMI which is significantly associated with less liver function test abnormality, the previous study showed Males have greater risk of abnormal liver function by controlling hepatitis B, obesity. Being patients infected by hepatic virus(s) the odds of developing ALT abnormality is 3.85 (95% CI: 1.23, 12.02) times likely than other type hepatic disease similar with the previous study in China [17][19], Serbia [6], and Korea [20]. The virus can multiply in the host and more tissue are damaged in short time leads to release more ALT enzyme [21].

Patient's status of blood transfusion history, vegetable dietary habit, and regular physical exercise, and ALD were showed statistically significant association with rising of AST. Patients having blood transfusion history less likely to happened AST abnormality by 55% (95%CI: 0.21, 0.95) than non-transfused which similar with the study conducted in USA [22]. Liver disease patients who lacked of vegetable feeding habit had makes AST abnormality by 2.73 (95%CI: 1.34, 5.59) times than study participant has habit of feeding vegetables, it similar with the study done in Iran [23]. Vegetables contain antioxidants may reduce the value of AST. The odds of developing AST rising were 2.97 (95%CI: 1.44, 6.11) more likely among patients lack of doing regular exercise than doing. Diet and weight loss are essential things in enhancing liver enzymes, with antioxidants provided by vegetables [24]. Exercise decreases stress on the liver, fats, increases energy levels and helps to prevent obesity which is risk factor for liver disease [25]. Lastly alcoholic liver disease patients were 17.09 (95%CI: 1.96, 148.98) times more likely to rise AST the others similar with study in Sweden [26]. Because vitamin B6 is used as coenzyme for liver activities, but alcoholism reduces the level of vitamin B6 [27].

Patient's status of anemia, vegetable dietary habit, physical exercises and blood transfusion were showed statistically significant association with rising total bilirubin. The odds of rising total bilirubin were 3.33 (95%CI: 1.58, 7.08) times more likely in anemic patients than non-anemic. Bilirubin is formed when hemoglobin is broken down. High bilirubin levels in the circulation may indicate hemolytic anemia. High amounts of this chemical are also associated with certain liver and gallbladder illnesses. Erythrocyte abnormalities were clinically important and frequent findings in patients with chronic disease [28]. In this study, patients who lack vegetable in dietary habit 1.98 (95%CI: 1.06, 3.69) times more likely to rising total bilirubin which may markers associated with cardiovascular problems. The controversial ideas between vegetables and bilirubin. When vegetables and balanced date is abnormal the red cells and immunity is affected causes to cellular damage, infection, and protein abnormalities. Therefore, balanced diet and vegetarians has good prognosis among liver disease patients [29]. Lack of physical exercise was raising the total bilirubin 4.03 (95%CI: 1.90, 8.57) times than its counterpart which is contradict with study done in Columbia [30].

Strength and Limitation of the study

This study has its strengths and limitations. The strength of this study is that it is the first study on the determination of the magnitude and associated factors of liver function abnormalities in liver disease patients in Ethiopia. However, the limitation of this study was that we could not perform parasitic infection screening, which may interfere with the finding of the study. In addition, detailed liver function tests not performed due to budget constraints.

Conclusion and Recommendation

In liver diseased patients' majority of patients rise AST followed by bilirubin. About 1/4th of the patients indicates all liver enzymes abnormalities. Females, having hepatic virus, blood transfusion history, lack vegetable dietary habit, and lack regular physical exercise, and ALD, anemic risk factors for liver function tests. And there is a moderate correlation between AST, ALT, and bilirubin with APTT and PT. We recommended, healthcare providers should monitor and consider

those factors for better improving and drug of choice.

List of abbreviations

ALD: Alcoholic Liver Disease, **ALT**: Alanine Transaminase, **AST**: Aspartate Transaminase, **CLD**: Chronic Liver Disease, **HBV**: Hepatitis B virus, **HCV**: Hepatitis C virus, **LFT**: Liver function test, **SPSS**: Statistical Package for the Social Sciences, **UoG-CSH**: University of Gondar Comprehensive Specialized Hospital.

Statements and Declarations

Acknowledgments

We'd like to thank the Department of Hematology and Immunohematology, the School of Biomedical and Laboratory Sciences, and the University of Gondar. We would also want to thank and appreciate the University of Gondar Hospital for their willingness and efforts during this data collection. Finally, we thank all study participants for their willingness to participate and for giving the essential information during data collection.

Ethical approval and consent to participate

This research was carried out by the Helsinki Declaration^[31]. An ethical clearance was obtained from the Ethics Review Committee of the School of Biomedical and Laboratory Science, College of Medicine and Health Science, University of Gondar Research, with reference number SBMLS/384/2022. Then, a permission letter was secured from the UoG-CSH medical director. Before starting data collection, the data collectors were informed to ask permission, explain the purpose of the study and its importance and benefits, and offer to answer all the participants' questions to confirm their willingness. Then, written informed consent, or assent, was obtained. Participation in the study and refusals were possible. To ensure confidentiality of data, study participants were identified using codes, and unauthorized persons had no access to the data. All abnormal laboratory findings were linked to health professionals for proper management and treatment of patients.

Availability of data and materials

All the data supporting these findings are contained within the manuscript.

Conflict of interest

The authors declare that there is no conflict of interest related to the publication of this manuscript.

Funding

The authors did not receive any funding for this work

Authors' contributions

AM: data collection, feeding, performing the statistical analysis, drafting the manuscript, YM: Data analysis, drafting and editing the manuscript, AY, ML, BM, GA, AA, AA: Review and editing the manuscript. All authors read and approved the final manuscript.

References

- [^] Monroe D, Hoffman M, Roberts H. *Williams hematology*. New York NY: McGraw-Hill Professional Publishing; 2010. p. 2191.
- ^{a, b} Cheemerla S, Balakrishnan M. *Global Epidemiology of Chronic Liver Disease*. *Clinical liver disease*. 2021;17(5):365-70.
- [^] Krishnan S. *Liver diseases-an overview*. *World journal of pharmacy and pharmaceutical sciences*. 2019;8:1385-95.
- [^] Lopes D SH. *Hepatic failure*. [Updated 2022 Jul 18] In: *StatPearls [Internet] Treasure Island (FL): StatPearls Publishing; 2022 Jan. 2022*.
- [^] Rahmioglu N, Andrew T, Cherkas L, Surdulescu G, Swaminathan R, Spector T, et al. *Epidemiology and genetic epidemiology of the liver function test proteins*. *PloS one*. 2009;4:e4435.
- ^{a, b} Mijač D, Krstić MN, Marković AP, Popović DD, Krstić JM, Milosavljević T. *Abnormal Liver blood tests: Primary care approach*. *Digestive Diseases*. 2022;40(2):215-22.
- [^] Mcclatchey KD. *Clinical laboratory medicine: Lippincott Williams & Wilkins; 2002*.
- [^] Devarbhavi H, Asrani SK, Arab JP, Nartey YA, Pose E, Kamath PS. *Global burden of liver disease: 2023 update*. *J Hepatol*. 2023;79(2):516-37.
- [^] Atsedo D. *Tegegne MAN MKD, Kumela G. Nedessa, Hone M. Belaye* https://mdl.donau-uni.ac.at/ses/pluginfile.php/314/mod_page/content/4/CITY%20PROFILE%20GONDAR.pdf. *City profile Gondar* { 29 October 2022).
- [^] Woldekidan NA, Mohammed AS. *Clinical Knowledge and Practice of "Ketofol" at University of Gondar Comprehensive Specialized Hospital*. *Frontiers in Medicine*. 2021:1518.
- [^] Dong X-X, Wang R-R, Liu J-Y, Ma Q-H, Pan C-W. *Habitual tea consumption and 5-year incident metabolic syndrome among older adults: a community-based cohort study*. *BMC Geriatrics*. 2021;21(1):728.
- [^] Diaz KM, Shimbo D. *Physical activity and the prevention of hypertension*. *Curr Hypertens Rep*. 2013;15(6):659-68.
- [^] Pem D, Jeewon R. *Fruit and Vegetable Intake: Benefits and Progress of Nutrition Education Interventions- Narrative Review Article*. *Iran J Public Health*. 2015;44(10):1309-21.
- [^] Lofgren PA. *Meat, Poultry, and Meat Products: Nutritional Value*. In: *Caballero B, editor. Encyclopedia of Human Nutrition (Third Edition)*. Waltham: Academic Press; 2013. p. 160-7.
- ^{a, b, c} Rifai N. *Tietz textbook of clinical chemistry and molecular diagnostics*. 7 ed: Elsevier Health Sciences; 2017. 1637

p.

16. [^]Rifai N. *Tietz textbook of clinical chemistry and molecular diagnostics: Elsevier Health Sciences; 2017.*
17. ^{a, b}Zhang H, He SM, Sun J, Wang C, Jiang YF, Gu Q, et al. *Prevalence and etiology of abnormal liver tests in an adult population in Jilin, China. International journal of medical sciences. 2011;8(3):254-62.*
18. [^]Association for the advancement of blood biotherapies. *Whole blood and red blood cell components 2022 [cited 2022 /28/10]. Available from: <https://www.aabb.org/regulatory-and-advocacy/regulatory-affairs/regulatory-for-blood/whole-blood-and-red-blood-cell-components>.*
19. [^]Sarin SK, Kumar M. *Should chronic HBV infected patients with normal ALT treated: debate. Hepatology international. 2008;2(2):179-84.*
20. [^]Oh JS, Choi JS, Lee YH, Ko KO, Lim JW, Cheon EJ, et al. *The Relationships between Respiratory Virus Infection and Aminotransferase in Children. Pediatric gastroenterology, hepatology & nutrition. 2016;19(4):243-50.*
21. [^]Liu B, Yang X, Zhao D, Mao X. *Viral Hepatitis. In: Li H, editor. Radiology of Infectious Diseases: Volume 1. Dordrecht: Springer Netherlands; 2015. p. 589-653.*
22. [^]Nachnani JS, Hamid F, Pandya P, Clarkston W, Alba LM. *Transfusion-related acute hepatic enzyme elevation: a new disease entity? European Journal of Gastroenterology & Hepatology. 2010;22(3):378.*
23. [^]Abazarfard Z, Eslamian G, Salehi M, Keshavarzi S. *A Randomized Controlled Trial of the Effects of an Almond-enriched, Hypocaloric Diet on Liver Function Tests in Overweight/Obese Women. Iranian Red Crescent medical journal. 2016;18(3):e23628.*
24. [^]Rafiq N, Younossi ZM. *Effects of weight loss on nonalcoholic fatty liver disease. Seminars in liver disease. 2008;28(4):427-33.*
25. [^]van der Windt DJ, Sud V, Zhang H, Tsung A, Huang H. *The Effects of Physical Exercise on Fatty Liver Disease. Gene Expr. 2018;18(2):89-101.*
26. [^]Nyblom H, Berggren U, Balldin J, Olsson R. *High AST/ALT ratio may indicate advanced alcoholic liver disease rather than heavy drinking. Alcohol and alcoholism (Oxford, Oxfordshire). 2004;39(4):336-9.*
27. [^]Vech RL, Lumeng L, Li TK. *Vitamin B6 metabolism in chronic alcohol abuse The effect of ethanol oxidation on hepatic pyridoxal 5'-phosphate metabolism. J Clin Invest. 1975;55(5):1026-32.*
28. [^]Ueda T, Kawakami R, Horii M, Sugawara Y, Matsumoto T, Okada S, et al. *High mean corpuscular volume is a new indicator of prognosis in acute decompensated heart failure. Circulation Journal. 2013;77(11):2766-71.*
29. [^]Loprinzi PD, Mahoney SE. *Association between flavonoid-rich fruit and vegetable consumption and total serum bilirubin. Angiology. 2015;66(3):286-90.*
30. [^]Swift DL, Johannsen NM, Earnest CP, Blair SN, Church TS. *Effect of different doses of aerobic exercise training on total bilirubin levels. Medicine and science in sports and exercise. 2012;44(4):569-74.*
31. [^]World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *The Journal of the American College of Dentists. 2014;81(3):14-8.*