

vi: 27 December 2024

Research Article

# Row Configuration and Soil Amendment Affect Performance of Popcorn (*Zea mays* L. var. *Everta*) in Semi-Arid South Africa

Peer-approved: 27 December 2024

© The Author(s) 2024. This is an Open Access article under the CC BY 4.0 license.

Qeios, Vol. 6 (2024)  
ISSN: 2632-3834

Oyeyemi Adigun Dada<sup>1</sup>, Funso R. Kutu<sup>2</sup>, Sydney Mavengahama<sup>1</sup>

1. Food Security and Safety Focus Area, North-West University, Potchefstroom, South Africa; 2. School of Agricultural Sciences, University of Mpumalanga, Nelspruit, South Africa

Appropriate plant spacing is vital to maximizing marginal field and expediting equitable distribution of applied plant nutrients for improved popcorn yield. Two field experiments were carried out during the summer seasons of 2017/2018 and 2018/2019 at the North-West University (NWU) research farm, Mafikeng campus. The study investigated the response of popcorn to different rates of compost and NPK 20-7-3 fertilizer and plant spacing in the semi-arid region of South Africa. The trial consisted of twenty treatments laid out as a split-plot in a randomized complete block design ( $r=3$ ). The main plot and subplot effects were amendment rates (4 and 8 t/ha compost, 90 and 180 kg N/ha, while the unamended field served as the control); and four intra-row spacing (cm): 15×15 (SP<sub>1</sub>), 20×20 (SP<sub>2</sub>), 25×25 (SP<sub>3</sub>), and 30×30 (SP<sub>4</sub>). Data were collected on growth and yield components. Results showed that popcorn had the highest number of leaves (12.75) in plots fertilized with 8 t/ha compost under SP<sub>4</sub>, while the tallest plant (205.64 cm) was recorded in plots intra-spaced at SP<sub>2</sub> and fertilized with 8 t/ha compost. The leaf area index was highest (5.1) in plots amended with 90 kg N/ha under SP<sub>4</sub>. The chlorophyll content of popcorn in plots supplied with 90 kg N/ha under SP<sub>3</sub> was significantly higher, with 56.1% more than the leaf chlorophyll from unfertilized plots under SP<sub>1</sub>. Biomass (178.33 g/plant) and ear number (2.08) were highest in plots treated with 180 kg N/ha under SP<sub>3</sub>. The kernel yield of 3.28 t/ha and harvest index of 0.32 were lowest in unfertilized plots under SP<sub>1</sub>. Popcorn yield improved in plots amended with 8 t/ha compost at SP<sub>2</sub> plant spacing, similar to the observations in plant configured at SP<sub>2</sub> and fertilized with 90 kg N/ha mineral fertilizer. Nevertheless, the provision of balanced nutrients and the eco-friendliness of applying organic fertilizer favoured the preference for the use of compost for promoting increased popcorn production.

Corresponding author: Oyeyemi A. Dada,  
[oadada24.7@yahoo.com](mailto:oadada24.7@yahoo.com)

# 1. Introduction

Popcorn (*Zea mays* L. *everta*) is an annual cereal crop of economic importance cultivated for its kernels that pop, puff up, or expand in size when slightly heated<sup>[1]</sup>. This attribute distinguishes it from other varieties of corn. The popped kernels are consumed as snacks in many households and recreational parks for their low carbohydrate and nutritional benefits<sup>[2]</sup>. There has been tremendous growth in the popcorn industry around the world, particularly in the United States of America<sup>[3]</sup> and South Africa, where it is projected to increase by 0.7% above the current net worth of ZAR 174 million by 2024<sup>[4]</sup>. Most popcorn industries in South Africa rely on imported kernels to complement the little harvested from about 2000 ha of local cultivation<sup>[5]</sup>. The major constraint to popcorn production in South Africa is the paucity of information on appropriate agronomic practices, perhaps because the crop is not native to the country. Most importantly, relevant information on its nitrogen requirement and the appropriate density required per unit of land area for its cultivation is not well documented.

The value of popcorn in improving food and nutrition security cannot be overemphasized. Kumar and Prabhaskar<sup>[6]</sup> and Rebello et al.<sup>[7]</sup> reported that it is an important snack that supplies high-quality nutrients like fiber, low cholesterol<sup>[8]</sup>, low carbohydrates<sup>[9]</sup> and quality proteins. It is a whole grain that supplies essential minerals and is very rich in antioxidants<sup>[10]</sup>, aids digestion<sup>[11]</sup>, enhances weight loss, and prevents diseases<sup>[2]</sup>. Besides, it commands triple the market price of other maize cultigens<sup>[12]</sup>. Notwithstanding these benefits, many popcorn growers lack basic information on its agronomic characteristics, especially in semi-arid regions where the grains are highly consumed. Information on the cultivation of popcorn in South Africa is still limited to a few provinces in spite of the fact that the crop can be cultivated in all the maize-growing regions, while cultivation is generally low compared to its market demand<sup>[13]</sup>. An estimated 55,000 t/year was harvested on 10,000 ha, and the crop was grown principally in the Free State and Northern Cape regions<sup>[13]</sup>. The annual popcorn produced in South Africa is approximately 1.8% of the total corn harvested on 3.1 million ha/annum<sup>[14][15]</sup>. One of the major limiting constraints to popcorn cultivation in South Africa is inadequate information

on the agronomic characteristics of this crop. This is a common phenomenon in the sub-Saharan agro-ecosystem, particularly with popcorn<sup>[16]</sup>.

Crops perform optimally under favourable environmental conditions such as adequate nutrition, moisture availability, and appropriate spacing with negligible or minimal intra- or inter-competition<sup>[17]</sup>. As a major agronomic practice, crops are supplied with either organic or mineral fertilizer to augment the native nutrients, which in most cases are inadequate to support the crop all through the statutory phenological stages<sup>[18]</sup>. This inadequacy may be a result of soil characteristics or depletion/removal of soil mineral nutrients by previous crops. Singh and Ryan<sup>[19]</sup> and Badu-Apraku and Fakorede<sup>[16]</sup> reported that in a soil-plant system, the response of maize to applied materials varies depending on several factors.

Utilization of diverse types and forms of organic fertilizers as an alternative to mineral fertilizers is gaining more attention and relevance because of its low cost, positive environmental impact, plant nutrition security, residual effect, and good mineralization values<sup>[20]</sup>. These views have been buttressed by Ghorbani et al.<sup>[21]</sup> on tomato, Matallana Gonzalez et al.<sup>[22]</sup> on vegetables, Zhao et al.<sup>[23]</sup> on rice, and Cihangir and Oktem<sup>[24]</sup> on popcorn. However, the derived benefits of the application of organic fertilizer are often location-specific, based on crop genetic potentials as well as other biotic and abiotic influences according to Waldrip et al.<sup>[25]</sup>.

Generally, average N recommendations for growing corn in South Africa ranged from 55 to 180 kg N/ha<sup>[26][27][28]</sup>. Application of farmyard manure for large-scale corn production is rare in South Africa, but few smallholders use organic manures like chicken or cattle manure to fertilize their arable fields. Information is sketchy on standard or recommended quantities of compost or farmyard manure for growing popcorn in South Africa. Information is scant regarding the response of popcorn to either compost or mineral fertilizer application in the tropical semi-arid region. The need to evolve appropriate rates of organic as well as mineral fertilizers that would be adequate for growing popcorn in this region is very pertinent for improving its production.

Uniformity of the crop in the field is premised on appropriate spacing, which plays an important role in optimizing solar utilization efficiency<sup>[29]</sup> and equal

distribution of applied or available resources. High photosynthetic efficiency is a function of adequate plant architectural pattern and/or spatial arrangement. According to Brodrick et al. (2013)<sup>[30]</sup> and Dong et al. (2010)<sup>[31]</sup>, inadequate architectural arrangement affects biomass accumulation which ultimately impinges photosynthate assimilation and partitioning negatively. The lesser the intra or inter competition within crops, the better the photoassimilate that is partitioned into economic yield. According to Dai et al. (2015)<sup>[32]</sup>, manipulation of spatial crop arrangement is a common agronomic practice engaged to improve physiological performance of crops.

Furthermore, the cultivable land is becoming marginal by the day owing to competition from different fronts like the proliferation of industries, infrastructural needs, and residential development<sup>[33]</sup><sup>[34]</sup>. This suggests the need for effective utilization of the available agricultural land for cultivating crops of high economic value necessary for human and livestock survival. While doing this, it is expedient to apply scientific approaches necessary to explore the potentials of all the inputs to maximize the economic gain from the process. There is a dearth of information on the appropriate mineral fertilizer or compost rate, as well as the plant density that would be adequate for enhancing the performance of popcorn with respect to growth and yield in the dry land regions of South Africa. The hypothesis of this study was that the grain yield of popcorn is affected by soil amendments and row configuration. Therefore, we investigated the influence of intra-row spacing, compost, and NPK 20-7-3 fertilizer rates on the growth, kernel yield, yield components, and biomass accumulation by *Zea mays* L. *Everta* in the tropical semi-arid region of South Africa.

## 2. Materials and methods

### *Description of the experimental site*

The experiment was conducted at the experimental farm of North West University (NWU), Mafikeng Campus (25°49'39''S, 25°36'3''E, 1280 m above sea level) during the 2017/2018 and 2018/2019 summer planting seasons. The farm is located in a savanna semi-arid climate in the North West Province of South Africa. The Province receives approximately 350–400 mm of rainfall annually in winter and averages 250–300 mm during the summer season, with an average temperature range of 18 °C – 22 °C in winter and 22 °C

– 37 °C during summer. The relative humidity varied from 13.5–81%.

### *Pre-trial soil samples*

Pre-trial soil samples collected at a depth of 0–30 cm and compost were analysed at the Department of Crop Science laboratory using LECO CNS TruMac analyzer for total carbon, nitrogen and sulphur contents. The P concentration was determined using Olsen's method as described by Koralage et al. (2015)<sup>[35]</sup>. This was achieved by extracting phosphorus from the soil sample with an acid solution and reacting it with a colorimetric reagent. The absorbance of the resulting solution was measured using spectrophotometer, and the phosphorus content was determined using a calibration curve. The K, was determined by atomic absorption spectrometry. The pH (Soil: H<sub>2</sub>O, 1:1) of the compost and soil samples were determined using a pH meter (Cole-Parmer Digi-Sense® Model No. 5938-00), while the percent particle size distribution analyses were determined using modified Bouyoucos methods described by Beretta et al. (2014)<sup>[36]</sup>. The soil was clay loam with pH of 6.8; total N of 6.96 mg/kg; Bray-1 P was 80.0 mg/kg and K was 235.0 mg/kg.

### *Treatments and experimental design*

The study was a 4 × 5 factorial comprising four intra-row spacings and five fertilizer treatments planted during 2017/18 and 2018/2019–summer growing seasons. The trial was laid out as a split plot with each treatment arranged in a Randomized Complete Block Design (RCBD) and replicated three times. The main plot contained fertilizer treatment, while the different within row spacings was the subplot effect. The four intra-row (within plants) spacings were 15 cm × 15 cm (SP<sub>1</sub>), 20 cm × 20 cm (SP<sub>2</sub>), 25 cm × 25 cm (SP<sub>3</sub>) and 30 cm × 30 cm (SP<sub>4</sub>), while the inter row (between plants) was spaced at 70 cm apart. The five fertilizer treatments consisted of 90 kg N/ha and 180 kg N/ha, while the compost was applied at the rates of 4 t/ha and 8 t/ha with unfertilized plot served as the control. The NPK fertilizer used contained (kg/100 kg); N = 20, P = 7, K = 3, Zn = 0.5, S = 5, Ca = 1. The compost was prepared from, sorted municipal solid waste at NWU Farm using the modified heap method described by Karak et al. (2018)<sup>[37]</sup>. The applied compost had 36 g/kg nitrogen, 1.41 g/kg phosphorus, 10.22 g/kg potassium and 134 g/kg organic carbon. The equivalent plant population per plot of 3 m × 2.1 m size with four rows each was 84, 64, 52 and 44 plants,

representing 1333,333; 101,587; 82,540 and 69,841 plants/ha for SP<sub>1</sub>, SP<sub>2</sub>, SP<sub>3</sub> and SP<sub>4</sub>, respectively.

### *Land preparation, sowing, and field management*

The experimental field was ploughed and harrowed using disc plough and harrow with working depth of 25-30 cm. Thereafter, the field was marked and demarcated based on the experimental design to accommodate the treatments and blocks. The dimension of the entire experimental field was 25.5 m × 24 m, while each experimental unit (plot) had a dimension of 3 m × 2.1 m. Mineral fertilizer was applied at planting, while organic fertilizer (compost) was incorporated into the soil two weeks before sowing. Compost was applied two weeks before sowing to enhance mineralization and ease mineral uptake. Seeds of mid-altitude popcorn, obtained from the International Institute of Tropical Agriculture (IITA, Nigeria), were sown in December 2017 for the first trial and in December 2018 for the repeat experiment. One seed was sown per hole at a depth of 2 cm. The field was kept weed-free throughout the experimental period through manual hand hoeing with a Dutch hoe. The trial was rainfed; however, supplementary irrigation was applied when the soil moisture level was low, especially at the tasseling and silking stages, using visual soil examination and evaluation methods of<sup>[38]</sup>. The overhead sprinkler method was adopted for irrigation purposes.

### *Data collection*

Four plants from the middle rows of each plot were randomly selected and tagged for data collection in each plot. Data collection followed standard procedures for determining the number of leaves and plant height. The Leaf Area Index (LAI) was evaluated following the procedure of Fageria et al.<sup>[39]</sup> and Berdjour et al.<sup>[40]</sup> as  $LAI = (P \times L \times A)/(GA)$ .

Where: LAI = Leaf area index, P = Plant population/ground area (ha), L = Number of fully expanded green leaves/plant, A = Single leaf area (cm<sup>2</sup>), and GA = Ground area (m<sup>2</sup>).

The stem diameter was measured with a vernier caliper (Mastercraft GS5071522). Chlorophyll content was collected *in situ* from fully expanded leaves per plant with a hand-held chlorophyll meter (model CCM-200 plus). Yield and its components: ear number, ear mass per plant, and 1000 seeds mass were collected following standard procedures

described by Abebe and Feyisa<sup>[41]</sup>. The harvest index was evaluated as:

$$\text{Harvest index} = \frac{\text{Mass of grain yield (g)}}{\text{Biological biomass (g)}} \quad [41].$$

The kernel yield was determined from the 1.4 m<sup>2</sup> area at harvest maturity. Grain yield was evaluated by harvesting ears from a 1.4 m<sup>2</sup> area of the plot. The period of data collection covered the vegetative (V6), tasseling (VT/R5), and maturity (R6) growth stages in corn described by Lee<sup>[42]</sup>.

### *Statistical analysis*

The data for the two planting seasons were pooled together according to Gomez and Gomez<sup>[43]</sup> and analysed using analysis of variance (ANOVA) of the general linear model (GLM) of the Statistical Analysis System (SAS, version 6.0) according to O'Rourke and Hatcher<sup>[44]</sup>. Different means were separated using the least significant difference (LSD) test at  $p \leq 0.05$  following the methods of Gomez and Gomez<sup>[43]</sup>.

## **3. Results**

Number of leaves, diameter of stem, and ear height were not significantly ( $p > 0.05$ ) influenced by within-row plant spacing. However, height, LAI, chlorophyll content, ear number, ear mass/plant, and biomass were significantly affected by intra-row spacing (Table 1). The rates of fertilizer applied had a significant ( $p < 0.05$ ) effect on height, leaf area index, chlorophyll content, and ear height, as shown in Table 1. The interaction between soil amendments and inter-row spacing had a significant effect on the number of leaves, leaf area index, chlorophyll content, and number of ears.

Mean squares								
Sources of variation	Df	NL	PHT	LAI	DS	CHL	EH	NE
Spacing (A)	3	51.67	3892.31*	1.94*	20.58	711.50*	470.65	3.43*
Amendment (B)	4	214.71	3270.40*	0.92*	75.47	1641.85*	2219.19*	0.77
A × B	12	7.29*	1872.88	2.62*	26.46	658.25*	649.67	2.13*
CV (%)		17.86	20.50	31.95	94.17	43.80	30.74	39.79
Mean		11.48	184.80	69.35	14.19	33.71	83.82	1.64
	Df	Ear/plant (g)	Ear/plot (Kg)	Cob (g)	1000 seeds (g)	Grain yield (t/ha)	Biomass (g)	Harvest index
Spacing (A)	3	15901.79*	0.52	961.69	272590.30	42.70	24062.58*	0.12
Amendment (B)	4	5275.51	1.85	639.55	237613.20	36.03	8103.74	0.13
A × B	12	41531.10	4.63	684.16	55113.08	7.03	4901.38	0.09
CV (%)		38.55	72.16	65.20	20.68	58.13	55.08	122.15
Mean		160.55	1.45	46.46	1580.60	5.10	129.70	0.47

**Table 1.** Mean squares, coefficient of variation and means of growth and yield characteristics of popcorn as influenced by intra-row configuration and soil amendments.


\*Significant at  $p \leq 0.05$ ; Df = degree of freedom; NL= number of leaves, PHT = plant height, LAI= leaf area index, DS = diameter of stem, CHL=chlorophyll content, EH= ear height, NE = number of ears.

### Growth response of popcorn to different intra-row configuration

Different intra-row spacings did not affect leaf formation and height of popcorn plants significantly ( $p > 0.05$ ) both at vegetative (V6) and tasseling (VT/R5) growth stages. However, these growth parameters were significantly ( $p \leq 0.05$ ) affected by intra-row spacing at the physiological maturity stage (R6). The number of leaves produced at the narrow spacing of 15 cm x 15 cm was significantly lower than the number of leaves recorded in plants with other spacings (Figure 1A). On the other hand, the height of the crop was not affected by intra-row spacing at the V6 and VT/R5 stages. Nevertheless, at R6, the height of the crop was significantly taller at the 20 cm x 20 cm spacing but not significantly taller than plants spaced at 25 cm x 25 cm or 30 cm x 30 cm, as indicated in Figure 1B. The leaf area index (LAI) was statistically affected by intra-row spacing throughout the growth stages

(Figure 1C). The LAI was significantly higher under the narrow intra-spacing (15 cm x 15 cm) than under the other intra-row spacings. The widest intra-row spacing (30 cm x 30 cm) had the lowest LAI.

The diameter of popcorn stems was not affected by different intra-row spacings at the vegetative and tasseling stages, except at the physiological maturity period, when the diameter was highest (18.40 mm) in plants spaced at 25 cm x 25 cm, as shown in Figure 1D. Intra-row spacing exerted a significant effect on leaf chlorophyll concentration at varying phenological stages. The highest concentration of leaf chlorophyll was obtained in plants intra-spaced at 25 cm x 25 cm at the vegetative phase. The lowest chlorophyll content was, however, recorded in plants intra-spaced at 15 cm x 15 cm at the tasseling and maturity phases (Figure 1E). The ear height was significantly affected by intra-row spacing, as plants spaced at 20 cm x 20 cm recorded the tallest ear height of 87.61 cm relative to other spacing treatments (Figure 1F).




**Figure 1.** Effect of intra-row configuration on the growth of popcorn.

### Growth response of popcorn to different rates of soil amendments

Different rates of soil amendments had a significant influence on the growth of popcorn. The effects of the two types of soil amendments on all measured growth parameters, except stem diameter, were statistically comparable relative to the control. The effect of organic and nitrogen fertilizers was significantly similar on the number of leaves produced across the growth stages. Leaf formation improved significantly at the tasseling and physiological maturity stages in plots supplied with 8 t/ha of compost, but this was not statistically different from the number of leaves formed in plots supplied with different rates of NPK 20-7-3 fertilizer (Figure 2A). The height of the crop increased significantly at the vegetative stage in plots fertilized with 90 kg N/ha (NPK 20-7-3) compared to the unfertilized plot (Figure 2B). At the tasseling stage, the height was not influenced by soil amendments, but at physiological maturity, the crop had the tallest plants in plots supplied with 8 t/ha of compost over and above other rates of either organic or inorganic fertilizers. At the early growth stage, soil amendments had no effect on LAI. However, over time, LAI was significantly affected by different rates of amendments, with the highest LAI recorded in plots supplied with 90 kg N/ha, which was significantly similar to the LAI recorded in fields fertilized with 4 t/ha of compost at tasseling. At physiological maturity, LAI was highest in plots supplied with mineral fertilizer at the rate of 90 kg

N/ha relative to the LAI recorded in the unfertilized field, which recorded the lowest LAI (Figure 2c).

The stem diameter was not affected by varying rates of organic or inorganic fertilizer (Figure 2d). Rates of soil amendments influenced chlorophyll content in popcorn plants. The plants had the lowest chlorophyll concentration on field fertilized with 4 t/ha compost which was statistically similar to the concentration obtained in unfertilized the field. There was no significant difference in chlorophyll concentration of popcorn grown in plots supplied with 8 t/ha of compost, 90, or 180 kg N/ha (Figure 2e). The tallest ear height (91.66 cm) was recorded in the field amended with 8 t/ha of compost, which was not significantly different from the ear height recorded in the field amended with 90 kg N/ha (Figure 2f).



**Figure 2.** Effect of different rates of soil amendments on the growth of popcorn.

### Effects of intra-row configuration and soil amendment rates on the growth of popcorn

The effects of interaction between intra-row spacing and soil amendments showed that the application of 90 kg N/ha enhanced the number of leaves produced by popcorn sown at 15 cm × 15 cm at the vegetative and tasseling stages (Table 2). Although this was not significantly higher than the number of leaves recorded in the other treatments, except in the unfertilized plots with narrower spacing, where the number of leaves produced was significantly lower compared to other treatments. However, at maturity, the plant formed the highest number of leaves in plots supplied with 8 t/ha of compost and intra-spaced at 30 cm × 30 cm.

The crop had the tallest plants in plots supplied with 8 t/ha and intra-spaced at 20 cm × 20 cm, both at tasseling and maturity. The shortest plants were recorded under the narrower spacing of 15 cm x 15 cm in the unamended field (Table 2). The effect of soil amendments and intra-row spacings on LAI varies across the three phenological phases. At the vegetative stage, plots fertilized with 180 kg N/ha and intra-spaced at 20 cm × 20 cm had the highest LAI of 2.08. At the tasseling stage, the narrow intra-spacing (15 cm × 15 cm) and lower compost rate (4 t/ha) promoted the highest LAI. However, at maturity, plots fertilized with 180 kg N/ha and intra-spaced at 15 cm × 15 cm had an LAI of 7.04, which was significantly higher than the LAI (2.76) obtained in unfertilized plots under 25 cm x 25 cm intra-row spacing.

Interactions		Number of leaves			Plant height (cm)			Leaf area index		
Amendments	Spacing	Veg.	Tasl.	Mat.	Veg.	Tasl.	Mat.	Veg.	Tasl.	Mat.
Control	SP1	6.50bc	11.00dc	10.33de	32.71c	107.64b	171.30c-e	1.07ab	5.13c-f	4.63d-h
	SP2	6.08c	11.00dc	11.00b-e	32.71c	120.07b	186.46a-d	0.45b	3.17gh	3.22i-k
	SP3	6.42bc	10.92d	10.17e	33.77c	119.23b	174.48b-e	0.68b	2.86gh	2.76k
	SP4	6.67abc	11.17b-d	10.83b-e	33.32c	114.26b	182.33a-e	0.50b	3.27f-h	3.66g-k
4 t/ha compost	SP1	7.58a	11.67a-d	10.92b-e	38.61a-c	129.36b	181.08a-e	1.05ab	7.62a	6.68ab
	SP2	7.08abc	11.50a-d	11.08-e	41.23a-c	133.91b	187.84a-d	0.52b	4.45c-g	5.65b-e
	SP3	6.92abc	12.00a-d	11.92a-d	36.03bc	131.33b	185.76a-d	0.92ab	3.53e-h	3.70g-k
	SP4	7.17ab	11.83a-d	12.00a-c	34.95bc	130.21b	195.87a-c	0.33b	2.33h	3.12i-k
8 t/ha compost	SP1	7.33ab	11.58a-d	11.08b-e	33.77a-c	144.23b	188.18a-d	0.87ab	5.31b-e	5.87a-d
	SP2	7.25ab	12.00a-d	11.83a-e	35.71bc	153.06a	205.64a	0.64b	5.32b-e	5.34c-f
	SP3	7.00abc	11.58a-d	12.33ab	33.32a-c	137.44b	202.21ab	0.69b	4.10d-h	4.62d-h
	SP4	7.08abc	12.17a-d	12.75a	34.37a-c	137.92b	190.68a-c	0.40b	2.96gh	3.55g-k
90 kg N/ha (NPK)	SP1	7.67a	12.50a	11.50a-e	36.03a-c	149.22b	182.27a-e	0.98ab	7.03ab	6.50a-c
	SP2	7.33ab	11.67a-d	11.08b-e	38.61ab	147.88b	193.58a-c	0.74b	5.24b-e	6.04a-c
	SP3	6.67abc	11.00dc	12.42ab	35.26bc	110.85b	157.78de	0.46b	4.36c-g	4.70d-g
	SP4	7.17ab	11.17b-d	11.83a-d	35.71a-c	133.78b	193.75a-c	0.67b	3.23gh	3.17i-k
180 kg N/ha (NPK)	SP1	6.33bc	11.67a-d	10.08e	34.95a-c	140.62b	152.01e	1.16ab	5.60b-d	7.04a
	SP2	7.17ab	12.00a-d	12.25ab	34.38a-c	145.10b	188.16a-d	2.08a	4.41c-g	4.32f-j
	SP3	7.08abc	11.42a-d	11.00b-e	41.23a	139.32b	170.37c-e	0.95ab	3.59e-h	3.41g-k
	SP4	6.67abc	11.25a-d	12.08a-c	34.37c	139.83b	185.32a-d	0.26b	2.70gh	3.03jk
LSD ( $p \leq 0.05$ )		1.07	1.27	1.65	8.98	71.79	30.46	1.35	0.78	1.32

**Table 2.** Effect interactions between intra-row configuration and different rates of soil amendments on number of leaves, plant height and leaf area index of popcorn.

LSD = Least significant difference at  $p < 0.05$ , SP1 = 15 cm × 15 cm; SP2 = 20 cm × 20 cm; SP3 = 25 cm × 25 cm; SP4 = 30 cm × 30 cm, Veg. = vegetative; Tasl. = tasseling and Mat. = physiological maturity. Values with different letters are not significantly different at  $p < 0.05$  using LSD.

The interaction between spacing and fertilizer application had no effect on stem diameter as shown

in Table 3. Popcorn plants intra-spaced at 25 cm × 25 cm and 20 cm × 20 cm and fertilized with 8 t/ha compost had highest chlorophyll content (25.48  $\mu\text{mol}/\text{m}^2$  and 42.92  $\mu\text{mol}/\text{m}^2$ ) at vegetative and tasseling phases, respectively. At maturity however, popcorn had highest chlorophyll content of 47.03  $\mu\text{mol}/\text{m}^2$  in plots where plants were intra-spaced at



25 cm × 25 cm and fertilized with 90 kg N/ha which statistically similar to chlorophyll content recorded in plant intra-spaced at 30 cm × 30 cm and fertilized with 180 kg N/ha (Table 3). Plant had lowest chlorophyll content in unamended plots with plants intra-spaced at 15 cm × 15 cm. Plants intra-spaced at

20 cm × 20 cm and amended with 90 kg N/ha had highest ear height (93.98 cm) which was statistically similar to ear height recorded in plots intra-spaced at 25 cm × 25 cm or 20 cm × 20 cm and amended with 8 t/ha (Table 3).

Interaction	Spacing	Stem diameter (mm)			Chlorophyll content ( $\mu\text{mol/m}$ )			Ear height (cm)
		Veg.	Tasl.	Mat.	Veg.	Tasl.	Mat.	
Amendments	Spacing							
Control	SP1	3.11b	10.71	10.66	13.13d	24.23g	20.66f	72.67ab
	SP2	4.49ab	11.80	11.02	16.84b-d	28.15de-g	30.68b-f	81.40ab
	SP3	4.35ab	11.66	11.94	12.33d	36.14d-g	29.46c-f	70.76b
	SP4	3.57b	13.15	14.30	15.23cd	24.69c-g	33.47b-d	76.84ab
4 t/ha compost	SP1	4.91ab	12.99	13.13	15.52cd	24.69fg	23.73d-f	84.50ab
	SP2	3.65b	11.33	14.28	14.93cd	28.02b-f	28.97c-f	83.91ab
	SP3	5.88ab	8.38	12.86	17.82a-d	30.35d-g	20.20f	81.89ab
	SP4	5.02ab	12.73	13.67	18.36a-d	26.01fg	29.56c-f	84.37ab
8 t/ha compost	SP1	3.91ab	13.06	15.32	21.52a-c	33.72d-g	37.43a-c	90.08ab
	SP2	7.77ab	12.84	14.55	13.41d	42.92a	37.30a-c	93.85a
	SP3	4.74ab	11.88	15.78	25.48a	29.03d-g	39.85a-c	93.88a
	SP4	4.07ab	12.95	15.00	16.42cd	26.01b-g	36.81a-c	88.83ab
90 kg N/ha (NPK)	SP1	3.79ab	13.13	14.61	16.00cd	30.35b-d	33.66b-d	86.48ab
	SP2	4.28ab	14.61	13.96	14.06cd	33.72a-c	36.05a-c	93.98a
	SP3	4.22ab	15.17	16.57	14.86cd	29.03b-e	47.03a	93.90a
	SP4	5.65ab	14.37	15.68	15.30cd	28.15b-f	39.15a-c	79.56ab
180 kg N/ha (NPK)	SP1	4.66ab	13.94	13.70	16.21cd	36.14ab	42.49ab	71.47b
	SP2	4.08ab	13.88	14.56	18.86a-d	28.02b-f	30.20c-f	82.46ab
	SP3	7.77ab	15.54	16.17	21.43a-c	36.14ab	39.74a-c	73.49ab
	SP4	3.28b	12.93	14.95	18.93a-d	24.23c-g	45.90a	76.42ab
LSD ( $p \leq 0.05$ )		4.97	9.23	18.74	7.74	18.01	11.87	20.27
		ns	ns	ns	*	*	*	*

**Table 3.** Effect of intra-row configuration, compost and nitrogen fertilizer and their interactions on stem diameter, chlorophyll content and ear height of popcorn.

LSD = Least significant difference at  $p < 0.05$ , SP1 = 15 cm  $\times$  15 cm; SP2 = 20 cm  $\times$  20 cm; SP3 = 25 cm  $\times$  25 cm; SP4 = 30 cm  $\times$  30 cm, Veg. = vegetative; Tasl. = tasseling and Mat. = physiological maturity. Values with different letters are not significantly different at  $p \leq 0.05$  using LSD, ns = not significant.

### *Yield response popcorn to different inter-row configuration*

Mass of ear per plot, mass of cob and harvest index were not significantly affected by intra-row spacing. Nevertheless, highest ear mass per plant and 1000 seed mass were obtained in plots intra-spaced at 30 cm × 30 cm and this was significantly similar to ear or 1000 seed mass obtained in plot intra-spaced at either 20 cm × 20 cm or 25 cm × 25 cm (Table 4).

The grain yield was significantly influenced by intra-row spacing as highest grain yield was obtained in plots intra spaced at 20 cm × 20 cm. Also, biomass yield was highest in plot intra-spaced at 25 cm × 25 cm but not significantly higher than from biomass obtained in plots intra-spaced at 30 cm × 30 cm (Table 4). Plant intra-spaced at 30 cm × 30 cm had highest number of ears (1.85) which was significantly higher than number of ears from plants intra-spaced at 15 cm × 15 cm or 20 cm × 20 cm as shown in Table 4.

### *Yield response of popcorn to different rates of soil amendments*

It was observed that highest ear mass (195.46 g) per plant was obtained in plots supplied with 90 kg N/ha. Similarly, plots fertilized with 8 t/ha compost produced ear with highest mass (1.82 kg) per plot, 1000 seed mass (1,695.54 g) and total biomass (146.95

g), but these were not significantly higher than the values obtained in plots fertilized with other rates of organic or inorganic fertilizers. The grain yield was not significantly affected by fertilizer application, but highest grain yield (6.12 t/ha) was obtained on plots amended with 90 kg N/ha. Nevertheless, lowest kernel yield was obtained in the unfertilized field (Table 4). Number of ears per plant was significantly higher in field amended with 180 kg N/ha than number of ears recorded on plots amended with 4 t/ha compost and unamended plots.

### *Effect of inter-row configuration and soil amendment rates on yield components of popcorn*

Interactions between intra-row spacing and soil amendments showed that ear mass (2.23 kg/plot) were highest in plots fertilized with 8 t/ha compost and intra-spaced at 20 cm × 20 cm. The results indicated that 1000 seeds mass was highest in plants intra-spaced at 30 cm × 30 cm and amended with 4 t/ha compost. The kernel yield (8.18 t/ha) and harvest index (0.91) were highest in plants intra-spaced at 20 cm × 20 cm and fertilized with 90 kg N/ha (Table 4). Highest biomass and number of ears were highest on field intra-spaced at 25 cm × 25 cm and fertilized with 180 kg N/ha as shown in Table 4.

		Weight					Ear number	Harvest index	
		Ear/plant (g)	Ear/ plot (kg)	Cob (g)	1000 seeds (g)	Grain Yield (t/ha)			Biomass (g)
<b>Intra spacing</b>									
15 cm × 15 cm		131.85b	1.97	38.96	1485.31b	5.31ab	114.26bc	1.36c	0.34
20 cm × 20 cm		159.11ab	2.81	47.23	1537.54ab	5.80a	103.49c	1.58b	0.54
25 cm × 25 cm		178.81a	2.32	49.19	1616.55ab	4.97ab	160.63a	1.78ab	0.56
30 cm × 30 cm		180.41a	1.89	50.43	1683.01a	4.30b	140.21ab	1.85a	0.44
LSD (p≤0.05)		28.85	1.28	14.12	1.52	1.38	33.31	0.21	0.27
			ns	ns				ns	ns
<b>Amendments</b>									
Control		148.20b	1.30ab	41.14	1515.58ab	4.53	116.48ab	1.52b	0.46
4 t/ha compost		148.31b	1.57ab	51.20	1632.06a	4.95	122.39ab	1.54b	0.48
8 t/ha compost		185.18a	1.82a	51.02	1695.54a	5.14	146.95a	1.73ab	0.40
90 kg NPK		195.46a	1.39ab	46.85	1537.57ab	6.12	103.95b	1.60ab	0.58
180 kg NPK		149.31b	1.02b	38.45	1421.81bc	4.78	137.39ab	1.85a	0.34
LSD (p≤0.05)		35.34	0.60	17.30	186.66	1.69	40.79	0.26	0.33
				ns		ns			ns
<b>Amendments</b>									
Control	SP1	121.60cd	1.03ab	32.02	1379.70b-d	3.28b	97.25c-f	1.08f	0.32ab
	SP2	146.33a-d	1.67ab	38.16	1486.30a-d	5.39ab	107.74b-f	1.33d-f	0.54ab
	SP3	153.87a-d	1.26ab	52.05	1611.10a-d	5.00ab	107.33b-f	1.50b-f	0.45ab
	SP4	170.44a-d	1.22ab	46.23	1585.30a-d	3.39b	156.04a-e	1.75a-e	0.43ab
4 t/ha compost	SP1	122.36cd	1.84ab	42.86	1473.20a-d	5.27ab	102.70b-f	1.42c-f	0.42ab
	SP2	144.92a-d	1.57ab	56.55	1607.90a-d	4.16b	90.95e-f	1.50b-f	0.53ab
	SP3	148.03a-d	1.61ab	52.24	1699.10a-c	4.93ab	176.98a-c	1.42c-f	0.57ab
	SP4	177.94a-d	1.29ab	53.16	1748.10a	3.76b	118.92b-f	1.83a-d	0.41ab
8 t/ha compost	SP1	166.82a-d	1.60ab	45.69	1702.50a-c	4.48b	165.39a-e	1.25ef	0.34ab
	SP2	211.90a	2.23a	52.48	1742.70ab	5.94ab	104.16b-f	1.58a-f	0.43ab
	SP3	190.72a-c	2.05a	54.48	1638.30a-d	5.99ab	168.53a-d	1.92a-c	0.62ab
	SP4	171.29a-d	1.39ab	51.41	1698.80a-c	4.15b	149.72a-f	1.67a-e	0.57ab
90 kg N/ha (NPK)	SP1	135.94b-d	1.56ab	39.28	1514.10a-d	6.51ab	98.74c-f	1.50b-f	0.34ab
	SP2	155.39a-d	1.47ab	45.00	1428.80a-d	8.18a	99.96b-f	1.42c-f	0.91a
	SP3	173.14a-d	1.36ab	52.52	1686.00a-c	4.72b	118.15b-f	2.00ab	0.60ab

		Weight						Ear number	Harvest index
		Ear/plant (g)	Ear/ plot (kg)	Cob (g)	1000 seeds (g)	Grain Yield (t/ha)	Biomass (g)		
	SP4	197.36ab	1.17ab	50.60	1665.50a-d	5.09ab	98.94c-f	2.00ab	0.48ab
180 kg N/ha (NPK)	SP1	115.34d	1.27ab	34.61	1322.60dc	6.32ab	152.09a-e	1.50b-f	0.18b
	SP2	149.23a-d	1.04ab	40.82	1284.60d	4.56ab	84.94ef	1.92a-c	0.30ab
	SP3	164.46a-d	0.64b	31.71	1450.70a-d	3.73b	178.33a	2.08a	0.62ab
	SP4	168.17a-d	1.14ab	46.66	1629.40a-d	4.51b	136.39a-f	1.92a-c	0.32ab
LSD ( $p \leq 0.05$ )		73.67	1.24	36.69	385.58	3.41	81.82	0.52	0.70
			ns	ns					

**Table 4.** Effect of intra-row spacing, compost and nitrogen fertilizers and their interactions on yield and yield components of popcorn.

LSD = Least significant difference at  $p < 0.05$ , SP1 = 15 cm × 15 cm; SP2 = 20 cm × 20 cm; SP3 = 25 cm × 25 cm; SP4 = 30 cm × 30 cm, Veg. – vegetative; Tasl. = tasseling and Mat. = physiological maturity. Values with different letters are not significantly different at  $p \leq 0.05$  using LSD, ns = not significant.

#### 4. Discussion

The poor leaf formation under narrow intra-spacing infers that closer spacing could affect leaf formation negatively. Bernhard and Below<sup>[45]</sup> have also reported similar observations in *Zea mays*. The results obtained in this study are, however, not in consonance with that of Moosavi et al.<sup>[46]</sup> on forage corn in Iran. The better performance observed under wide intra-row spacing implies the absence of competition for basic crop growth resources, particularly nutrients. Widdicombe and Thelen<sup>[47]</sup> and Murányi and Pepo<sup>[48]</sup> have also shown that the height of different maize hybrids was never affected by intra-row spacing. The height of popcorn was not adversely affected by plant configuration, which indicates that intra-row spacings may not play any important role in the tallness of popcorn plants. Nonetheless, closer spacing had been shown by Bernhard and Below<sup>[45]</sup> to improve the height of cereals like sweet corn by Williams II<sup>[49]</sup>. The observation in this trial is in consonance with that of Moosavi et al.<sup>[46]</sup> on corn.

The narrow spacing appeared adequate as it promoted the leaf area index better than other spacings. The leaf area index is a measure of leaf efficiency with respect to intercepting solar radiation per unit area of land. Under narrow spacing with a higher leaf area index, it would suffice to expect an improvement in photoassimilate production over treatments with wider spacing. The superior performance of popcorn with respect to leaf area index observed under narrow intra-row spacing is in consonance with the reports of the Board and Harville<sup>[50]</sup> and Widdicombe and Thelen<sup>[47]</sup> who suggest that rows with closer plant stands had a greater leaf area index compared with the optimal plantings in the short-season maize cultivar. Liu et al.<sup>[51]</sup> also reported that narrow planting configuration influenced light interception and radiation use efficiency in maize. However, the relatively poor performance in intra-row spacing beyond 15 cm × 15 cm could have meant a waste of important production inputs, especially fertilizer, a major input in highly weathered soil with huge monetary cost.

Plants configured in a narrow within-row spacing adversely affected the chlorophyll contents of popcorn. This perhaps could be adduced to a higher population per unit area, as intra-competition could impinge negatively on the concentration of chlorophyll produced. It appeared that chlorophyll synthesis improved with wider intra-row spacing than with narrow spacing. This could mean that wide spacing reduces intra-competition, which ultimately

enhanced chlorophyll synthesis in popcorn. Our observation agrees with that of Golada et al.<sup>[52]</sup>, who obtained the highest chlorophyll content (2.34 mg/g) in baby corn grown at wider spacing. Ear height is a parameter that showed a response to intra-row spacing, as it was affected by within-row plant spacing. This infers that ear height is a major parameter that could be negatively affected by intra-row spacing. The height of the ear has implications for calibrating harvesting machines; hence, the need to reassess the row width of corn is necessary for evolving optimal plant population density for popcorn. The better performance of the number of ears produced under wider within-plant spacing implies that a wider intra space will benefit ear development better than under narrow intra-spacing.

A fairly wide spacing promoted higher photoassimilate production better than narrow spacing, which explains why better grain yield was obtained at the medium spacing of 20 cm × 20 cm. Thus, popcorn appeared to be more efficient at partitioning assimilate into economic yield at wider widths than at somewhat close or narrow intra-spacing. A similar observation had been reported by Sheth et al.<sup>[53]</sup> on sweet potato yield.

It is clear from the trial that either organic or inorganic fertilizer has similar effects on the growth of popcorn; however, the quantity required to produce the expected response is very important. The comparative effect of organic and inorganic fertilizer on the performance of popcorn has been well documented in many arid environments, but there is scant information on such reports in South Africa. Our observations on improving the performance of popcorn with either organic or inorganic fertilizer agree with earlier findings of Laxminarayana et al.<sup>[54]</sup> on sweet potato and Dada et al.<sup>[55]</sup> *Amaranthus cruentus*.

The growth response of popcorn to different rates of soil amendments varies at different growth stages. For instance, the effects of either type of amendment did not show any observable variation at the early stage, possibly because the native nutrients were sufficient to support growth up to that point. It is also unlikely that popcorn is more efficient in fertilizer utilization than other corn genotypes. White et al.<sup>[56]</sup> have shown that popcorn does not demand high nutrients unlike other *Zea* species. This could possibly explain the negligible effect of the supplied soil amendments during the early growth stage. This presupposes that previous fertilization history, crop

requirements, and other factors such as soil inherent characteristics may influence the amount of fertilizer needed by certain crops across different growing phases<sup>[57]</sup>. The diameter of popcorn stems was similar irrespective of variation in fertilizer application. This is in line with different observations by Sener et al.<sup>[58]</sup> on corn and Gözübenli<sup>[59]</sup> on popcorn. The increase in the concentration of chlorophyll with increases in rates of organic or inorganic fertilizers suggests that adequate synthesis of chlorophyll in plants could be linked to the availability of sufficient mineral nutrients, especially nitrogen. A similar observation was recorded in the number of ears produced per plant. This implies that assimilate partitioning of photosynthate into economic yield may be linked to adequate chlorophyll synthesis by crops. Similar results were obtained by Sener et al.<sup>[58]</sup> and Meena et al.<sup>[57]</sup> on maize and popcorn, respectively.

The positive response of popcorn to mineral and organic fertilizers could be linked to the mineral nutrient constituents of the applied fertilizers. The comparable positive influence of 8 t/ha compost or 90 kg N/ha inorganic soil amendments on components of yield, such as the number of ears and harvest index, proposes that either rate was adequate for popcorn cultivation in dryland environments. The compost had superfluous essential mineral nutrients comparable with those supplied by inorganic NPK 20-7-3 fertilizer. It is equally possible that the compost was excellently mineralized, thereby making the minerals available for uptake by the crop for growth and yield. Pérez-Lomas et al.<sup>[60]</sup> and Vargas-García and Suárez-Estrella<sup>[61]</sup> have reported similar effects of sewage co-compost on agricultural fields and rice-wheat cropping systems.

The superior performance of popcorn plants in a well-fertilized field with closer plant stands suggests that when crops are grown in an adequately fertilized field, despite the high population density, leaf formation may not be adversely affected. This is similar to the view of Hamzei and Soltani<sup>[62]</sup> on rapeseed biomass accumulation. Conversely, popcorn plants appeared sensitive to spacing and nutrient availability, considering the fact that the crop grew taller under wide row spacing in an adequately manured field. This may imply that popcorn plants fared better under wide intra-row spacing, perhaps due to minimal intra-plant competition in contrast to closer spacing.

The leaf area index was affected by intra-row spacing and soil amendments, particularly at wider intra-row

spacing and maximum fertilizer application. Wider row spacing under well-fertilized conditions had been shown to improve canopy formation and photosynthate production at the maize silking stage, according to Liu et al.<sup>[63]</sup>. The implication of the results is that popcorn responds to fertilizer application and spacing differently at each phenological phase.

Chlorophyll content of popcorn was affected by intra-spacing and fertilizer application, but this varied across the phenological phases. This shows that popcorn will synthesize chlorophyll content better under a wider intra-row spacing that is adequately fertilized. The same trend recorded for chlorophyll content was observed with the number of ears per plant. Popcorn tends to show better performance with much wider spacing in an adequately fertilized environment than with a narrower spacing, irrespective of fertilizer type, with regards to photoassimilate partitioning<sup>[24]</sup>.

The interaction between intra-row spacing and fertilizer treatments influenced components of yield and the grain yield of popcorn significantly. This suggests that at much wider intra-spacing, popcorn requires minimal fertilizer to produce bigger grains. This implies that better partitioning of photosynthate into economic benefits may also be achieved with a lower fertilizer rate, but certainly not at wider intra-spacing. Hence, the need to equilibrate plant density with soil fertility management cannot be overemphasized. The observed results at the 90 kg N/ha rate and 20 cm × 20 cm plant spacing indicate a better combination that could promote popcorn yield in the region. The greater harvest index of close to a unit indicated that a higher percent of the biomass synthesized was converted to economic value, which makes this combination more appropriate for popcorn cultivation in semi-arid regions. It emphasizes that the crop may have used the applied amendments efficiently under medium intra-row competition to form economic yield. Sakariyawo et al.<sup>[64]</sup> have reported a high harvest index and nitrogen use efficiency in *Zea mays* fertilized with CaC<sub>2</sub> and NPK fertilizers in derived savannah ecology.

## 5. Conclusion

The performance of popcorn improved tremendously with the application of compost or NPK fertilizer. The growth and kernel yield response of the crop to different intra-row spacing varied across the vegetative, tasseling, and maturity stages. The

damaging effect of poor within-row plant configuration may go unnoticed at the early stage of popcorn phenology, as shown in this study. It became clearer at the physiological maturity phase that intra-row spacing affected the performance of popcorn adversely. Yield and its components improved with the application of 8 t/ha of compost supplied to plots intra-spaced at 20 cm × 20 cm. In the same vein, the application of 90 kg N/ha using NPK 20-7-3 fertilizer enhanced the vegetative and reproductive development of popcorn sown at 20 cm × 20 cm intra-row spacing in the dryland of South Africa. Nevertheless, since organic fertilizer could supply balanced nutrients and promote an eco-friendly agroecosystem, applying compost is more preferred to mineral fertilizers for promoting increased popcorn production.

## Statements and Declarations

**Authors' contributions:** Oyeyemi A. Dada conceived the research idea, prepared the proposal, collected and analysed data, and wrote the manuscript. Funso R. Kutu hosted and supervised the activities of O. A. Dada, while Sydney Mavengahama assisted in field experimentation, design, and data analysis. All authors contributed equally and approved the final manuscript.

**Funding:** There was no funding for this research.

**Data Availability:** The data sets used and/or analyzed during the current study are available from the corresponding author on request.

**Acknowledgments:** The technical support provided by the field staff of the Crop Science Department at North-West University is gratefully acknowledged. Equally, the support provided to the first author by the Food Security and Safety Research Niche of the Mafikeng Campus, North-West University, is well appreciated. The South African National Research Foundation (NRF) provided financial support for the Compost project (NRF UID: 90762), to which the study is linked.

**Conflicts of Interest:** We complied with standard ethical practices while conducting the trial. In the same vein, the authors declare that there is no conflict of interest relating to this study.

## References

- <sup>^</sup>Hansen R, Brester G (2009). *Popcorn profile*. *Agricultural Marketing Resource Center*. <http://www.agmr>

- c. org/commodities-products/grains-oilseeds/corn-grain/popcorn-profile/(accessed day 1 Feb. 2016).
2. <sup>a</sup>, <sup>b</sup>Nguyen V, Cooper L, Lowndes J, Melanson K, Angelopoulos TJ, Rippe JM, Reimers K (2012). Popcorn is more satiating than potato chips in normal-weight adults. *Nutrition Journal* 11, 1-6.
  3. <sup>Δ</sup>Ranum P, Peña-Rosas JP, Garcia-Casal MN (2014). Global maize production, utilization, and consumption. *Annals of the new York academy of sciences* 1312, 105-112.
  4. <sup>Δ</sup>Serna-Saldivar SO, Carrillo EP (2019). Food uses of whole corn and dry-milled fractions. In "Corn", pp. 435-467. Elsevier.
  5. <sup>Δ</sup>Fisheries DAaF (2011). Maize market value chain profile 2010-2011. DAFF Pretoria.
  6. <sup>Δ</sup>Kumar SB, Prabhasankar P (2014). Low glycemic index ingredients and modified starches in wheat based food processing: A review. *Trends in Food Science & Technology*. 35: 32-41.
  7. <sup>Δ</sup>Rebello CJ, Greenway FL, Finley JW (2014). Whole grains and pulses: A comparison of the nutritional and health benefits. *Journal of agricultural and food chemistry* 62, 7029-7049.
  8. <sup>Δ</sup>Grandjean AC, Fulgoni III VL, Reimers KJ, Agarwal S (2008). Popcorn consumption and dietary and physiological parameters of US children and adults: analysis of the National Health and Nutrition Examination Survey (NHANES) 1999-2002 dietary survey data. *Journal of the American Dietetic Association*. 108: 853-856.
  9. <sup>Δ</sup>Park D, Allen KG, Stermitz FR, Maga JA (2000). Chemical composition and physical characteristics of unpopped popcorn hybrids. *Journal of Food Composition and Analysis* 13, 921-934.
  10. <sup>Δ</sup>Shahidi F (2015). "Handbook of antioxidants for food preservation," Woodhead Publishing.
  11. <sup>Δ</sup>Rayes P (2010). "Fiber: Bulk of life." Oklahoma Cooperative Extension Service.
  12. <sup>Δ</sup>Júnior SdPF, do Amaral Júnior AT, Rangel RM, Viana AP (2009). Genetic gains in popcorn by full-sib recurrent selection. *Crop Breeding and Applied Biotechnology*. 9.
  13. <sup>a</sup>, <sup>b</sup>Jele P, Derera J, Siwela M (2014). Assessment of popping ability of new tropical popcorn hybrids. *Australian Journal of Crop Science*. 8: 831-839.
  14. <sup>Δ</sup>Du Plessis J (2003). Maize Production; Department of Agriculture: Pretoria, South Africa, 2003.
  15. <sup>Δ</sup>Pardey PG, Greyling JC (2019). Measuring maize in South Africa: The shifting structure of production during the twentieth century, 1904-2015. *Agrekon* 58, 21-41.
  16. <sup>a</sup>, <sup>b</sup>Badu-Apraku B, Fakorede M (2017). Maize in Sub-Saharan Africa: importance and production constraints. In "Advances in genetic enhancement of early and extra-early maize for Sub-Saharan Africa", pp. 3-10. Springer.
  17. <sup>Δ</sup>Tokatlidis IS (2017). Crop adaptation to density to optimize grain yield: breeding implications. *Euphytica* 213, 1-25.
  18. <sup>Δ</sup>Wu W, Ma B (2015). "Integrated nutrient management (INM) for sustaining crop productivity and reducing environmental impact: A review". *Science of the Total Environment*. 512: 415-427.
  19. <sup>Δ</sup>Singh B, Ryan J (2015). Managing fertilizers to enhance soil health. *International Fertilizer Industry Association, Paris, France* 1.
  20. <sup>Δ</sup>Powelson D, Jenkinson D, Johnston A, Poulton P, Gledinning M, Goulding K, Mulvaney R, Khan S, Ellsworth T (2010). Comments on "Synthetic Nitrogen Fertilizers Deplete Soil Nitrogen: A Global Dilemma for Sustainable Cereal Production," by RL Mulvaney, SA Khan, and TR Ellsworth in the *Journal of Environmental Quality* 2009 38: 2295-2314/Reply to Comments on "Synthetic Nitrogen Fertilizers Deplete Soil Nitrogen: A Global Dilemma for Sustainable Cereal Production," by RL Mulvaney, SA Khan, and TR Ellsworth in the *Journal of Environmental Quality* 2009 38: 2295-2314. *Journal of environmental quality* 39, 749.
  21. <sup>Δ</sup>Ghorbani R, Koocheki A, Jahan M, Asadi GA (2008). Impact of organic amendments and compost extracts on tomato production and storability in agroecological systems. *Agronomy for sustainable Development*. 28: 307-311.
  22. <sup>Δ</sup>Matallana Gonzalez M, Martinez-Tome M, Torija Isasa M (2010). Nitrate and nitrite content in organically cultivated vegetables. *Food Additives and Contaminants: Part B* 3, 19-29.
  23. <sup>Δ</sup>Zhao J, Ni T, Li J, Lu Q, Fang Z, Huang Q, Zhang R, Li R, Shen B, Shen Q (2016). "Effects of organic-inorganic compound fertilizer with reduced chemical fertilizer application on crop yields, soil biological activity and bacterial community structure in a rice-wheat cropping system". *Applied Soil Ecology*. 99: 1-12.
  24. <sup>a</sup>, <sup>b</sup>Cihangir H, Oktem A (2019). The effect of different organic nutrients on some quality properties of popcorn (*Zea mays* L. everta). *Asian Food Science Journal*. 7: 1-9.
  25. <sup>Δ</sup>Waldrip HM, He Z, Erich MS (2011). Effects of poultry manure amendment on phosphorus uptake by rye grass, soil phosphorus fractions and phosphatase activity. *Biology and Fertility of Soils* 47, 407-418.



26. <sup>^</sup>Heffer P, Gruère A, Roberts T (2009). Assessment of fertilizer use by crop at the global level. International Fertilizer Industry Association, Paris, [www.fertilizer.org/ifa/Home-Page/LIBRARY/Publication-database.html/Assessment-of-Fertilizer-Use-by-Crop-at-the-Global-Level-2006-07-2007-08.html2](http://www.fertilizer.org/ifa/Home-Page/LIBRARY/Publication-database.html/Assessment-of-Fertilizer-Use-by-Crop-at-the-Global-Level-2006-07-2007-08.html2).
27. <sup>^</sup>Jude JOO, Vidah NM (2008). An assessment of the use of mineral and organic fertilizers by smallholder farmers in Vhembe district, Limpopo province, South Africa. *African Journal of Agricultural Research*. 3: 357-362.
28. <sup>^</sup>Smaling E, Toure M, Ridder N, Sanginga N, Breman H (2006). Fertilizer use and the environment in Africa: friends or foes. Background paper: African fertilizer summit: nourish the soil, feed the continent 9, 13.
29. <sup>^</sup>Sangoi L (2001). Understanding plant density effects on maize growth and development: an important issue to maximize grain yield. *Ciência rural* 31, 159-168.
30. <sup>^</sup>Brodrick R, Bange M, Milroy S, Hammer G (2013). Physiological determinants of high yielding ultra-narrow row cotton: Canopy development and radiation use efficiency. *Field Crops Research*. 148: 86-94.
31. <sup>^</sup>Dong H, Kong X, Li W, Tang W, Zhang D (2010). Effects of plant density and nitrogen and potassium fertilization on cotton yield and uptake of major nutrients in two fields with varying fertility. *Field Crops Research*. 119: 106-113.
32. <sup>^</sup>Dai J, Li W, Tang W, Zhang D, Li Z, Lu H, Eneji AE, Dong H (2015). Manipulation of dry matter accumulation and partitioning with plant density in relation to yield stability of cotton under intensive management. *Field Crops Research*. 180: 207-215.
33. <sup>^</sup>Dahlman C (2007). Technology, globalization, and international competitiveness: Challenges for developing countries. *Industrial development for the 21st century: Sustainable development perspectives*, 29-83.
34. <sup>^</sup>Lambin EF, Meyfroidt P (2011). Global land use change, economic globalization, and the looming land scarcity. *Proceedings of the National Academy of Sciences*. 108: 3465-3472.
35. <sup>^</sup>Koralage I, Silva N, De Silva C. (2015). The determination of available phosphorus in soil: a quick and simple method. *Open Univ Sri Lanka J*. 8:1-17.
36. <sup>^</sup>Beretta AN, Silbermann AV, Paladino L, Torres D, Kassahun D, Musselli R, Lamohte AG (2014). Soil texture analyses using a hydrometer: modification of the Bouyoucos method. *Ciencia e investigación agraria: revista latinoamericana de ciencias de la agricultura*. 41: 263-271.
37. <sup>^</sup>Karak T, Abollino O, Paul RK, Dutta AK, Giacomino A, Khare P, Boruah RK (2018). Achievability of municipal solid waste compost for tea cultivation with special reference to cadmium. *CLEAN-Soil, Air, Water*. 46: 1800093.
38. <sup>^</sup>Emmet-Booth J, Forristal D, Fenton O, Bondi G, Creamer R, Holden N (2017). Impacts of soil moisture content on visual soil evaluation. In "EGU General Assembly Conference Abstracts", pp. 8129.
39. <sup>^</sup>Fageria NK, Baligar VC, Clark R (2006). "Physiology of crop production," CRC Press.
40. <sup>^</sup>Berdjour A, Dugje I, Rahman NA, Odoom DA, Karama A, Ajala S (2020). Direct estimation of maize leaf area index as influenced by organic and inorganic fertilizer rates in Guinea savanna.
41. <sup>a, b</sup> Abebe Z, Feyisa H (2017). Effects of nitrogen rates and time of application on yield of maize: rainfall variability influenced time of N application. *International Journal of Agronomy*. 2017.
42. <sup>^</sup>Lee C (2007). Corn growth and development. Melal ui [http://www.uky.edu/ag/grain\\_crops](http://www.uky.edu/ag/grain_crops).
43. <sup>a, b</sup> Gomez KA, Gomez AA (1984). "Statistical procedures for agricultural research," John Wiley & sons.
44. <sup>^</sup>O'Rourke N, Hatcher L (2013). "A step-by-step approach to using SAS for factor analysis and structural equation modeling," Sas Institute.
45. <sup>a, b</sup> Bernhard BJ, Below FE (2020). Plant population and row spacing effects on corn: Plant growth, phenology, and grain yield. *Agronomy Journal*. 112: 2456-2465.
46. <sup>a, b</sup> Moosavi S, Seghatoleslami M, Moazeni A (2012). Effect of planting date and plant density on morphological traits, LAI and forage corn (Sc. 370) yield in second cultivation. *International Research Journal of Applied and Basic Sciences* 3, 57-63.
47. <sup>a, b</sup> Widdicombe WD, Thelen KD (2002). Row width and plant density effects on corn grain production in the northern Corn Belt. *Agronomy journal* 94, 1020-1023.
48. <sup>^</sup>Murányi E, Pepo P (2013). The effects of plant density and row spacing on the height of maize hybrids of different vegetation time and genotype. *International Journal of Agricultural and Biosystems Engineering* 7, 1011-1014.
49. <sup>^</sup>Williams II MM (2012). "Agronomics and economics of plant population density on processing sweet corn". *Field Crops Research*. 128: 55-61.
50. <sup>^</sup>Board J, Harville B (1992). Explanations for greater light interception in narrow-vs. wide-row. *Crop science*

- ce. 32: 198–202.
51. <sup>△</sup>Liu T, Song F, Liu S, Zhu X (2012). Light interception and radiation use efficiency response to narrow-wide row planting patterns in maize. *Australian Journal of Crop Science* 6, 506–513.
  52. <sup>△</sup>Golada S, Sharma G, Nepalia V (2017). Effect of spacing, nitrogen and growth regulators on yield and nutrient uptake by baby corn (*Zea mays* L.). *Annals of Plant and Soil Research*. 19: 148–153.
  53. <sup>△</sup>Sheth S, Desai K, Patil S, Navya K, Chaudhari V (2017). Effect of integrated nutrient management on growth, yield and quality of sweet potato [*Ipomoea batatas*(L.) Lam]. *International Journal of Chemical Studies* 5, 346–349.
  54. <sup>△</sup>Laxminarayana K, John KS, Ravindran C, Naskar S (2011). Effect of lime, inorganic, and organic sources on soil fertility, yield, quality, and nutrient uptake of sweet potato in alfisols. *Communications in soil science and plant analysis* 42, 2515–2525.
  55. <sup>△</sup>Dada OA, Imade F, Anifowose E (2017). Growth and proximate composition of *Amaranthus cruentus* L. on poor soil amended with compost and arbuscular mycorrhiza fungi. *International Journal of Recycling of Organic Waste in Agriculture*. 6: 195–202.
  56. <sup>△</sup>White G, Ross I, Poneleit C (1981). Fully-exposed drying of popcorn. *Transactions of the ASAE* 24, 466–468.
  57. <sup>△</sup><sup>a, b</sup>Meena B, Kumar A, Shiva Dhar PS, Kumar A (2016). Productivity, nutrient uptake and quality of popcorn and potato in relation to organic nutrient management practices. *Annals of Agricultural Research New Series* 37, 72–79.
  58. <sup>△</sup><sup>a, b</sup>Sener O, Gozubenli H, Konuskan O, Kilinc M (2004). The effects of intra-row spacings on the grain yield and some agronomic characteristics of maize (*Zea mays* L.) hybrids. *Asian Journal of Plant Sciences*.
  59. <sup>△</sup>Gözübenli H (2010). Nitrogen dose and plant density effects on popcorn grain yield. *African Journal of Biotechnology*. 9: 3828–3832.
  60. <sup>△</sup>Pérez-Lomas AL, Delgado G, Párraga J, Delgado R, Almendros G, Aranda V (2010). Evolution of organic matter fractions after application of co-compost of sewage sludge with pruning waste to four Mediterranean agricultural soils. A soil microcosm experiment. *Waste Management* 30, 1957–1965.
  61. <sup>△</sup>Vargas-García M, Suárez-Estrella F (2008). Effect of compost on biological properties of the soil. *Composting*. (Moreno-Cascos J, & Moral-Herrero J, eds), Mundi-Prensa, Madrid, 329–350.
  62. <sup>△</sup>Hamzei J, Soltani J (2012). Deficit irrigation of rape seed for water-saving: Effects on biomass accumulation, light interception and radiation use efficiency under different N rates. *Agriculture, ecosystems & environment*. 155: 153–160.
  63. <sup>△</sup>Liu T, Song F, Liu S, Zhu X (2011). Canopy structure, light interception, and photosynthetic characteristics under different narrow-wide planting patterns in maize at silking stage. *Spanish Journal of Agricultural Research* 9, 1249–1261.
  64. <sup>△</sup>Sakariyawo O, Oyeledun K, Adeyemi N, Atayese M (2020). Nitrogen use efficiency and performance of maize (*Zea mays* L.) cultivars as influenced by calcium carbide and inorganic nitrogen application rates in a derived savanna. *Journal of Plant Nutrition* 43, 784–797.

## Declarations

**Funding:** No specific funding was received for this work.

**Potential competing interests:** No potential competing interests to declare.